

**BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid
bandage liquid
AmerisourceBergen Drug Corporation**

GNP Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

- first aid to prevent bacteria contamination in minor cuts and scrapes

Warnings

For external use only.

Flammable:

do not use while smoking or near heat or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor
- on infected areas or wounds that are draining
- with other first aid products such as lotions and creams
- over sutures
- on mucous membranes

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious bleeding
- diabetes
- poor circulation
- serious burns

When using this product

- do not store at temperature above 120 °F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

Stop use and ask a doctor if

- condition persists or gets worse
- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- Clean affected area
- apply a small amount on the area 1-3 times daily
- let dry
- a second coating may be applied for extra protection
- to remove, apply more Liquid Bandage and quickly wipe off
- finger nail polish remover may dissolve Liquid Bandage

Other information

- ***cap bottle tightly***
- store at room temperature away from heat
- may temporarily sting upon application
- do not allow to come in contact with floors, countertops, or other finished surfaces - will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel**Good Neighbor Pharmacy****For small cuts & wounds****Liquid Bandage**

Antiseptic to Help Prevent Infection

- Invisible
- Flexible

- Easy to use

Use liquid bandage for:

blisters, chapped & cracked fingertips, hangnails, paper cuts, shaving nicks & help in preventing the formulation of calluses.

Used by:

athletes, active lifestyles, outdoor sportsman & musicians

NET 0.3 FL OZ (9 mL)



NDC 46122-525-23

For small cuts & wounds

Liquid Bandage

Antiseptic to Help Prevent Infection

- Invisible
- Flexible
- Easy to use

NET 0.3 FL OZ (9mL)



Use liquid bandage for:

blisters, chapped & cracked finger tips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

Used by:

athletes, active lifestyles, outdoor sportsman & musicians.

53-033GN-01

For small cuts & wounds

Liquid Bandage

Antiseptic to Help Prevent Infection

For small cuts & wounds

Liquid Bandage

Antiseptic to Help Prevent Infection

NET 0.3 FL OZ (9mL)



NO VARNISH AREA



Distributed By AmerisourceBergen
 1300 Morris Drive, Chesterbrook, PA 19087
 Visit us at www.goodneighborpharmacy.com
 Made in USA with U.S. and imported parts

ABC# 364-998



Questions? Call 1-866-964-0939

Inactive ingredients acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Liquid Bandage and quickly wipe off

Other information ■ cap bottle tightly ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply more

Directions ■ for adults and children 2 years of age and older ■ clean affected area ■ apply a small amount on the area 1 to 3 times daily and let dry ■ a second coating may be applied for extra protection ■ children under 2 years of age: consult a doctor

Warnings ■ avoid contact with eyes ■ avoid contact with eyes ■ animal bites ■ serious burns

Uses ■ first aid to prevent bacterial contamination in minor cuts, scrapes and burns ■ for the temporary relief of pain

Drug Facts

Active ingredients Benzethonium chloride 0.2% ■ first aid antiseptic Dyclonine hydrochloride 0.75% ■ topical analgesic

Purpose

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

■ symptoms persist for more than 7 days or clear up and occur again within a few days

Stop use and ask a doctor if ■ condition persists or gets worse ■ infection occurs

When using this product ■ avoid contact with eyes ■ animal bites ■ serious burns

Ask a doctor before use if you have ■ deep or puncture wounds ■ deep cuts ■ longer than 1 week unless directed by a doctor

Do not use ■ in the eyes ■ over large areas of the body

For external use only. Flammable. Keep away from fire or flame.

DO NOT USE IF NECKBAND IS BROKEN



BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE

liquid bandage liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-525
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

Inactive Ingredients

Ingredient Name	Strength
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y219G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	
ACETONE (UNII: 1364PS73AF)	

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:46122-525-23	9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product	12/21/2017	
Marketing Information				
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	OTC Monograph Drug	M003	12/21/2017	

Labeler - AmerisourceBergen Drug Corporation (007914906)

Revised: 11/2024

AmerisourceBergen Drug Corporation