

**MUCUS RELIEF SEVERE COLD DAYTIME MAXIMUM STRENGTH- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid
TARGET Corporation**

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Antihistamine/Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm(mucus)

Ask a doctor or pharmacist before use if you are

are taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition,

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
- store between 20-25°C (68-77°). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue 1, FD&C red 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call **1-800-910-6874**

Principal Display Panel

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® DayTime Severe Cold*

Maximum Strength

daytime severe cold

Acetaminophen

(pain reliever / fever reducer)

dextromethorphan HBr (cough Suppressant)

guaifenesin (expectorant)

phenylephrine HCl (nasal decongestant)

relieves aches, fever and sore throat controls cough

relieves nasal and chest congestion thins and loosens mucus

AGES 12 + YEARS

FL OZ (mL)

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® DayTime Severe Cold.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

Dist. by Target Corp.

Minneapolis, MN 55403

Package Label

NDC 11673-410-06

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Day Time Severe Cold*

maximum strength
daytime
severe cold

acetaminophen

(pain reliever/fever reducer)
dextromethorphan HBr (cough suppressant)
guaifenesin (expectorant)
phenylephrine HCl (nasal decongestant)

relieves aches, fever and sore throat
controls cough
relieves nasal and chest congestion
thins and loosens mucus



AGES
12+
YEARS

6 FL OZ (177 mL)

PLD-B346B LB004227

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Day Time Severe Cold.



094 14 0185 R01
C-000274-01-049
Dist. by Target Corp.
Minneapolis, MN 55403
TM & ©2021 Target Brands, Inc.

PLD-B346B
LB004228

PEEL CORNER FOR DRUG FACTS ▲

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe ▶

Drug Facts (continued)

skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ▶

PEEL CORNER FOR MORE DRUG FACTS ▲

Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. ▶

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-800-910-6874

TARGET Mucus Relief Severe Cold

MUCUS RELIEF SEVERE COLD DAYTIME MAXIMUM STRENGTH

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-410
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-410-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/31/2016	

