

FOSAMPRENAVIR CALCIUM- fosamprenavir calcium tablet, coated
Sun Pharmaceutical Industries, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use FOSAMPRENAVIR CALCIUM TABLETS safely and effectively. See full prescribing information for FOSAMPRENAVIR CALCIUM TABLETS.

FOSAMPRENAVIR CALCIUM tablets, for oral use

Initial U.S. Approval: 2003

----- **RECENT MAJOR CHANGES** -----

Dosage and Administration, Adults (2.2) 03/2019

Contraindications (4) 03/2019

----- **INDICATIONS AND USAGE** -----

Fosamprenavir calcium tablet are HIV protease inhibitor indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection. (1) (1)

----- **DOSAGE AND ADMINISTRATION** -----

- Therapy-Naive Adults: Fosamprenavir calcium tablets 1,400 mg twice daily; fosamprenavir calcium tablets 1,400 mg once daily plus ritonavir 200 mg once daily; fosamprenavir calcium tablets 1,400 mg once daily plus ritonavir 100 mg once daily; fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily. (2.2)
- Protease Inhibitor-Experienced Adults: Fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily. (2.2)
- Pregnant Patients: Fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily should only be considered in women who are already on a stable twice-daily regimen of fosamprenavir calcium /ritonavir 700 mg/100 mg prior to pregnancy and who are virologically suppressed (HIV-1 RNA less than 50 copies per mL). (2.2)
- Pediatric Patients (aged at least 4 weeks to 18 years): Dosage should be calculated based on body weight (kg) and should not exceed adult dose. (2.3)
- Hepatic Impairment: Recommended adjustments for patients with mild, moderate, or severe hepatic impairment. (2.4)

Dosing Considerations (2)

- Fosamprenavir calcium tablets may be taken with or without food. (2.1)
- Fosamprenavir Calcium Oral Suspension: Adults should take without food; pediatric patients should take with food. (2.1)

----- **DOSAGE FORMS AND STRENGTHS** -----

700 mg tablets (3) (3)

----- **CONTRAINDICATIONS** -----

- Hypersensitivity to fosamprenavir calcium or amprenavir (e.g., Stevens-Johnson syndrome). (4)
- Drugs highly dependent on cytochrome P450 (CYP)3A4 for clearance and for which elevated plasma levels may result in serious and/or life-threatening events. (4)
- Review ritonavir contraindications when used in combination. (4)

----- **WARNINGS AND PRECAUTIONS** -----

- The concomitant use of fosamprenavir calcium with ritonavir and certain other drugs may result in known or potentially significant drug interactions. Consult the full prescribing information prior to and during treatment for potential drug interactions. (5.1, 7.3)
- Fosamprenavir calcium should be discontinued for severe skin reactions, including Stevens-Johnson syndrome. (5.2)
- Fosamprenavir calcium should be used with caution in patients with a known sulfonamide allergy. (5.3)
- Use of higher-than-approved doses may lead to transaminase elevations. Patients with hepatitis B or C are at increased risk of transaminase elevations. (5.4)
- Patients receiving fosamprenavir calcium may develop new onset or exacerbations of diabetes mellitus, hyperglycemia (5.5), immune reconstitution syndrome (5.6), increase of body fat (5.7), and elevated triglyceride and cholesterol concentrations (5.8). Monitor cholesterol and triglycerides prior to therapy and periodically thereafter.

- Acute hemolytic anemia has been reported with amprenavir. (5.9)
- Hemophilia: Spontaneous bleeding may occur, and additional factor VIII may be required. (5.10)
- Nephrolithiasis: Cases of nephrolithiasis have been reported with fosamprenavir. (5.11)

----- **ADVERSE REACTIONS** -----

- In adults the most common adverse reactions (incidence greater than or equal to 4%) are diarrhea, rash, nausea, vomiting, and headache. (6.1)
- Vomiting and neutropenia were more frequent in pediatrics than in adults. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Sun Pharmaceutical Industries, Inc. at 1-800-406-7984 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. (6)

----- **DRUG INTERACTIONS** -----

- Coadministration of fosamprenavir calcium with drugs that induce CYP3A4 may decrease amprenavir (active metabolite) concentrations leading to potential loss of virologic activity. (7, 12.3)
- Coadministration with drugs that inhibit CYP3A4 may increase amprenavir concentrations. (7, 12.3)
- Coadministration of fosamprenavir calcium or fosamprenavir calcium and ritonavir may result in clinically significant interactions with drugs metabolized by CYP3A4. (7)
- Coadministration of fosamprenavir calcium and ritonavir may result in clinically significant interactions with drugs metabolized by CYP2D6. (7)

----- **USE IN SPECIFIC POPULATIONS** -----

- Lactation: Breastfeeding is not recommended due to potential for HIV transmission. (8.2)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 12/2019

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Information

2.2 Adults

2.3 Pediatric Patients (Aged at Least 4 Weeks to 18 Years)

2.4 Patients with Hepatic Impairment

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Serious Adverse Reactions Due to Drug Interactions

5.2 Skin Reactions

5.3 Sulfa Allergy

5.4 Hepatic Toxicity

5.5 Diabetes/Hyperglycemia

5.6 Immune Reconstitution Syndrome

5.7 Increase in Body Fat

5.8 Lipid Elevations

5.9 Hemolytic Anemia

5.10 Patients with Hemophilia

5.11 Nephrolithiasis

5.12 Resistance/Cross-Resistance

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Cytochrome P450 Inhibitors and Inducers

7.2 Established and Other Potentially Significant Drug Interactions

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

14.1 Therapy-Naive Adult Trials

14.2 Protease Inhibitor-Experienced Adult Trials

14.3 Pediatric Trials

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Fosamprenavir calcium tablets are indicated in combination with other antiretroviral agents for the treatment of human immunodeficiency virus (HIV-1) infection.

The following points should be considered when initiating therapy with fosamprenavir calcium tablets plus ritonavir in protease inhibitor-experienced patients:

- The protease inhibitor-experienced patient trial was not large enough to reach a definitive conclusion that fosamprenavir calcium tablets plus ritonavir and lopinavir plus ritonavir are clinically equivalent [see *Clinical Studies (14.2)*].
- Once-daily administration of fosamprenavir calcium tablets plus ritonavir is not recommended for adult protease inhibitor-experienced patients or any pediatric patients [see *Dosage and Administration (2.2,2.3), Clinical Studies (14.2, 14.3)*].
- Dosing of fosamprenavir calcium tablets plus ritonavir is not recommended for protease inhibitor-experienced pediatric patients younger than 6 months [see *Clinical Pharmacology (12.3)*].

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Information

Fosamprenavir calcium tablets may be taken with or without food.

Adults should take fosamprenavir calcium oral suspension without food. Pediatric patients should take fosamprenavir calcium oral suspension with food [see *Clinical Pharmacology (12.3)*]. If emesis occurs within 30 minutes after dosing, re-dosing of fosamprenavir calcium oral suspension should occur.

Higher-than-approved dose combinations of fosamprenavir calcium tablets plus ritonavir are not recommended due to an increased risk of transaminase elevations [see *Overdosage (10)*].

When fosamprenavir calcium tablets are used in combination with ritonavir, prescribers should consult the full prescribing information for ritonavir.

2.2 Adults

Therapy-Naive Adults

- Fosamprenavir calcium tablets 1,400 mg twice daily (without ritonavir).
- Fosamprenavir calcium tablets 1,400 mg once daily plus ritonavir 200 mg once daily.
- Fosamprenavir calcium tablets 1,400 mg once daily plus ritonavir 100 mg once daily.
- Dosing of fosamprenavir calcium tablets 1,400 mg once daily plus ritonavir 100 mg once daily is supported by pharmacokinetic data [see *Clinical Pharmacology (12.3)*].
- Fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily.
- Dosing of fosamprenavir calcium tablets 700 mg twice daily plus 100 mg ritonavir twice daily is supported by pharmacokinetic and safety data [see *Clinical Pharmacology (12.3)*].

Protease Inhibitor-Experienced Adults

- Fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily.

Pregnancy

- Fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily.
- Dosing of fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily should only be considered in pregnant patients who are already on a stable twice-daily regimen of fosamprenavir calcium/ritonavir 700 mg/100 mg prior to pregnancy and who are virologically suppressed (HIV-1 RNA less than 50 copies per mL). Lower exposures of amprenavir were observed during pregnancy; therefore, viral load should be monitored closely to ensure viral suppression is maintained [see *Use in Specific Populations (8.1), Clinical Pharmacology (12.3)*]. Data regarding use of other regimens of fosamprenavir calcium tablets (with or without ritonavir) in pregnancy are not available.

2.3 Pediatric Patients (Aged at Least 4 Weeks to 18 Years)

The recommended dosage of fosamprenavir calcium in patients aged at least 4 weeks to 18 years should be calculated based on body weight (kg) and should not exceed the recommended adult dose (Table 1).

Table 1. Twice-Daily Dosage Regimens by Weight for Protease Inhibitor-Naive Pediatric Patients (Aged 4 Weeks and Older) and for Protease Inhibitor-Experienced Pediatric Patients (Aged 6 Months and Older) Using Fosamprenavir Calcium Oral Suspension with Concurrent Ritonavir

Weight	Twice-Daily Dosage Regimen
< 11 kg	Fosamprenavir Calcium 45 mg/kg plus ritonavir 7 mg/kg ^a
11 kg - < 15 kg	Fosamprenavir Calcium 30 mg/kg plus ritonavir 3 mg/kg ^a
15 kg - < 20 kg	Fosamprenavir Calcium 23 mg/kg plus ritonavir 3

	mg/kg ^a
≥ 20 kg	Fosamprenavir Calcium 18 mg/kg plus ritonavir 3 mg/kg ^a

^a When dosing with ritonavir, do not exceed the adult dose of fosamprenavir calcium 700 mg/ritonavir 100 mg twice-daily dose.

Alternatively, protease inhibitor-naïve children aged 2 years and older can be administered fosamprenavir calcium (without ritonavir) 30 mg per kg twice daily.

Fosamprenavir calcium should only be administered to infants born at 38 weeks' gestation or greater and who have attained a postnatal age of 28 days.

For pediatric patients, pharmacokinetic and clinical data:

- do not support once-daily dosing of fosamprenavir calcium alone or in combination with ritonavir [see *Clinical Studies (14.3)*].
- do not support administration of fosamprenavir calcium alone or in combination with ritonavir for protease inhibitor-experienced children younger than 6 months [see *Clinical Pharmacology (12.3)*].
- do not support twice-daily dosing of fosamprenavir calcium without ritonavir in pediatric patients younger than 2 years [see *Clinical Pharmacology (12.3)*].

Other Dosing Considerations

- When administered without ritonavir, the adult regimen of fosamprenavir calcium tablets 1,400 mg twice daily may be used for pediatric patients weighing at least 47 kg.

When administered in combination with ritonavir, fosamprenavir calcium tablets may be used for pediatric patients weighing at least 39 kg; ritonavir capsules may be used for pediatric patients weighing at least 33 kg.

2.4 Patients with Hepatic Impairment

See *Clinical Pharmacology (12.3)*.

Mild Hepatic Impairment (Child-Pugh Score Ranging from 5 to 6)

Fosamprenavir calcium tablets should be used with caution at a reduced dosage of 700 mg twice daily without ritonavir (therapy-naïve) or 700 mg twice daily plus ritonavir 100 mg once daily (therapy-naïve or protease inhibitor-experienced).

Moderate Hepatic Impairment (Child-Pugh Score Ranging from 7 to 9)

Fosamprenavir calcium tablets should be used with caution at a reduced dosage of 700 mg twice daily without ritonavir (therapy-naïve) or 450 mg twice daily plus ritonavir 100 mg once daily (therapy-naïve or protease inhibitor-experienced).

Severe Hepatic Impairment (Child-Pugh Score Ranging from 10 to 15)

Fosamprenavir calcium tablets should be used with caution at a reduced dosage of 350 mg twice daily without ritonavir (therapy-naïve) or 300 mg twice daily plus ritonavir 100 mg once daily (therapy-naïve or protease inhibitor-experienced).

There are no data to support dosing recommendations for pediatric patients with hepatic impairment.

3 DOSAGE FORMS AND STRENGTHS

Fosamprenavir calcium tablets, 700 mg, are pink colored, coated, oval-shaped tablets with “**RJ47**” imprinted on one side with black ink and plain on the other side.

4 CONTRAINDICATIONS

- Fosamprenavir calcium is contraindicated in patients with previously demonstrated clinically significant hypersensitivity (e.g., Stevens-Johnson syndrome) to any of the components of this product or to amprenavir.
- Fosamprenavir calcium is contraindicated when coadministered with drugs that are highly dependent on cytochrome P450 (CYP)3A4 for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events. These drugs and other contraindicated drugs (which may lead to reduced efficacy of fosamprenavir calcium and possible resistance) are listed below [see *Drug Interactions (7)*, *Clinical Pharmacology (12.3)*]. The list of contraindicated drugs applies to the use of fosamprenavir calcium with or without ritonavir, unless otherwise indicated. If fosamprenavir calcium is coadministered with ritonavir, reference should be made to the full prescribing information for ritonavir for additional contraindications.
- Fosamprenavir calcium is contraindicated when coadministered with the following drugs:
 - Alpha 1-adrenoreceptor antagonist: Alfuzosin
 - Antiarrhythmics: Flecainide (with **ritonavir**), propafenone (with **ritonavir**)
 - Antimycobacterial: Rifampin
 - Antipsychotic: Lurasidone (with **ritonavir**), pimozide
 - Ergot derivatives: Dihydroergotamine, ergonovine, ergotamine, methylergonovine
 - GI motility agent: Cisapride
 - Herbal product: St. John's wort (*Hypericum perforatum*)
 - Lipid modifying agents: Lomitapide, lovastatin, simvastatin
 - Non-nucleoside reverse transcriptase inhibitor: Delavirdine
 - PDE5 inhibitor: Sildenafil (REVATIO) (for treatment of pulmonary arterial hypertension)
 - Sedative/hypnotics: Midazolam, triazolam

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Serious Adverse Reactions Due to Drug Interactions

Initiation of fosamprenavir calcium/ritonavir, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A, or initiation of medications metabolized by CYP3A in patients already receiving fosamprenavir calcium/ritonavir may increase plasma concentrations of medications metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of fosamprenavir calcium/ritonavir, respectively. These interactions may lead to:

- clinically significant adverse reactions, potentially leading to severe, life-threatening, or fatal events from greater exposures of concomitant medications.
- clinically significant adverse reactions from greater exposures of fosamprenavir calcium/ritonavir.
- loss of therapeutic effect of fosamprenavir calcium/ritonavir and possible development of resistance.

See Table 6 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations [see *Drug Interactions (7)*]. Consider the potential for drug interactions prior to and during therapy with fosamprenavir calcium/ritonavir; review concomitant medications during therapy with fosamprenavir calcium/ritonavir, and monitor for the adverse reactions associated with the concomitant medications [see *Contraindications (4)*, *Drug Interactions (7)*].

5.2 Skin Reactions

Severe and life-threatening skin reactions, including 1 case of Stevens-Johnson syndrome among 700 subjects treated with fosamprenavir calcium in clinical trials. Treatment with fosamprenavir calcium should be discontinued for severe or life-threatening rashes and for moderate rashes accompanied by systemic symptoms [see *Adverse Reactions (6)*].

5.3 Sulfa Allergy

Fosamprenavir calcium should be used with caution in patients with a known sulfonamide allergy. Fosamprenavir contains a sulfonamide moiety. The potential for cross-sensitivity between drugs in the sulfonamide class and fosamprenavir is unknown. In a clinical trial of fosamprenavir calcium used as the sole protease inhibitor, rash occurred in 2 of 10 subjects (20%) with a history of sulfonamide allergy compared with 42 of 126 subjects (33%) with no history of sulfonamide allergy. In 2 clinical trials of fosamprenavir calcium plus low-dose ritonavir, rash occurred in 8 of 50 subjects (16%) with a history of sulfonamide allergy compared with 50 of 412 subjects (12%) with no history of sulfonamide allergy.

5.4 Hepatic Toxicity

Use of fosamprenavir calcium with ritonavir at higher-than-recommended dosages may result in transaminase elevations and should not be used [see *Dosage and Administration (2)*, *Overdosage (10)*]. Patients with underlying hepatitis B or C or marked elevations in transaminases prior to treatment may be at increased risk for developing or worsening of transaminase elevations. Appropriate laboratory testing should be conducted prior to initiating therapy with fosamprenavir calcium and patients should be monitored closely during treatment.

5.5 Diabetes/Hyperglycemia

New onset diabetes mellitus, exacerbation of pre-existing diabetes mellitus, and hyperglycemia have been reported during postmarketing surveillance in HIV-1-infected patients receiving protease inhibitor therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued protease inhibitor therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and causal relationships between protease inhibitor therapy and these events have not been established.

5.6 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including fosamprenavir calcium. During the initial phase of combination antiretroviral treatment, patients whose immune systems respond may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, and Guillain-Barré syndrome) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.7 Increase in Body Fat

Increase of body fat has been observed in patients receiving protease inhibitors, including fosamprenavir calcium. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

5.8 Lipid Elevations

Treatment with fosamprenavir calcium plus ritonavir has resulted in increases in the concentration of

triglycerides and cholesterol [see *Adverse Reactions (6)*]. Triglyceride and cholesterol testing should be performed prior to initiating therapy with fosamprenavir calcium and at periodic intervals during therapy. Lipid disorders should be managed as clinically appropriate [see *Drug Interactions (7)*].

5.9 Hemolytic Anemia

Acute hemolytic anemia has been reported in a patient treated with amprenavir.

5.10 Patients with Hemophilia

There have been reports of spontaneous bleeding in patients with hemophilia A and B treated with protease inhibitors. In some patients, additional factor VIII was required. In many of the reported cases, treatment with protease inhibitors was continued or restarted. A causal relationship between protease inhibitor therapy and these episodes has not been established.

5.11 Nephrolithiasis

Cases of nephrolithiasis were reported during postmarketing surveillance in HIV-1-infected patients receiving fosamprenavir calcium. Because these events were reported voluntarily during clinical practice, estimates of frequency cannot be made. If signs or symptoms of nephrolithiasis occur, temporary interruption or discontinuation of therapy may be considered.

5.12 Resistance/Cross-Resistance

Because the potential for HIV cross-resistance among protease inhibitors has not been fully explored, it is unknown what effect therapy with fosamprenavir calcium will have on the activity of subsequently administered protease inhibitors. Fosamprenavir calcium has been studied in patients who have experienced treatment failure with protease inhibitors [see *Clinical Studies (14.2)*].

6 ADVERSE REACTIONS

- Severe or life-threatening skin reactions have been reported with the use of fosamprenavir calcium [see *Warnings and Precautions (5.2)*].
- The most common moderate to severe adverse reactions in clinical trials of fosamprenavir calcium were diarrhea, rash, nausea, vomiting, and headache.
- Treatment discontinuation due to adverse events occurred in 6.4% of subjects receiving fosamprenavir calcium and in 5.9% of subjects receiving comparator treatments. The most common adverse reactions leading to discontinuation of fosamprenavir calcium (incidence less than or equal to 1% of subjects) included diarrhea, nausea, vomiting, AST increased, ALT increased, and rash.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Adult Trials

The data for the 3 active-controlled clinical trials described below reflect exposure of 700 HIV-1-infected subjects to fosamprenavir calcium tablets, including 599 subjects exposed to fosamprenavir calcium for greater than 24 weeks, and 409 subjects exposed for greater than 48 weeks. The population age ranged from 17 to 72 years. Of these subjects, 26% were female, 51% white, 31% black, 16% American Hispanic, and 70% were antiretroviral-naive. Sixty-one percent received fosamprenavir calcium 1,400 mg once daily plus ritonavir 200 mg once daily; 24% received fosamprenavir calcium 1,400 mg twice daily; and 15% received fosamprenavir calcium 700 mg twice daily plus ritonavir 100 mg twice daily.

Selected adverse reactions reported during the clinical efficacy trials of fosamprenavir calcium are shown in Tables 2 and 3. Each table presents adverse reactions of moderate or severe intensity in subjects treated with combination therapy for up to 48 weeks.

Table 2. Selected Moderate/Severe Clinical Adverse Reactions Reported in Greater than or Equal to 2% of Antiretroviral-Naive Adult Subjects

Adverse Reaction	APV30001 ^a		APV30002 ^a	
	Fosamprenavir Calcium 1,400 mg Twice Daily (n = 166)	Nelfinavir 1,250 mg Twice Daily (n = 83)	Fosamprenavir Calcium 1,400 mg and Ritonavir 200 mg Once Daily (n = 322)	Nelfinavir 1,250 mg Twice Daily (n = 327)
Gastrointestinal				
Diarrhea	5%	18%	10%	18%
Nausea	7%	4%	7%	5%
Vomiting	2%	4%	6%	4%
Abdominal pain	1%	0%	2%	2%
Skin				
Rash	8%	2%	3%	2%
General disorders				
Fatigue	2%	1%	4%	2%
Nervous system				
Headache	2%	4%	3%	3%

^a All subjects also received abacavir and lamivudine twice daily.

Table 3. Selected Moderate/Severe Clinical Adverse Reactions Reported in Greater than or Equal to 2% of Protease Inhibitor-Experienced Adult Subjects (Trial APV30003)

Adverse Reaction	Fosamprenavir Calcium 700 mg and Ritonavir 100 mg Twice Daily ^a (n = 106)	Lopinavir 400 mg and Ritonavir 100 mg Twice Daily ^a (n = 103)
Gastrointestinal		
Diarrhea	13%	11%
Nausea	3%	9%
Vomiting	3%	5%
Abdominal pain	< 1%	2%
Skin		
Rash	3%	0%
Nervous system		
Headache	4%	2%

^a All subjects also received 2 reverse transcriptase inhibitors.

Skin rash (without regard to causality) occurred in approximately 19% of subjects treated with fosamprenavir calcium in the pivotal efficacy trials. Rashes were usually maculopapular and of mild or moderate intensity, some with pruritus. Rash had a median onset of 11 days after initiation of fosamprenavir calcium and had a median duration of 13 days. Skin rash led to discontinuation of fosamprenavir calcium in less than 1% of subjects. In some subjects with mild or moderate rash, dosing with fosamprenavir calcium was often continued without interruption; if interrupted, reintroduction of fosamprenavir calcium generally did not result in rash recurrence.

The percentages of subjects with Grade 3 or 4 laboratory abnormalities in the clinical efficacy trials of fosamprenavir calcium are presented in Tables 4 and 5.

Table 4. Grade 3/4 Laboratory Abnormalities Reported in Greater than or Equal to 2% of Antiretroviral-Naive Adult Subjects in Trials APV30001 and APV30002

	APV30001^a		APV30002^a	
	Fosamprenavir	Nelfinavir	Fosamprenavir	Nelfinavir

Laboratory Abnormality	Calcium 1,400 mg Twice Daily (n = 166)	1,250 mg Twice Daily (n = 83)	Calcium 1,400 mg and Ritonavir 200 mg Once Daily (n = 322)	1,250 mg Twice Daily (n = 327)
ALT (> 5 x ULN)	6%	5%	8%	8%
AST (> 5 x ULN)	6%	6%	6%	7%
Serum lipase (> 2 x ULN)	8%	4%	6%	4%
Triglycerides ^b (> 750 mg/dL)	0%	1%	6%	2%
Neutrophil count, absolute (< 750 cells/mm ³)	3%	6%	3%	4%

^a All subjects also received abacavir and lamivudine twice daily.

^b Fasting specimens.

ULN = Upper limit of normal.

The incidence of Grade 3 or 4 hyperglycemia in antiretroviral-naive subjects who received fosamprenavir calcium in the pivotal trials was less than 1%.

Table 5. Grade 3/4 Laboratory Abnormalities Reported in Greater than or Equal to 2% of Protease Inhibitor-Experienced Adult Subjects in Trial APV30003

Laboratory Abnormality	Fosamprenavir Calcium 700 mg and Ritonavir 100 mg Twice Daily ^a (n = 104)	Lopinavir 400 mg and Ritonavir 100 mg Twice Daily ^a (n = 103)
Triglycerides ^b (> 750 mg/dL)	11% ^c	6% ^c
Serum lipase (> 2 x ULN)	5%	12%
ALT (> 5 x ULN)	4%	4%
AST (> 5 x ULN)	4%	2%
Glucose (> 251 mg/dL)	2% ^c	2% ^c

^a All subjects also received 2 reverse transcriptase inhibitors.

^b Fasting specimens.

^c n = 100 for fosamprenavir calcium plus ritonavir, n = 98 for lopinavir plus ritonavir.

ULN = Upper limit of normal.

Pediatric Trials

Fosamprenavir calcium with and without ritonavir was studied in 237 HIV-1-infected pediatric subjects aged at least 4 weeks to 18 years in 3 open-label trials; APV20002, APV20003, and APV29005 [see *Clinical Studies (14.3)*]. Vomiting and neutropenia occurred more frequently in pediatric subjects compared with adults. Other adverse events occurred with similar frequency in pediatric subjects compared with adults.

The frequency of vomiting among pediatric subjects receiving fosamprenavir calcium twice daily with ritonavir was 20% in subjects aged at least 4 weeks to younger than 2 years and 36% in subjects aged 2 to 18 years compared with 10% in adults. The frequency of vomiting among pediatric subjects receiving fosamprenavir calcium twice daily without ritonavir was 60% in subjects aged 2 to 5 years compared with 16% in adults.

The median duration of drug-related vomiting episodes in APV29005 was 1 day (range: 1 to 3 days), in APV20003 was 16 days (range: 1 to 38 days), and in APV20002 was 9 days (range: 4 to 13 days). Vomiting was treatment limiting in 4 pediatric subjects across all 3 trials.

The incidence of Grade 3 or 4 neutropenia (neutrophils less than 750 cells per mm³) seen in pediatric subjects treated with fosamprenavir calcium with and without ritonavir was higher (15%) than the incidence seen in adult subjects (3%). Grade 3/4 neutropenia occurred in 10% (5 of 51) of subjects aged at least 4 weeks to younger than 2 years and 16% (28 of 170) of subjects aged 2 to 18 years.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of fosamprenavir calcium. Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

These reactions have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to fosamprenavir calcium.

Cardiac Disorders

Myocardial infarction.

Metabolism and Nutrition Disorders

Hypercholesterolemia.

Nervous System Disorders

Oral paresthesia.

Skin and Subcutaneous Tissue Disorders

Angioedema.

Urogenital

Nephrolithiasis.

7 DRUG INTERACTIONS

7.1 Cytochrome P450 Inhibitors and Inducers

Amprenavir, the active metabolite of fosamprenavir, is an inhibitor of CYP3A4 metabolism and therefore should not be administered concurrently with medications with narrow therapeutic windows that are substrates of CYP3A4. Data also suggest that amprenavir induces CYP3A4.

Amprenavir is metabolized by CYP3A4. Coadministration of fosamprenavir calcium and drugs that induce CYP3A4, such as rifampin, may decrease amprenavir concentrations and reduce its therapeutic effect. Coadministration of fosamprenavir calcium and drugs that inhibit CYP3A4 may increase amprenavir concentrations and increase the incidence of adverse effects.

The potential for drug interactions with fosamprenavir calcium changes when fosamprenavir calcium is coadministered with the potent CYP3A4 inhibitor ritonavir. The magnitude of CYP3A4-mediated drug interactions (effect on amprenavir or effect on coadministered drug) may change when fosamprenavir calcium is coadministered with ritonavir. Because ritonavir is a CYP2D6 inhibitor, clinically significant interactions with drugs metabolized by CYP2D6 are possible when coadministered with fosamprenavir calcium plus ritonavir. Ritonavir also appears to induce CYP3A, CYP1A2, CYP2C9, CYP2C19, and CYP2B6, as well as other enzymes, including glucuronosyl transferase.

There are other agents that may result in serious and/or life-threatening drug interactions [*see Contraindications (4)*].

7.2 Established and Other Potentially Significant Drug Interactions

If fosamprenavir calcium is used in combination with ritonavir, see full prescribing information for ritonavir for additional information on drug interactions [see *Contraindications (4)*, *Clinical Pharmacology (12.3)*].

Table 6 provides a listing of established or potentially clinically significant drug interactions. Information in the table applies to fosamprenavir calcium with or without ritonavir, unless otherwise indicated.

Table 6. Established and Other Potentially Significant Drug Interactions

Concomitant Drug Class: Drug Name	Effect on Concentration of Amprenavir or Concomitant Drug	Clinical Comment
<i>HCV/HIV-Antiviral Agents</i>		
HCV protease inhibitor: Boceprevir	Fos amprenavir calcium: ↓ Amprenavir (predicted) ↔ or ↓ Boceprevir (predicted) Fos amprenavir calcium /ritonavir: ↓ Amprenavir (predicted) ↓ Boceprevir (predicted)	Coadministration of fosamprenavir calcium or fosamprenavir calcium/ ritonavir and boceprevir is not recommended.
HCV protease inhibitor: Simeprevir	Fos amprenavir calcium: ↔ Amprenavir (predicted) ↑ or ↓ Simeprevir (predicted) Fos amprenavir calcium /ritonavir: ↔ Amprenavir (predicted) ↑ Simeprevir (predicted)	Coadministration of fosamprenavir calcium or fosamprenavir calcium/ritonavir and simeprevir is not recommended.
HCV protease inhibitor: Paritaprevir (coformulated with ritonavir and ombitasvir and coadministered with dasabuvir)	Fos amprenavir calcium: ↑ Amprenavir (predicted) ↑ or ↔ Paritaprevir (predicted) Fos amprenavir calcium /ritonavir: ↑ or ↔ Amprenavir (predicted) ↑ Paritaprevir (predicted)	Appropriate doses of the combinations with respect to safety and efficacy have not been established. Fosamprenavircalcium 1400 mg once daily may be considered when coadministered with paritaprevir/ritonavir/ombitasvir/ dasabuvir. Coadministration of Fosamprenavircalcium /ritonavir and paritaprevir/ritonavir/ombitasvir/ dasabuvir is not recommended.
Non-nucleoside reverse transcriptase inhibitor: Delavirdine ^a	Fos amprenavir calcium: ↑ Amprenavir ↓ Delavirdine Fos amprenavir	Coadministration is contraindicated as it may lead to loss of virologic response and possible resistance to delavirdine [see <i>Contraindications (4)</i>].

<p>Non-nucleoside reverse transcriptase inhibitor: Efavirenz^a</p>	<p>calcium/ritonavir: ↑ Amprenavir ↓ Delavirdine</p> <p>Fos amprenavir calcium: ↓ Amprenavir</p> <p>Fos amprenavir calcium/ritonavir: ↓ Amprenavir</p>	<p>Appropriate doses of the combinations with respect to safety and efficacy have not been established.</p> <p>An additional 100 mg/day (300 mg total) of ritonavir is recommended when efavirenz is administered with fosamprenavir calcium/ritonavir once daily. No change in the ritonavir dose is required when efavirenz is administered with Fosamprenavir calcium plus ritonavir twice daily.</p>
<p>Non-nucleoside reverse transcriptase inhibitor: Nevirapine^a</p>	<p>Fos amprenavir calcium: ↓ Amprenavir ↑ Nevirapine</p> <p>Fos amprenavir calcium/ritonavir: ↓ Amprenavir ↑ Nevirapine</p>	<p>Coadministration of nevirapine and fosamprenavir calcium without ritonavir is not recommended.</p> <p>No dosage adjustment required when nevirapine is administered with fosamprenavir calcium/ritonavir twice daily.</p> <p>The combination of nevirapine administered with fosamprenavir calcium/ritonavir once-daily regimen has not been studied.</p>
<p>HIV protease inhibitor: Atazanavir^a</p>	<p>Fos amprenavir calcium: Interaction has not been evaluated.</p> <p>Fos amprenavir calcium/ritonavir: ↓ Atazanavir ↔ Amprenavir</p>	<p>Appropriate doses of the combinations with respect to safety and efficacy have not been established.</p>
<p>HIV protease inhibitors: Indinavir^a, nelfinavir^a</p>	<p>Fos amprenavir calcium: ↑ Amprenavir Effect on indinavir and nelfinavir is not well established.</p> <p>Fos amprenavir calcium/ritonavir: Interaction has not been evaluated.</p>	<p>Appropriate doses of the combinations with respect to safety and efficacy have not been established.</p>
<p>HIV protease inhibitors: Lopinavir/ritonavir^a</p>	<p>↓ Amprenavir ↓ Lopinavir</p>	<p>An increased rate of adverse events has been observed. Appropriate doses of the combinations with respect to safety and efficacy have not been established.</p>
<p>HIV protease inhibitor: Saquinavir^a</p>	<p>Fos amprenavir calcium: ↓ Amprenavir Effect on saquinavir is not well established.</p> <p>Fos amprenavir calcium/ritonavir: Interaction has not been evaluated.</p>	<p>Appropriate doses of the combination with respect to safety and efficacy have not been established.</p>
<p>HIV integrase inhibitor: Raltegravir^a</p>	<p>Fos amprenavir calcium: ↓ Amprenavir ↓ Raltegravir</p>	<p>Appropriate doses of the combination with respect to safety and efficacy have not been established.</p>

	Fosamprenavir calcium/ritonavir: ↓ Amprenavir ↓ Raltegravir	
HIV integrase inhibitor: Dolutegravir ^a	Fosamprenavir calcium/ritonavir: ↓Dolutegravir	The recommended dose of dolutegravir is 50 mg twice daily when coadministered with Fosamprenavir calcium /ritonavir. Use an alternative combination where possible in patients with known or suspected integrase inhibitor resistance.
HIV CCR5 co-receptor antagonist: Maraviroc ^a	Fosamprenavir calcium/ritonavir: ↓ Amprenavir ↑ Maraviroc	No dosage adjustment required for fosamprenavir calcium /ritonavir. The recommended dose of maraviroc is 150 mg twice daily when coadministered with fosamprenavir calcium/ritonavir. Fosamprenavir calcium should be given with ritonavir when coadministered with maraviroc.
Other Agents		
Alpha 1-adrenoreceptor antagonist: Alfuzosin	↑ Alfuzosin	Coadministration is contraindicated due to potential hypotension [see <i>Contraindications (4)</i>].
Antacid: MAALOX TC	↓ Amprenavir	Use with caution when administered at the same time. Fosamprenavir calcium may be less effective due to decreased amprenavir plasma concentrations. Staggered coadministration of antacids and fosamprenavir calcium has not been evaluated.
Antiarrhythmics: Amiodarone, disopyramide, lidocaine (systemic), and quinidine Flecainide, propafenone	↑ Antiarrhythmics ↑ Antiarrhythmics	Therapeutic concentration monitoring, if available, is recommended for antiarrhythmics. Coadministration is contraindicated due to potential for serious and/or life-threatening reactions such as cardiac arrhythmias if fosamprenavir calcium is co-prescribed with ritonavir [see <i>Contraindications (4)</i>].
Anticoagulant: Warfarin		Concentrations of warfarin may be affected. It is recommended that INR (international normalized ratio) be monitored.
Anticonvulsants: Carbamazepine, phenobarbital, phenytoin Phenytoin ^a	Fosamprenavir calcium: ↓ Amprenavir Fosamprenavir calcium/ritonavir: ↑ Amprenavir ↓ Phenytoin	Use with caution. Fosamprenavir calcium may be less effective due to decreased amprenavir plasma concentrations in patients taking these agents concomitantly. Plasma phenytoin concentrations should be monitored and phenytoin dose should be increased as appropriate. No change in fosamprenavir calcium/ritonavir dose is recommended.
Antidepressant: Paroxetine, trazodone	↓ Paroxetine	Any paroxetine dose adjustment should be guided by clinical effect (tolerability and efficacy).

	↑ Trazodone	Adverse events of nausea, dizziness, hypotension, and syncope have been observed following coadministration of trazodone and ritonavir. If trazodone is used with a CYP3A4 inhibitor such as fosamprenavir calcium, the combination should be used with caution and a lower dose of trazodone should be considered.
Antifungals: Ketoconazole ^a , itraconazole	↑ Ketoconazole ↑ Itraconazole	Increase monitoring for adverse events. Fosamprenavir calcium: Dose reduction of ketoconazole or itraconazole may be needed for patients receiving more than 400 mg ketoconazole or itraconazole per day. Fosamprenavir calcium/ritonavir: High doses of ketoconazole or itraconazole (greater than 200 mg/day) are not recommended.
Anti-gout: Colchicine	↑ Colchicine	Patients with renal or hepatic impairment should not be given colchicine with fosamprenavir calcium/ritonavir. Fosamprenavir calcium/ritonavir and coadministration of colchicine: Treatment of gout flares: 0.6 mg (1 tablet) x 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Dose to be repeated no earlier than 3 days. Prophylaxis of gout flares: If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once every other day. Treatment of familial Mediterranean fever (FMF): Maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day). Fosamprenavir calcium and coadministration of colchicine: Treatment of gout flares: 1.2 mg (2 tablets) x 1 dose. Dose to be repeated no earlier than 3 days. Prophylaxis of gout flares: If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg twice a day or 0.6 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once a day. Treatment of FMF: Maximum daily dose of 1.2 mg (may be given as 0.6 mg twice a day).

<p>Antimycobacterial: Rifabutin^a</p>	<p>↑ Rifabutin and rifabutin metabolite</p>	<p>A complete blood count should be performed weekly and as clinically indicated to monitor for neutropenia. Fosamprenavir calcium: A dosage reduction of rifabutin by at least half the recommended dose is required. Fosamprenavir calcium/ritonavir: Dosage reduction of rifabutin by at least 75% of the usual dose of 300 mg/day is recommended (a maximum dose of 150 mg every other day or 3 times per week).</p>
<p>Antimycobacterial: Rifampin^a</p>	<p>↓ Amprenavir</p>	<p>Coadministration is contraindicated as it may lead to loss of virologic response and possible resistance to Fosamprenavir calcium or to the class of protease inhibitors [see <i>Contraindications (4)</i>].</p>
<p>Antineoplastics: Dasatinib, nilotinib, ibrutinib, vinblastine, everolimus</p>	<p>↑ Antineoplastics</p>	<p>Fosamprenavir calcium or fosamprenavir calcium/ritonavir may increase plasma concentrations of antineoplastics metabolized by CYP3A, potentially increasing the risk of adverse events typically associated with these medications. In case of coadministration, please refer to relevant prescribing information for these medications.</p>
<p>Antipsychotics: Lurasidone Pimozide Quetiapine</p>	<p>↑ Lurasidone ↑ Pimozide ↑ Quetiapine</p>	<p>Fosamprenavir calcium: If coadministration is necessary, reduce the lurasidone dose. Refer to the lurasidone prescribing information for concomitant use with moderate CYP3A4 inhibitors. Fosamprenavir calcium /ritonavir: Use of lurasidone is contraindicated due to potential for serious and/or life-threatening reactions [see <i>Contraindications (4)</i>]. Coadministration is contraindicated due to potential for serious and/or life-threatening reactions such as cardiac arrhythmias [see <i>Contraindications (4)</i>]. Fosamprenavir calcium /ritonavir: <u>Initiation of fosamprenavir calcium with ritonavir in patients taking quetiapine:</u> Consider alternative antiretroviral therapy to avoid increases in quetiapine drug exposures. If coadministration is necessary, reduce the quetiapine dose to 1/6 of the current dose and monitor for quetiapine-associated adverse reactions. Refer to the quetiapine prescribing information for recommendations on adverse reaction monitoring. <u>Initiation of quetiapine in patients taking fosamprenavir calcium with ritonavir:</u> Refer</p>

		to the quetiapine prescribing information for initial dosing and titration of quetiapine.
Benzodiazepines: Alprazolam, clorazepate, diazepam, flurazepam	↑ Benzodiazepines	Clinical significance is unknown. A decrease in benzodiazepine dose may be needed.
Calcium channel blockers: Diltiazem, felodipine, nifedipine, nicardipine, nimodipine, verapamil, amlodipine, nisoldipine, isradipine	↑ Calcium channel blockers	Use with caution. Clinical monitoring of patients is recommended.
Corticosteroid: Dexamethasone	↓ Amprenavir	Use with caution. Fosamprenavir calcium may be less effective due to decreased amprenavir plasma concentrations.
Endothelin-receptor antagonist: Bosentan	↑ Bosentan	<p><u>Coadministration of bosentan in patients on fosamprenavir calcium:</u></p> <p>In patients who have been receiving fosamprenavir calcium for at least 10 days, start bosentan at 62.5 mg once daily or every other day based upon individual tolerability.</p> <p><u>Coadministration of fosamprenavir calcium in patients on bosentan:</u></p> <p>Discontinue use of bosentan at least 36 hours prior to initiation of fosamprenavir calcium.</p> <p>After at least 10 days following the initiation of fosamprenavir calcium, resume bosentan at 62.5 mg once daily or every other day based upon individual tolerability.</p>
Ergot derivatives: Dihydroergotamine, ergonovine, ergotamine, methylergonovine	↑ Ergot derivatives	Coadministration is contraindicated due to potential for serious and/or life-threatening reactions such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues [see <i>Contraindications (4)</i>].
GI motility agent: Cisapride	↑ Cisapride	Coadministration is contraindicated due to potential for serious and/or life-threatening reactions such as cardiac arrhythmias [see <i>Contraindications (4)</i>].
Herbal product: St. John's wort (<i>Hypericum perforatum</i>)	↓ Amprenavir	Coadministration is contraindicated as it may lead to loss of virologic response and possible resistance to Fosamprenavir

		calcium or to the class of protease inhibitors [see <i>Contraindications (4)</i>].
Histamine H₂-receptor antagonists: Cimetidine, famotidine, nizatidine, ranitidine ^a	Fosamprenavir calcium: ↓ Amprenavir Fosamprenavir calcium/ritonavir: Interaction not evaluated	Use with caution. Fosamprenavir calcium may be less effective due to decreased amprenavir plasma concentrations.
Immunosuppressants: Cyclosporine, tacrolimus, sirolimus	↑ Immunosuppressants	Therapeutic concentration monitoring is recommended for immunosuppressant agents.
Inhaled beta-agonist: Salmeterol	↑ Salmeterol	Concurrent administration of salmeterol with fosamprenavir calcium is not recommended. The combination may result in increased risk of cardiovascular adverse events associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia.
Inhaled/nasal steroid: Fluticasone	Fosamprenavir calcium: ↑ Fluticasone Fosamprenavir calcium/ritonavir: ↑ Fluticasone	Use with caution. Consider alternatives to fluticasone, particularly for long-term use. May result in significantly reduced serum cortisol concentrations. Systemic corticosteroid effects including Cushing's syndrome and adrenal suppression have been reported during postmarketing use in patients receiving ritonavir and inhaled or intranasally administered fluticasone. Coadministration of fluticasone and fosamprenavir calcium/ritonavir is not recommended unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side effects.
Lipid Modifying Agents: <u>HMG-CoA reductase inhibitors:</u> Atorvastatin ^a Lovastatin, simvastatin <u>Other lipid modifying agents:</u> Lomitapide	↑ Atorvastatin ↑ Lovastatin ↑ Simvastatin ↑ Lomitapide	Titrate atorvastatin dose carefully and use the lowest necessary dose; do not exceed atorvastatin 20 mg/day. Coadministration with lovastatin or simvastatin is contraindicated due to potential for serious reactions such as risk of myopathy including rhabdomyolysis [see <i>Contraindications (4)</i>]. Coadministration with lomitapide is contraindicated due to potential for markedly increased transaminases.
Narcotic analgesic: Methadone Fentanyl	↓ Methadone ↑ Fentanyl	Data suggest that the interaction is not clinically relevant; however, patients should be monitored for opiate withdrawal symptoms. Careful monitoring of therapeutic effects and adverse effects of fentanyl (including potentially fatal respiratory depression) is recommended.
Oral contraceptives: Ethinyl		Alternative methods of non-hormonal contraception are recommended.

<p>estradiol/norethindrone^a</p>	<p>Fosamprenavir calcium: ↓ Amprenavir ↓ Ethinyl estradiol</p> <p>Fosamprenavir calcium/ritonavir: ↓ Ethinyl estradiol</p>	<p>May lead to loss of virologic response.^a</p> <p>Increased risk of transaminase elevations. No data are available on the use of fosamprenavir calcium/ritonavir with other hormonal therapies, such as hormone replacement therapy (HRT) for postmenopausal women.</p>
<p>PDE5 inhibitors: Sildenafil, tadalafil, vardenafil</p>	<p>↑ Sildenafil ↑ Tadalafil ↑ Vardenafil</p>	<p>May result in an increase in PDE5 inhibitor-associated adverse events, including hypotension, syncope, visual disturbances, and priapism.</p> <p><u>Use of PDE5 inhibitors for pulmonary arterial hypertension (PAH):</u></p> <ul style="list-style-type: none"> • Use of sildenafil (REVATIO) is contraindicated when used for the treatment of PAH. A safe and effective dose has not been established when used with fosamprenavir calcium. There is increased potential for sildenafil-associated adverse events (which include visual disturbances, hypotension, prolonged erection, and syncope) [see Contraindications (4)]. • <u>The following dose adjustments are recommended for use of tadalafil (ADCIRCA[®]) with fosamprenavir calcium:</u> <u>Coadministration of ADCIRCA in patients on fosamprenavir calcium:</u> In patients receiving fosamprenavir calcium for at least one week, start ADCIRCA at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability. <u>Coadministration of fosamprenavir calcium in patients on ADCIRCA:</u> Avoid use of ADCIRCA during the initiation of fosamprenavir calcium. Stop ADCIRCA at least 24 hours prior to starting fosamprenavir calcium. After at least one week following the initiation of fosamprenavir calcium, resume ADCIRCA at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability. <p><u>Use of PDE5 inhibitors for erectile</u></p>

		<u>dysfunction:</u> Fosamprenavir calcium: Sildenafil: 25 mg every 48 hours. Tadalafil: no more than 10 mg every 72 hours. Vardenafil: no more than 2.5 mg every 24 hours. Fosamprenavir calcium/ritonavir: Sildenafil: 25 mg every 48 hours. Tadalafil: no more than 10 mg every 72 hours. Vardenafil: no more than 2.5 mg every 72 hours. Use with increased monitoring for adverse events.
Proton pump inhibitors: Esomeprazole ^a , lansoprazole, omeprazole, pantoprazole, rabeprazole	Fosamprenavir calcium: ↔ Amprenavir ↑ Esomeprazole Fosamprenavir calcium/ritonavir: ↔ Amprenavir ↔ Esomeprazole	Proton pump inhibitors can be administered at the same time as a dose of fosamprenavir calcium with no change in plasma amprenavir concentrations.
Sedative/hypnotics: Midazolam, triazolam	↑ Midazolam ↑ Triazolam	Coadministration is contraindicated due to potential for serious and/or life-threatening reactions such as prolonged or increased sedation or respiratory depression [see <i>Contraindications (4)</i>].
Tricyclic antidepressants: Amitriptyline, imipramine	↑ Tricyclics	Therapeutic concentration monitoring is recommended for tricyclic antidepressants.

^a See *Clinical Pharmacology (12.3) Tables 10, 11, 12, or 13 for magnitude of interaction.*

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to fosamprenavir calcium during pregnancy. Healthcare providers are encouraged to register patients by calling the Antiretroviral Pregnancy Registry (APR) at 1-800-258-4263.

Risk Summary

Limited data are available for use of fosamprenavir calcium in pregnancy. fosamprenavir calcium 700 mg twice daily taken with ritonavir 100 mg twice daily should only be considered in pregnant patients who are already on a stable twice-daily regimen of fosamprenavir calcium/ritonavir 700 mg/100 mg prior to pregnancy, and who are virologically suppressed (HIV-1 RNA less than 50 copies per mL) (*see Clinical Considerations and Data*).

There are insufficient human data on the use of fosamprenavir during pregnancy to adequately assess a drug-associated risk for birth defects and miscarriage. Given the limited number of pregnancies exposed to fosamprenavir-based regimens, no conclusions can be drawn on the safety of fosamprenavir

in pregnancy. The background risk for major birth defects and miscarriage for the indicated population is unknown. The background rate for major birth defects in a U.S. reference population of the Metropolitan Atlanta Congenital Defects Program (MACDP) is 2.7% (*see Data*). The estimated background rate of miscarriage in clinically recognized pregnancies in the U.S. general population is 15% to 20%.

In animal reproduction studies, no evidence of major adverse developmental outcomes was observed following oral administration of fosamprenavir. Systemic exposure to amprenavir (the active ingredient) was less than (rabbits) or up to 2 times (rats) those in humans at the maximum recommended human dose (MRHD) with or without ritonavir. In contrast, oral administration of amprenavir was associated with abortions in pregnant rabbits at doses that produced approximately one-twentieth the human exposure at the MRHD.

In the rat pre- and postnatal development study, toxicities to the offspring, including reduced survival and reproductive performance, were observed at maternal systemic exposures (AUC) to amprenavir that were approximately 2 times the exposure in humans at the MRHD of fosamprenavir alone or approximately the same as those seen in humans following administration of the MRHD of fosamprenavir in combination with ritonavir (*see Data*).

Clinical Considerations

Virologic Monitoring During Pregnancy and the Postpartum Period: Based on limited data on the use of fosamprenavir calcium during pregnancy, no dosage adjustments are required for pregnant patients who are already on a stable twice-daily regimen of fosamprenavir calcium 700 mg taken with ritonavir 100 mg prior to pregnancy, and who are virologically suppressed (HIV-1 RNA less than 50 copies per mL) [*see Dosage and Administration (2.3), Clinical Pharmacology (12.3)*]. In a clinical trial of 10 HIV-1–infected pregnant women treated with fosamprenavir calcium 700 mg taken with ritonavir 100 mg twice daily through postpartum, total amprenavir exposures were lower during pregnancy compared with the postpartum period. Therefore, viral load should be monitored closely to ensure viral suppression is maintained [*see Data, Dosage and Administration (2.2), Clinical Pharmacology (12.3)*]. Pregnancy data with other dosage regimens of fosamprenavir calcium (with or without ritonavir) are not available.

Data

Human Data: fosamprenavir calcium 700 mg taken with ritonavir 100 mg twice daily in combination with a background regimen was evaluated in a clinical trial of 10 HIV-1–infected pregnant women during the second and third trimesters and postpartum. Subjects initiated fosamprenavir calcium /ritonavir during pregnancy at a median of 19 weeks' gestation; 4 had undetectable HIV-1 RNA viral load (less than 50 copies/mL) at the time of initiation. Amprenavir pharmacokinetics and placental transfer were studied during the second trimester (n = 6) or third trimester (n = 9) and postpartum (n = 9). Pregnancy outcomes were available for all 10 subjects, among which 1 twin pregnancy was included. Compared to the postpartum period, geometric mean amprenavir AUC was 35% lower in the second trimester and 25% lower in the third trimester. The amprenavir geometric mean ratio (95% CI) of fetal cord to maternal peripheral plasma concentration (n = 7) was 0.27 (0.24 to 0.30) [*see Clinical Pharmacology (12.3)*]. At delivery, 9 subjects had HIV-1 viral load less than 50 copies/mL, and 1 subject had a viral load of 111 copies/mL. All 11 infants born had test results that were negative for HIV-1 at the time of delivery and through 12 months postpartum. There were no new safety findings compared with the known safety profile of fosamprenavir calcium/ritonavir in HIV-1–infected adults.

Based on prospective reports to the APR of approximately 146 live births following exposure to fosamprenavir-containing regimens, there were 2 birth defects reported in 109 first trimester exposures and 2 birth defects reported in 36 second and third trimester exposures. The background rate for major birth defects is 2.7% in a U.S. reference population of the MACDP. Prospective reports from the APR of overall major birth defects in pregnancies exposed to fosamprenavir are compared with the U.S. background major birth defect rate. Methodological limitations of the APR include the use of MACDP as the external comparator group. Limitations of using an external comparator include differences in methodology and populations as well as confounding due to the underlying disease.

Animal Data: Fosamprenavir was administered orally to pregnant rats (300, 820, or 2,240 mg per kg per day) and rabbits (74.8, 224.3, or 672.8 mg per kg per day) on Gestation Days 6 to 17 and Days 7 to 20, respectively. No major adverse effects on embryo-fetal development were observed at these dose levels, resulting in exposures ($AUC_{0-24\text{ h}}$) approximately 2 times (rats) and 0.8 times (rabbits) human exposures at the MRHD of fosamprenavir alone or 0.7 times (rats) and 0.3 times (rabbits) human exposures at the MRHD of fosamprenavir in combination with ritonavir. However, increased incidence of abortion was observed in rabbits administered a maternally toxic dose of fosamprenavir (672.8 mg per kg per day). In a study where amprenavir was administered orally to pregnant rabbits (25, 50, or 100 mg per kg per day) on Gestation Days 8 to 20, increased abortions and an increased incidence of minor skeletal variations (deficient ossification of the femur, humerus, and trochlea) were observed at doses that produced approximately one-twentieth the exposure seen at the MRHD.

In the rat pre- and postnatal development study, fosamprenavir was administered orally (300, 820, or 2,240 mg per kg per day) on Gestation Day 6 to Lactation/Postpartum Day 20. Fosamprenavir caused a reduction in pup survival and body weights. In surviving female offspring from the high-dose group, an increased time to successful mating, an increased length of gestation, a reduced number of uterine implantation sites per litter, and reduced gestational body weights were observed. Systemic exposure ($AUC_{0-24\text{ h}}$) to amprenavir in rats was approximately 2 times the exposures in humans at the MRHD of fosamprenavir alone or approximately the same as those seen in humans at the MRHD of fosamprenavir in combination with ritonavir.

8.2 Lactation

Risk Summary

The Centers for Disease Control and Prevention recommends that HIV-1-infected mothers in the United States not breastfeed their infants to avoid risking postnatal transmission of HIV-1 infection.

There is no information available on the presence of amprenavir in human milk, the effects of the drug on the breastfed infant, or the effects of the drug on milk production. When administered to lactating rats, amprenavir was present in milk (*see Data*). Because of the potential for (1) HIV-1 transmission (in HIV-negative infants), (2) developing viral resistance (in HIV-positive infants), and (3) adverse reactions in a breastfed infant, instruct mothers not to breastfeed if they are receiving fosamprenavir calcium.

Data

Amprenavir was excreted into the milk of lactating rats following a single dose of amprenavir (100 mg per kg); a maximal milk concentration was achieved 2 hours post-administration at a milk concentration approximately 1.2 times that of maternal plasma concentrations.

8.3 Females and Males of Reproductive Potential

Contraception

Use of fosamprenavir calcium may reduce the efficacy of combined hormonal contraceptives. Advise patients using combined hormonal contraceptives to use an effective alternative contraceptive method or an additional barrier method of contraception [*see Drug Interactions (7.2)*].

8.4 Pediatric Use

The safety, pharmacokinetic profile, and virologic and immunologic responses of fosamprenavir calcium with and without ritonavir were evaluated in protease inhibitor-naïve and -experienced HIV-1-infected pediatric subjects aged at least 4 weeks to younger than 18 years and weighing at least 3 kg in 3 open-label trials [*see Adverse Reactions (6.1), Clinical Pharmacology (12.3), Clinical Studies (14.3)*].

Treatment with fosamprenavir calcium is not recommended in protease inhibitor-experienced pediatric patients younger than 6 months. The pharmacokinetics, safety, tolerability, and efficacy of

fosamprenavir calcium in pediatric patients younger than 4 weeks have not been established [see *Clinical Pharmacology (12.3)*]. Available pharmacokinetic and clinical data do not support once-daily dosing of fosamprenavir calcium alone or in combination with ritonavir for any pediatrics or twice-daily dosing without ritonavir in pediatric patients younger than 2 years [see *Clinical Pharmacology (12.3)*, *Clinical Studies (14.3)*]. See *Dosage and Administration (2.3)* for dosing recommendations for pediatric patients.

8.5 Geriatric Use

Clinical studies of fosamprenavir calcium did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger adults. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

8.6 Hepatic Impairment

Amprenavir is principally metabolized by the liver; therefore, caution should be exercised when administering fosamprenavir calcium to patients with hepatic impairment because amprenavir concentrations may be increased [see *Clinical Pharmacology (12.3)*]. Patients with impaired hepatic function receiving fosamprenavir calcium with or without concurrent ritonavir require dose reduction [see *Dosage and Administration (2.4)*].

There are no data to support dosing recommendations for pediatric subjects with hepatic impairment.

10 OVERDOSAGE


In a healthy volunteer repeat-dose pharmacokinetic trial evaluating high-dose combinations of fosamprenavir calcium plus ritonavir, an increased frequency of Grade 2/3 ALT elevations (greater than 2.5 x ULN) was observed with fosamprenavir calcium 1,400 mg twice daily plus ritonavir 200 mg twice daily (4 of 25 subjects). Concurrent Grade 1/2 elevations in AST (greater than 1.25 x ULN) were noted in 3 of these 4 subjects. These transaminase elevations resolved following discontinuation of dosing.

There is no known antidote for fosamprenavir calcium. It is not known whether amprenavir can be removed by peritoneal dialysis or hemodialysis, although it is unlikely as amprenavir is highly protein bound. If overdosage occurs, the patient should be monitored for evidence of toxicity and standard supportive treatment applied as necessary.

11 DESCRIPTION

Fosamprenavir calcium is a prodrug of amprenavir, an inhibitor of HIV protease. The chemical name of fosamprenavir calcium is (3S)-tetrahydrofuran-3-yl (1S,2R)-3-[[[(4-aminophenyl) sulfonyl](isobutyl)amino]-1-benzyl-2-(phosphonoxy) propyl]carbamate monocalcium salt.

Fosamprenavir calcium is a single stereoisomer with the (3S)(1S,2R) configuration. It has a molecular formula of C₂₅H₃₄CaN₃O₉PS and a molecular weight of 623.7. It has the following structural formula:



Fosamprenavir calcium (amorphous) is a white to cream color powder which is soluble in methanol.

Fosamprenavir calcium tablets are available for oral administration in a strength of 700 mg of fosamprenavir as fosamprenavir calcium (amorphous) (equivalent to approximately 600 mg of amprenavir). Each 700 mg tablet contains the inactive ingredients colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, and povidone. The tablet coating contains the inactive ingredients ethylcellulose, ferric oxide red, hypromellose, talc, and titanium dioxide. The imprinting ink contains the inactive ingredients ferric oxide black, propylene glycol, and shellac.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Fosamprenavir is an HIV-1 antiretroviral agent [see *Microbiology (12.4)*].

12.3 Pharmacokinetics

The pharmacokinetic properties of amprenavir after administration of fosamprenavir calcium, with or without ritonavir, have been evaluated in both healthy adult volunteers and in HIV-1-infected subjects; no substantial differences in steady-state amprenavir concentrations were observed between the 2 populations.

The pharmacokinetic parameters of amprenavir after administration of fosamprenavir calcium (with and without concomitant ritonavir) are shown in Table 7.

Table 7. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters in Adults

Regimen	C _{max} (mcg/mL)	T _{max} (hours) ^a	AUC ₂₄ (mcg•h/mL)	C _{tau} ^b (mcg/mL)
Fosamprenavir calcium 1,400 mg Twice Daily (n = 22) ^c	4.82 (4.06 to 5.72)	1.3 (0.8 to 4)	33 ^d (27.6 to 39.2)	0.35 (0.27 to 0.46)
Fosamprenavir calcium 1,400 mg Once Daily plus Ritonavir 200 mg	7.24 (6.32 to 8.28)	2.1 (0.8 to 5)	69.4 (59.7 to 80.8)	1.45 (1.16 to 1.81)

Once Daily (n = 22) ^e				
Fosamprenavir calcium 1,400 mg Once Daily plus Ritonavir 100 mg Once Daily (n = 36) ^e	7.93 (7.25 to 8.68)	1.5 (0.75 to 5)	66.4 (61.1 to 72.1)	0.86 (0.74 to 1.01)
Fosamprenavir calcium 700 mg Twice Daily plus Ritonavir 100 mg Twice Daily (n = 24) ^e	6.08 (5.38 to 6.86)	1.5 (0.75 to 5)	79.2 ^d (69 to 90.6)	2.12 (1.77 to 2.54)

^a Data shown are median (range).

^b C_{tau} is the concentration at the end of the dose interval.


^c HIV-infected adults.

^d AUC₂₄ was calculated from AUC₁₂ summary data x 2.

^e Healthy adults.

The mean plasma amprenavir concentrations of the dosing regimens over the dosing intervals are displayed in Figure 1.

Figure 1. Mean (±SD) Steady-State Plasma Amprenavir Concentrations and Mean EC₅₀ Values against HIV from Protease Inhibitor-Naive Subjects (in the Absence of Human Serum)



Absorption

After administration of a single dose of fosamprenavir calcium to HIV-1-infected subjects, the time to peak amprenavir concentration (T_{max}) occurred between 1.5 and 4 hours (median 2.5 hours). The absolute oral bioavailability of amprenavir after administration of fosamprenavir calcium in humans has not been established.

After administration of a single 1,400-mg dose in the fasted state, fosamprenavir calcium oral suspension (50 mg per mL) and fosamprenavir calcium tablets (700 mg) provided similar amprenavir exposures (AUC); however, the C_{max} of amprenavir after administration of the suspension formulation was 14.5% higher compared with the tablet.

Amprenavir is both a substrate for and inducer of P-glycoprotein.

Effects of Food on Oral Absorption

Administration of a single 1,400 mg dose of fosamprenavir calcium tablets in the fed state (standardized high-fat meal: 967 kcal, 67 grams fat, 33 grams protein, 58 grams carbohydrate) compared with the fasted state was associated with no significant changes in amprenavir C_{max} , T_{max} , or $AUC_{0-\infty}$ [see *Dosage and Administration (2)*].

Administration of a single 1,400 mg dose of fosamprenavir calcium oral suspension in the fed state (standardized high-fat meal: 967 kcal, 67 grams fat, 33 grams protein, 58 grams carbohydrate) compared with the fasted state was associated with a 46% reduction in C_{max} , a 0.72-hour delay in T_{max} , and a 28% reduction in amprenavir $AUC_{0-\infty}$.

Distribution

In vitro, amprenavir is approximately 90% bound to plasma proteins, primarily to α_1 -acid glycoprotein. *In vitro*, concentration-dependent binding was observed over the concentration range of 1 to 10 mcg per mL, with decreased binding at higher concentrations. The partitioning of amprenavir into erythrocytes is low, but increases as amprenavir concentrations increase, reflecting the higher amount

of unbound drug at higher concentrations.

Metabolism

After oral administration, fosamprenavir is rapidly and almost completely hydrolyzed to amprenavir and inorganic phosphate prior to reaching the systemic circulation. This occurs in the gut epithelium during absorption. Amprenavir is metabolized in the liver by the CYP3A4 enzyme system. The 2 major metabolites result from oxidation of the tetrahydrofuran and aniline moieties. Glucuronide conjugates of oxidized metabolites have been identified as minor metabolites in urine and feces.

Elimination

Excretion of unchanged amprenavir in urine and feces is minimal. Unchanged amprenavir in urine accounts for approximately 1% of the dose; unchanged amprenavir was not detectable in feces. Approximately 14% and 75% of an administered single dose of ¹⁴C-amprenavir can be accounted for as metabolites in urine and feces, respectively. Two metabolites accounted for greater than 90% of the radiocarbon in fecal samples. The plasma elimination half-life of amprenavir is approximately 7.7 hours.

Specific Populations

Hepatic Impairment: The pharmacokinetics of amprenavir have been studied after the administration of fosamprenavir calcium in combination with ritonavir to adult HIV-1-infected subjects with mild, moderate, and severe hepatic impairment. Following 2 weeks of dosing with fosamprenavir calcium plus ritonavir, the AUC of amprenavir was increased by approximately 22% in subjects with mild hepatic impairment, by approximately 70% in subjects with moderate hepatic impairment, and by approximately 80% in subjects with severe hepatic impairment compared with HIV-1-infected subjects with normal hepatic function. Protein binding of amprenavir was decreased in subjects with hepatic impairment. The unbound fraction at 2 hours (approximate C_{max}) ranged between a decrease of -7% to an increase of 57% while the unbound fraction at the end of the dosing interval (C_{min}) increased from 50% to 102% [see *Dosage and Administration (2.4)*].

The pharmacokinetics of amprenavir have been studied after administration of amprenavir given as AGENERASE[®] capsules to adult subjects with hepatic impairment. Following administration of a single 600-mg oral dose, the AUC of amprenavir was increased by approximately 2.5-fold in subjects with moderate cirrhosis and by approximately 4.5-fold in subjects with severe cirrhosis compared with healthy volunteers [see *Dosage and Administration (2.4)*].

Patients with Renal Impairment: The impact of renal impairment on amprenavir elimination in adults has not been studied. The renal elimination of unchanged amprenavir represents approximately 1% of the administered dose; therefore, renal impairment is not expected to significantly impact the elimination of amprenavir.

Pregnant Women: Amprenavir pharmacokinetics were studied in pregnant women receiving fosamprenavir calcium (700 mg) plus ritonavir (100 mg) twice daily during the second trimester (n = 6) or third trimester (n = 9) and postpartum (n = 9). Compared to postpartum, geometric mean amprenavir AUC was 35% lower in the second trimester and 25% lower in the third trimester (Table 8). This decrease results in amprenavir concentrations that are within the range observed across regimens of fosamprenavir calcium in non-pregnant adults and lower concentrations compared with fosamprenavir calcium (700 mg) plus ritonavir (100 mg) twice daily in non-pregnant adults (Table 7, Table 8). This decrease is not expected to be considered clinically relevant in patients who are virologically suppressed; however, viral load should be monitored closely to ensure viral suppression is maintained [see *Dosage and Administration (2.2)*, *Use in Specific Populations (8.1)*]. The amprenavir geometric mean ratio (95% CI) of fetal cord to maternal peripheral plasma concentration (n = 7) was 0.27 (0.24 to 0.30).

Table 8. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters in Pregnant Women Receiving Fosamprenavir calcium with Ritonavir

Pharmacokinetic Parameter	Fosamprenavir calcium 700 mg Twice Daily plus Ritonavir 100 mg Twice Daily		
	Second Trimester (n = 6)	Third Trimester (n = 9)	Postpartum (n = 9)
AUC ₁₂ (mcg•h/mL)	26.0 (19.5, 34.6)	30.1 (21.6, 41.9)	39.9 (31.9, 50.1)
AUC ₂₄ (mcg•h/mL) ^a	52.0 (39.0, 69.2)	60.2 (43.2, 83.8)	79.8 (63.8, 100.2)
C _{max} (mcg/mL)	4.19 (3.19, 5.51)	5.36 (3.98, 7.22)	6.65 (5.24, 8.44)
C _{tau} (mcg/mL) ^b	1.31 (0.97, 1.77)	1.34 (0.95, 1.89)	2.03 (1.46, 2.83)

^a AUC₂₄ was calculated from AUC₁₂ summary data x 2.

^b C_{tau} represents the concentration at the end of the dose interval.

Pediatric Patients: The pharmacokinetics of amprenavir following administration of fosamprenavir calcium oral suspension and fosamprenavir calcium tablets, with or without ritonavir, have been studied in a total of 212 HIV-1-infected pediatric subjects enrolled in 3 trials. Fosamprenavir calcium without ritonavir was administered as 30 or 40 mg per kg twice daily to children aged 2 to 5 years.

Fosamprenavir calcium with ritonavir was administered as fosamprenavir calcium 30 mg per kg plus ritonavir 6 mg per kg once daily to children aged 2 to 18 years and as fosamprenavir calcium 18 to 60 mg per kg plus ritonavir 3 to 10 mg per kg twice daily to children aged at least 4 weeks to 18 years; body weights ranged from 3 to 103 kg.

Amprenavir apparent clearance decreased with increasing weight. Weight-adjusted apparent clearance was higher in children younger than 4 years, suggesting that younger children require higher mg-per-kg dosing of fosamprenavir calcium.

The pharmacokinetics of fosamprenavir calcium oral suspension in protease inhibitor-naive infants younger than 6 months (n = 9) receiving fosamprenavir calcium 45 mg per kg plus ritonavir 10 mg per kg twice daily generally demonstrated lower AUC₁₂ and C_{min} than adults receiving twice-daily fosamprenavir calcium 700 mg plus ritonavir 100 mg, the dose recommended for protease-experienced adults. The mean steady-state amprenavir AUC₁₂, C_{max}, and C_{min} were 26.6 mcg•hour per mL, 6.25 mcg per mL, and 0.86 mcg per mL, respectively. Because of expected low amprenavir exposure and a requirement for large volume of drug, twice-daily dosing of fosamprenavir calcium alone (without ritonavir) in pediatric subjects younger than 2 years was not studied.

Pharmacokinetic parameters for fosamprenavir calcium administered with food and with ritonavir in this patient population at the recommended weight-band-based dosage regimens are provided in Table 9.

Table 9. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters by Weight in Pediatric and Adolescent Subjects Aged at Least 4 Weeks to 18 Years Receiving Fosamprenavir Calcium with Ritonavir

Weight	Recommended Dosage Regimen	C _{max}		AUC ₂₄		C _{min}	
		n	(mcg/mL)	n	(mcg•h/mL)	n	(mcg/mL)
< 11 kg	Fosamprenavir calcium 45 mg/kg plus Ritonavir 7 mg/kg twice daily	12	6 (3.88, 9.29)	12	57.3 (34.1, 96.2)	27	1.65 (1.22, 2.24)
11 kg - < 15 kg	Fosamprenavir calcium 30 mg/kg	Not studied ^a					

	plus Ritonavir 3 mg/kg twice daily						
15 kg - < 20 kg	Fosamprenavir calcium 23 mg/kg plus Ritonavir 3 mg/kg twice daily	5	9.54 (4.63, 19.7)	5	121 (54.2, 269)	9	3.56 (2.33, 5.43)
20 kg - < 39 kg	Fosamprenavir calcium 18 mg/kg plus Ritonavir 3 mg/kg twice daily	13	6.24 (5.01, 7.77)	12	97.9 (77, 124)	23	2.54 (2.11, 3.06)
≥ 39 kg	Fosamprenavir calcium 700 mg plus Ritonavir 100 mg twice daily	15	5.03 (4.04, 6.26)	15	72.3 (59.6, 87.6)	42	1.98 (1.72, 2.29)

^a Recommended dose for pediatric patients weighing 11 kg to less than 15 kg is based on population pharmacokinetic analysis.

Subjects aged 2 to younger than 6 years receiving fosamprenavir calcium 30 mg per kg twice daily without ritonavir achieved geometric mean (95% CI) amprenavir C_{max} (n = 9), AUC_{12} (n = 9), and C_{min} (n = 19) of 7.15 (5.05, 10.1), 22.3 (15.3, 32.6), and 0.513 (0.384, 0.686), respectively.

Geriatric Patients: The pharmacokinetics of amprenavir after administration of fosamprenavir calcium to patients older than 65 years have not been studied [see *Use in Specific Populations (8.5)*].

Male and Female Patients: The pharmacokinetics of amprenavir after administration of fosamprenavir calcium do not differ between males and females.

Racial Groups: The pharmacokinetics of amprenavir after administration of fosamprenavir calcium do not differ between blacks and non-blacks.

Drug Interaction Studies

[See *Contraindications (4)*, *Warnings and Precautions (5.1)*, *Drug Interactions (7)*.]

Amprenavir, the active metabolite of fosamprenavir, is metabolized in the liver by the cytochrome P450 enzyme system. Amprenavir inhibits CYP3A4. Data also suggest that amprenavir induces CYP3A4. Caution should be used when coadministering medications that are substrates, inhibitors, or inducers of CYP3A4, or potentially toxic medications that are metabolized by CYP3A4. Amprenavir does not inhibit CYP2D6, CYP1A2, CYP2C9, CYP2C19, CYP2E1, or uridine glucuronosyltransferase (UDPGT). Amprenavir is both a substrate for and inducer of P-glycoprotein.

Drug interaction trials were performed with fosamprenavir calcium and other drugs likely to be coadministered or drugs commonly used as probes for pharmacokinetic interactions. The effects of coadministration on AUC , C_{max} , and C_{min} values are summarized in Table 10 (effect of other drugs on amprenavir) and Table 12 (effect of fosamprenavir calcium on other drugs). In addition, since fosamprenavir calcium delivers comparable amprenavir plasma concentrations as AGENERASE, drug interaction data derived from trials with AGENERASE are provided in Tables 11 and 13. For information regarding clinical recommendations, [see *Drug Interactions (7)*].

Table 10. Drug Interactions: Pharmacokinetic Parameters for Amprenavir after Administration of Fosamprenavir Calcium in the Presence of the Coadministered Drug(s)

Coadministered Drug(s) and Dose(s)	Dose of Fosamprenavir Calcium ^a	n	% Change in Amprenavir Pharmacokinetic Parameters (90% CI)		
			C_{max}	AUC	C_{min}

Antacid (MAALOX TC) 30 mL single dose	1,400-mg single dose	30		↓ 35 (↓ 24 to ↓ 42)	↓ 18 (↓ 9 to ↓ 26)	↑ 14 (↓ 7 to ↑ 39)
Atazanavir 300 mg once daily for 10 days	700 mg twice daily plus ritonavir 100 mg twice daily for 10 days	22		↔	↔	↔
Atorvastatin 10 mg once daily for 4 days	1,400 mg twice daily for 2 weeks	16		↓ 18 (↓ 34 to ↑ 1)	↓ 27 (↓ 41 to ↓ 12)	↓ 12 (↓ 27 to ↓ 6)
Atorvastatin 10 mg once daily for 4 days	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	16		↔	↔	↔
Efavirenz 600 mg once daily for 2 weeks	1,400 mg once daily plus ritonavir 200 mg once daily for 2 weeks	16		↔	↓ 13 (↓ 30 to ↑ 7)	↓ 36 (↓ 8 to ↓ 56)
Efavirenz 600 mg once daily plus additional ritonavir 100 mg once daily for 2 weeks	1,400 mg once daily plus ritonavir 200 mg once daily for 2 weeks	16		↑ 18 (↑ 1 to ↑ 38)	↑ 11 (0 to ↑ 24)	↔
Efavirenz 600 mg once daily for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	16		↔	↔	↓ 17 (↓ 4 to ↓ 29)
Esomeprazole 20 mg once daily for 2 weeks	1,400 mg twice daily for 2 weeks	25		↔	↔	↔
Esomeprazole 20 mg once daily for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	23		↔	↔	↔
Ethinyl estradiol/norethindrone	700 mg twice daily plus	25				

0.035 mg/0.5 mg once daily for 21 days	ritonavir ^b 100 mg twice daily for 21 days		↔ ^c	↔ ^c	↔ ^c
Ketoconazole ^d 200 mg once daily for 4 days	700 mg twice daily plus ritonavir 100 mg twice daily for 4 days	15	↔	↔	↔
Lopinavir/ritonavir 533 mg/133 mg twice daily	1,400 mg twice daily for 2 weeks	18	↓ 13 ^e	↓ 26 ^e	↓ 42 ^e
Lopinavir/ritonavir 400 mg/100 mg twice daily for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	18	↓ 58 (↓ 42 to ↓ 70)	↓ 63 (↓ 51 to ↓ 72)	↓ 65 (↓ 54 to ↓ 73)
Maraviroc 300 mg twice daily for 10 days	700 mg twice daily plus ritonavir 100 mg twice daily for 20 days	14	↓ 34 (↓ 25 to ↓41)	↓ 35 (↓ 29 to ↓ 41)	↓ 36 (↓ 27 to ↓ 43)
Maraviroc 300 mg twice daily for 10 days	1,400 mg once daily plus ritonavir 100 mg once daily for 20 days	14	↓ 29 (↓ 20 to ↓38)	↓30 (↓23 to ↓36)	↓ 15 (↓3 to ↓25)
Methadone 70 to 120 mg twice daily for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	19	↔ ^c	↔ ^c	↔ ^c
Nevirapine 200 mg twice daily for 2 weeks ^f	1,400 mg twice daily for 2 weeks	17	↓ 25 (↓ 37 to ↓ 10)	↓ 33 (↓ 45 to ↓ 20)	↓ 35 (↓ 50 to ↓ 15)
Nevirapine 200 mg twice daily. for 2 weeks ^f	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	17	↔	↓ 11 (↓ 23 to ↑ 3)	↓ 19 (↓ 32 to ↓ 4)
Phenytoin	700 mg twice	13			

300 mg once daily for 10 days	daily plus ritonavir 100 mg twice daily for 10 days		↔	↑ 20 (↑ 8 to ↑ 34)	↑ 19 (↑ 6 to ↑ 33)
Raltegravir 400 mg twice daily for 14 days	1,400 mg twice daily for 14 days (fasted)	14	↓ 27 (↓ 46 to ↔)	↓ 36 (↓ 53 to ↓ 13)	↓ 43 ^g (↓ 59 to ↓ 21)
	1,400 mg twice daily for 14 days ^h	14	↓ 15 (↓ 27 to ↓ 1)	↓ 17 (↓ 27 to ↓ 6)	↓ 32 ^g (↓ 53 to ↓ 1)
	700 mg twice daily plus ritonavir 100 mg twice daily for 14 days (fasted)	14	↓ 14 (↓ 39 to ↑ 20)	↓ 17 (↓ 38 to ↑ 12)	↓ 20 ^g (↓ 45 to ↑ 17)
	700 mg twice daily plus ritonavir 100 mg twice daily for 14 days ^h	12	↓ 25 (↓ 42 to ↓ 2)	↓ 25 (↓ 44 to ↔)	↓ 33 ^g (↓ 52 to ↓ 7)
Raltegravir 400 mg twice daily. for 14 days	1,400 mg once daily for 14 days (fasted)	13	↓ 18 (↓ 34 to ↔)	↓ 24 (↓ 41 to ↔)	↓ 50 ^g (↓ 64 to ↓ 31)
	1,400 mg once daily for 14 days ^h	14	↑ 27 (↓ 1 to ↑ 62)	↑ 13 (↓ 7 to ↑ 38)	↓ 17 ^g (↓ 45 to ↑ 26)
Ranitidine 300 mg single dose (administered 1 hour before fosamprenavir)	1,400 mg single dose	30	↓ 51 (↓ 43 to ↓ 58)	↓ 30 (↓ 22 to ↓ 37)	↔ (↓ 19 to ↑ 21)
Rifabutin 150 mg every other day for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	15	↑ 36 ^c (↑ 18 to ↑ 55)	↑ 35 ^c (↑ 17 to ↑ 56)	↑ 17 ^c (↓ 1 to ↑ 39)
Tenofovir 300 mg once daily for 4 to 48 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 4 to 48 weeks	45	NA	NA	↔ ⁱ
Tenofovir 300 mg once daily for 4 to 48 weeks	1,400 mg once daily plus ritonavir 200 mg once daily for 4 to 48 weeks	60	NA	NA	↔ ⁱ

1. Concomitant medication is also shown in this column where appropriate.

2. Ritonavir C_{max} , AUC, and C_{min} increased by 63%, 45%, and 13%, respectively, compared with historical control.
3. Compared with historical control.
4. Subjects were receiving fosamprenavir calcium/ritonavir for 10 days prior to the 4-day treatment period with both ketoconazole and fosamprenavir calcium/ritonavir.
5. Compared with fosamprenavir calcium 700 mg/ritonavir 100 mg twice daily for 2 weeks.
6. Subjects were receiving nevirapine for at least 12 weeks prior to trial.
7. C_{last} (C_{12h} or C_{24h}).
8. Doses of fosamprenavir calcium and raltegravir were given with food on pharmacokinetic sampling days and without regard to food all other days.
9. $N = 18$ for C_{min} .
10. Compared with parallel control group.

↑ = Increase; ↓ = Decrease; ↔ = No change (↑ or ↓ less than or equal to 10%), NA = Not applicable.

Table 11. Drug Interactions: Pharmacokinetic Parameters for Amprenavir after Administration of AGENERASE in the Presence of the Coadministered Drug(s)

Coadministered Drug(s) and Dose(s)	Dose of AGENERASE ^a	n	% Change in Amprenavir Pharmacokinetic Parameters (90% CI)		
			C_{max}	AUC	C_{min}
Abacavir 300 mg twice daily for 2 to 3 weeks	900 mg twice daily for 2 to 3 weeks	4	↔ ^a	↔ ^a	↔ ^a
Clarithromycin 500 mg twice daily for 4 days	1,200 mg twice daily for 4 days	12	↑ 15 (↑ 1 to ↑ 31)	↑ 18 (↑ 8 to ↑ 29)	↑ 39 (↑ 31 to ↑ 47)
Delavirdine 600 mg twice daily for 10 days	600 mg twice daily for 10 days	9	↑ 40 ^b	↑ 130 ^b	↑ 125 ^b
Ethinyl estradiol/norethindrone 0.035 mg/1 mg for 1 cycle	1,200 mg twice daily for 28 days	10	↔	↓ 22 (↓ 35 to ↓ 8)	↓ 20 (↓ 41 to ↑ 8)
Indinavir 800 mg 3 times a day for 2 weeks (fasted)	750 or 800 mg 3 times a day for 2 weeks (fasted)	9	↑ 18 (↑ 13 to ↑ 58)	↑ 33 (↑ 2 to ↑ 73)	↑ 25 (↓ 27 to ↑ 116)

Ketoconazole 400-mg single dose	1,200-mg single dose	12	↓ 16 (↓ 25 to ↓ 6)	↑ 31 (↑ 20 to ↑ 42)	NA
Lamivudine 150-mg single dose	600-mg single dose	11	↔	↔	NA
Methadone 44 to 100 mg once daily for > 30 days	1,200 mg mg once daily for 10 days	16	↓ 27 ^c	↓ 30 ^c	↓ 25 ^c
Nelfinavir 750 mg 3 times a day for 2 weeks (fed)	750 or 800 mg 3 times a day for 2 weeks (fed)	6	↓ 14 (↓ 38 to ↑ 20)	↔	↑ 189 (↑ 52 to ↑ 448)
Rifabutin 300 mg once daily for 10 days	1,200 mg twice daily for 10 days	5	↔	↓ 15 (↓ 28 to 0)	↓ 15 (↓ 38 to ↑ 17)
Rifampin 300 mg once daily for 4 days	1,200 mg twice daily for 4 days	11	↓ 70 (↓ 76 to ↓ 62)	↓ 82 (↓ 84 to ↓ 78)	↓ 92 (↓ 95 to ↓ 89)
Saquinavir 800 mg 3 times a day for 2 weeks (fed)	750 or 800 mg 3 times a day for 2 weeks (fed)	7	↓ 37 (↓ 54 to ↓ 14)	↓ 32 (↓ 49 to ↓ 9)	↓ 14 (↓ 52 to ↑ 54)
Zidovudine 300-mg single dose	600-mg single dose	12	↔	↑ 13 (↓ 2 to ↑ 31)	NA

1. Compared with parallel control group.
2. Median percent change; confidence interval not reported.
3. Compared with historical data.

↑ = Increase; ↓ = Decrease; ↔ = No change (↑ or ↓ less than 10%); NA = C_{min} not calculated for single-dose trial.

Table 12. Drug Interactions: Pharmacokinetic Parameters for Coadministered Drug in the

Presence of Amprenavir after Administration of Fosamprenavir Calcium

Coadministered Drug(s) and Dose(s)	Dose of Fosamprenavir Calcium ^a	n	% Change in Pharmacokinetic Parameters of Coadministered Drug (90% CI)		
			C _{max}	AUC	C _{min}
Atazanavir 300 mg once daily for 10 days ^b	700 mg twice daily plus ritonavir 100 mg twice daily for 10 days	21	↓ 24 (↓ 39 to ↓ 6)	↓ 22 (↓ 34 to ↓ 9)	↔
Atorvastatin 10 mg once daily for 4 days	1,400 mg twice daily for 2 weeks	16	↑ 304 (↑ 205 to ↑ 437)	↑ 130 (↑ 100 to ↑ 164)	↓ 10 (↓ 27 to ↑ 12)
Atorvastatin 10 mg once daily for 4 days	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	16	↑ 184 (↑ 126 to ↑ 257)	↑ 153 (↑ 115 to ↑ 199)	↑ 73 (↑ 45 to ↑ 108)
Esomeprazole 20 mg once daily. for 2 weeks	1,400 mg twice daily for 2 weeks	25	↔	↑ 55 (↑ 39 to ↑ 73)	ND
Esomeprazole 20 mg once daily for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	23	↔	↔	ND
Ethinyl estradiol ^c 0.035 mg once daily for 21 days	700 mg twice daily plus ritonavir 100 mg twice daily for 21 days	25	↓ 28 (↓ 21 to ↓ 35)	↓ 37 (↓ 30 to ↓ 42)	ND
Dolutegravir 50 mg once daily	700 mg twice daily plus ritonavir 100 mg twice daily.	12	↓ 24 (↓ 8 to ↓ 37)	↓ 35 (↓ 22 to ↓ 46)	↓ 49 (↓ 37 to ↓ 59)
Ketoconazole ^d 200 mg once daily for 4 days	700 mg twice daily plus ritonavir 100 mg twice daily for 4 days	15	↑ 25 (↑ 0 to ↑ 56)	↑ 169 (↑ 108 to ↑ 248)	ND
Lopinavir/ritonavir ^e 533 mg/133 mg twice daily for 2 weeks	1,400 mg twice daily for 2 weeks	18	↔ ^f	↔ ^f	↔ ^f

Lopinavir/ritonavir ^e 400 mg/100 mg twice daily for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	18	↑ 30 (↓ 15 to ↑ 47)	↑ 37 (↓ 20 to ↑ 55)	↑ 52 (↓ 28 to ↑ 82)
Maraviroc 300 mg twice daily for 10 days	700 mg twice daily plus ritonavir 100 mg twice daily for 20 days	14	↑ 52 (↑ 27 to ↑ 82)	↑ 149 (↑ 119 to ↑ 182)	↑ 374 (↑ 303 to ↑ 457)
Maraviroc 300 mg once daily for 10 days	1,400 mg once daily plus ritonavir 100 mg once daily for 20 days	14	↑ 45 (↑ 20 to ↑ 74)	↑ 126 (↑ 99 to ↑ 158)	↑ 80 (↑ 53 to ↑ 113)
Methadone 70 to 120 mg once daily for 2 weeks	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	19	R-Methadone (active)		
			↓ 21 ^g (↓ 30 to ↓ 12)	↓ 18 ^g (↓ 27 to ↓ 8)	↓ 11 ^g (↓ 21 to ↑ 1)
			S-Methadone (inactive)		
Nevirapine 200 mg twice daily for 2 weeks ^h	1,400 mg twice daily for 2 weeks	17	↓ 43 ^g (↓ 49 to ↓ 37)	↓ 43 ^g (↓ 50 to ↓ 36)	↓ 41 ^g (↓ 49 to ↓ 31)
			↑ 25 (↑ 14 to ↑ 37)	↑ 29 (↑ 19 to ↑ 40)	↑ 34 (↑ 20 to ↑ 49)
Nevirapine 200 mg twice daily for 2 weeks ^h	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	17	↑ 13 (↑ 3 to ↑ 24)	↑ 14 (↑ 5 to ↑ 24)	↑ 22 (↑ 9 to ↑ 35)
Norethindrone ^c 0.5 mg once daily for 21 days	700 mg twice daily plus ritonavir 100 mg twice daily for 21 days	25	↓ 38 (↓ 32 to ↓ 44)	↓ 34 (↓ 30 to ↓ 37)	↓ 26 (↓ 20 to ↓ 32)
Phenytoin 300 mg once daily for 10 days	700 mg twice daily plus ritonavir 100 mg twice daily. for 10 days	14	↓ 20 (↓ 12 to ↓ 27)	↓ 22 (↓ 17 to ↓ 27)	↓ 29 (↓ 23 to ↓ 34)
Rifabutin 150 mg every other day for 2 weeks ⁱ	700 mg twice daily plus ritonavir 100 mg twice daily for 2 weeks	15	↓ 14 (↓ 28 to ↑ 4)	↔	↑ 28 (↑ 12 to ↑ 46)

(25-O-desacetyl rifabutin metabolite)		↑ 579 (↑ 479 to ↑ 698)	↑ 1,120 (↑ 965 to ↑ 1,300)	↑ 2,510 (↑ 1,910 to ↑ 3,300)
Rifabutin + 25-O-desacetyl rifabutin metabolite		NA	↑ 64 (↑ 46 to ↑ 84)	NA
Rosuvastatin 10 mg single dose	700 mg twice daily plus ritonavir 100 mg twice daily for 7 days	(↑ 45)	(↑ 8)	NA

1. Concomitant medication is also shown in this column where appropriate.
2. Comparison arm of atazanavir 300 mg once daily plus ritonavir 100 mg once daily for 10 days.
3. Administered as a combination oral contraceptive tablet: ethinyl estradiol 0.035 mg/norethindrone 0.5 mg.
4. Subjects were receiving fosamprenavir calcium/ritonavir for 10 days prior to the 4-day treatment period with both ketoconazole and fosamprenavir calcium/ritonavir.
5. Data represent lopinavir concentrations.
6. Compared with lopinavir 400 mg/ritonavir 100 mg twice daily for 2 weeks.
7. Dose normalized to methadone 100 mg. The unbound concentration of the active moiety, R-methadone, was unchanged.
8. Subjects were receiving nevirapine for at least 12 weeks prior to trial.
9. Comparison arm of rifabutin 300 mg once daily for 2 weeks. AUC is AUC_(0-48 h).

↑ = Increase; ↓ = Decrease; ↔ = No change (↑ or ↓ less than 10%); ND = Interaction cannot be determined as C_{min} was below the lower limit of quantitation.

Table 13. Drug Interactions: Pharmacokinetic Parameters for Coadministered Drug in the Presence of Amprenavir after Administration of AGENERASE

Coadministered Drug(s) and Dose(s)	Dose of AGENERASE	n	% Change in Pharmacokinetic Parameters of Coadministered Drug (90% CI)		
			C _{max}	AUC	C _{min}
Abacavir 300 mg twice daily for 2 to 3	900 mg twice daily for 2 to 3 weeks	4	↔ ^a	↔ ^a	↔ ^a

weeks					
Clarithromycin 500 mg twice daily for 4 days	1,200 mg twice daily for 4 days	12	↓ 10 (↓ 24 to ↑ 7)	↔	↔
Delavirdine 600 mg twice daily for 10 days	600 mg twice daily for 10 days	9	↓ 47 ^b	↓ 61 ^b	↓ 88 ^b
Ethinyl estradiol 0.035 mg for 1 cycle	1,200 mg twice daily for 28 days	10	↔	↔	↑ 32 (↓ 3 to ↑ 79)
Indinavir 800 mg 3 times a day for 2 weeks (fasted)	750 mg or 800 mg 3 times a day for 2 weeks (fasted)	9	↓ 22 ^a	↓ 38 ^a	↓ 27 ^a
Ketoconazole 400 mg single dose	1,200 mg single dose	12	↑ 19 (↑ 8 to ↑ 33)	↑ 44 (↑ 31 to ↑ 59)	NA
Lamivudine 150 mg single dose	600 mg single dose	11	↔	↔	NA
Methadone 44 to 100 mg once daily for > 30 days	1,200 mg twice daily for 10 days	16	R-Methadone (active)		
			↓ 25 (↓ 32 to ↓ 18)	↓ 13 (↓ 21 to ↓ 5)	↓ 21 (↓ 32 to ↓ 9)
			S-Methadone (inactive)		
			↓ 48 (↓ 55 to ↓ 40)	↓ 40 (↓ 46 to ↓ 32)	↓ 53 (↓ 60 to ↓ 43)
Nelfinavir 750 mg 3 times a day for 2 weeks (fed)	750 mg or 800 mg 3 times a day for 2 weeks (fed)	6	↑ 12 ^a	↑ 15 ^a	↑ 14 ^a
Norethindrone 1 mg for 1 cycle	1,200 mg twice daily for 28 days	10	↔	↑ 18 (↑ 1 to ↑ 38)	↑ 45 (↑ 13 to ↑ 88)
Rifabutin	1,200 mg twice daily for 10	5	↑ 119 (↑ 82 to ↑ 164)	↑ 193 (↑ 156 to ↑ 235)	↑ 271 (↑ 171 to ↑

300 mg once daily for 10 days	days				409)
Rifampin 300 mg once daily for 4 days	1,200 mg twice daily for 4 days	11	↔	↔	ND
Saquinavir 800 mg 3 times a day for 2 weeks (fed)	750 mg or 800 mg 3 times a day for 2 weeks (fed)	7	↑ 21 ^a	↓ 19 ^a	↓ 48 ^a
Zidovudine 300 mg single dose	600 mg single dose	12	↑ 40 (↑ 14 to ↑ 71)	↑ 31 (↑ 19 to ↑ 45)	NA

1. Compared with historical data.
2. Median percent change; confidence interval not reported.

↑ = Increase; ↓ = Decrease; ↔ = No change (↑ or ↓ less than 10%); NA = C_{min} not calculated for single-dose trial; ND = Interaction cannot be determined as C_{min} was below the lower limit of quantitation.

12.4 Microbiology

Mechanism of Action

Fosamprenavir is a prodrug that is rapidly hydrolyzed to amprenavir by cellular phosphatases in the gut epithelium as it is absorbed. Amprenavir is an inhibitor of HIV-1 protease. Amprenavir binds to the active site of HIV-1 protease and thereby prevents the processing of viral Gag and Gag-Pol polyprotein precursors, resulting in the formation of immature non-infectious viral particles.

Antiviral Activity

Fosamprenavir has little or no antiviral activity in cell culture. The antiviral activity of amprenavir was evaluated against HIV-1 IIIB in both acutely and chronically infected lymphoblastic cell lines (MT-4, CEM-CCRF, H9) and in peripheral blood lymphocytes in cell culture. The 50% effective concentration (EC_{50}) of amprenavir ranged from 0.012 to 0.08 μM in acutely infected cells and was 0.41 μM in chronically infected cells (1 μM = 0.50 mcg per mL). The median EC_{50} value of amprenavir against HIV-1 isolates from clades A to G was 0.00095 μM in peripheral blood mononuclear cells (PBMCs). Similarly, the EC_{50} values for amprenavir against monocytes/macrophage tropic HIV-1 isolates (clade B) ranged from 0.003 to 0.075 μM in monocyte/macrophage cultures. The EC_{50} values of amprenavir against HIV-2 isolates grown in PBMCs were higher than those for HIV-1 isolates, and ranged from 0.003 to 0.11 μM . The anti-HIV-1 activity of amprenavir was not antagonistic in combination with the nucleoside reverse transcriptase inhibitors (NRTIs); abacavir, didanosine, lamivudine, stavudine, tenofovir, and zidovudine; the non-nucleoside reverse transcriptase inhibitors (NNRTIs) delavirdine, efavirenz, and nevirapine; the protease inhibitors (PIs) atazanavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir; and the gp41 fusion inhibitor enfuvirtide. These drug combinations have not been adequately studied in humans.

Resistance

HIV-1 isolates with decreased susceptibility to amprenavir have been selected in cell culture and obtained from subjects treated with fosamprenavir. Genotypic analysis of isolates from treatment-naive subjects failing amprenavir-containing regimens showed substitutions in the HIV-1 protease resulting in amino acid substitutions primarily at positions V32I, M46I/L, I47V, I50V, I54L/M, and I84V, as well as substitutions in the p7/p1 and p1/p6 Gag and Gag-Pol polyprotein precursor cleavage sites. Some of these amprenavir resistance-associated substitutions have also been detected in HIV-1 isolates from antiretroviral-naive subjects treated with fosamprenavir calcium. Of the 488 antiretroviral-naive subjects treated with fosamprenavir calcium 1,400 mg twice daily or fosamprenavir calcium 1,400 mg plus ritonavir 200 mg once daily in Trials APV30001 and APV30002, respectively, isolates from 61 subjects (29 receiving fosamprenavir calcium and 32 receiving fosamprenavir calcium/ritonavir) with virologic failure (plasma HIV-1 RNA greater than 1,000 copies per mL on 2 occasions on or after Week 12) were genotyped. Isolates from 5 of the 29 antiretroviral-naive subjects (17%) receiving fosamprenavir calcium without ritonavir in Trial APV30001 had evidence of genotypic resistance to amprenavir: I54L/M (n = 2), I54L + L33F (n = 1), V32I + I47V (n = 1), and M46I + I47V (n = 1). No amprenavir resistance-associated substitutions were detected in isolates from antiretroviral-naive subjects treated with fosamprenavir calcium/ritonavir for 48 weeks in Trial APV30002. However, the M46I and I50V substitutions were detected in isolates from 1 virologic failure subject receiving fosamprenavir calcium/ritonavir once daily at Week 160 (HIV-1 RNA greater than 500 copies per mL). Upon retrospective analysis of stored samples using an ultrasensitive assay, these resistant substitutions were traced back to Week 84 (76 weeks prior to clinical virologic failure).

Cross-Resistance

Varying degrees of cross-resistance among HIV-1 protease inhibitors have been observed. An association between virologic response at 48 weeks (HIV-1 RNA level less than 400 copies per mL) and protease inhibitor-resistance substitutions detected in baseline HIV-1 isolates from protease inhibitor-experienced subjects receiving fosamprenavir calcium/ritonavir twice daily (n = 88), or lopinavir/ritonavir twice daily (n = 85) in Trial APV30003 is shown in Table 14. The majority of subjects had previously received either one (47%) or 2 protease inhibitors (36%), most commonly nelfinavir (57%) and indinavir (53%). Out of 102 subjects with baseline phenotypes receiving twice-daily fosamprenavir calcium/ritonavir, 54% (n = 55) had resistance to at least one protease inhibitor, with 98% (n = 54) of those having resistance to nelfinavir. Out of 97 subjects with baseline phenotypes in the lopinavir/ritonavir arm, 60% (n = 58) had resistance to at least one protease inhibitor, with 97% (n = 56) of those having resistance to nelfinavir.

Table 14. Responders at Trial Week 48 by Presence of Baseline Protease Inhibitor Resistance-Associated Substitutions^a

Protease Inhibitor Resistance-Associated Substitutions^b	Fos amprenavir Calcium/Ritonavir Twice Daily (n = 88)		Lopinavir/Ritonavir Twice Daily (n = 85)	
D30N	21/22	95%	17/19	89%
N88D/S	20/22	91%	12/12	100%
L90M	16/31	52%	17/29	59%
M46I/L	11/22	50%	12/24	50%
V82A/F/T/S	2/9	22%	6/17	35%
I54V	2/11	18%	6/11	55%
I84V	1/6	17%	2/5	40%

1. Results should be interpreted with caution because the subgroups were small.
2. Most subjects had greater than 1 protease inhibitor resistance-associated substitution at baseline.

The virologic response based upon baseline phenotype was assessed. Baseline isolates from protease

inhibitor-experienced subjects responding to fosamprenavir calcium/ritonavir twice daily had a median shift in susceptibility to amprenavir relative to a standard wild-type reference strain of 0.7 (range: 0.1 to 5.4, n = 62), and baseline isolates from individuals failing therapy had a median shift in susceptibility of 1.9 (range: 0.2 to 14, n = 29). Because this was a select patient population, these data do not constitute definitive clinical susceptibility break points. Additional data are needed to determine clinically relevant break points for fosamprenavir calcium.

Isolates from 15 of the 20 subjects receiving twice-daily fosamprenavir calcium/ritonavir up to Week 48 and experiencing virologic failure/ongoing replication were subjected to genotypic analysis. The following amprenavir resistance-associated substitutions were found either alone or in combination: V32I, M46I/L, I47V, I50V, I54L/M, and I84V. Isolates from 4 of the 16 subjects continuing to receive twice-daily fosamprenavir calcium/ritonavir up to Week 96 who experienced virologic failure underwent genotypic analysis. Isolates from 2 subjects contained amprenavir resistance-associated substitutions: V32I, M46I, and I47V in 1 isolate and I84V in the other.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In long-term carcinogenicity studies, fosamprenavir was administered orally for up to 104 weeks at doses of 250, 400, or 600 mg per kg per day in mice and at doses of 300, 825, or 2,250 mg per kg per day in rats. Exposures at these doses were 0.3- to 0.7-fold (mice) and 0.7- to 1.4-fold (rats) those in humans given 1,400 mg twice daily of fosamprenavir alone, and 0.2- to 0.3-fold (mice) and 0.3- to 0.7-fold (rats) those in humans given 1,400 mg once daily of fosamprenavir plus 200 mg ritonavir once daily. Exposures in the carcinogenicity studies were 0.1- to 0.3-fold (mice) and 0.3- to 0.6-fold (rats) those in humans given 700 mg of fosamprenavir plus 100 mg ritonavir twice daily. There was an increase in hepatocellular adenomas and hepatocellular carcinomas at all doses in male mice and at 600 mg per kg per day in female mice, and in hepatocellular adenomas and thyroid follicular cell adenomas at all doses in male rats, and at 825 mg per kg per day and 2,250 mg per kg per day in female rats. The relevance of the hepatocellular findings in the rodents for humans is uncertain. Repeat-dose studies with fosamprenavir in rats produced effects consistent with enzyme induction, which predisposes rats, but not humans, to thyroid neoplasms. In addition, in rats only there was an increase in interstitial cell hyperplasia at 825 mg per kg per day and 2,250 mg per kg per day, and an increase in uterine endometrial adenocarcinoma at 2,250 mg per kg per day. The incidence of endometrial findings was slightly increased over concurrent controls, but was within background range for female rats. The relevance of the uterine endometrial adenocarcinoma findings in rats for humans is uncertain.

Fosamprenavir was not mutagenic or genotoxic in a battery of *in vitro* and *in vivo* assays. These assays included bacterial reverse mutation (Ames), mouse lymphoma, rat micronucleus, and chromosome aberrations in human lymphocytes.

The effects of fosamprenavir on fertility and general reproductive performance were investigated in male (treated for 4 weeks before mating) and female rats (treated for 2 weeks before mating through Postpartum Day 6) that received doses of 300, 820, or 2,240 mg per kg per day. Systemic exposures ($AUC_{0-24\text{ h}}$) to amprenavir in these studies were 3 (males) to 4 (females) times higher than exposures in humans following administration of the MRHD of fosamprenavir alone or similar to those seen in humans following administration of fosamprenavir in combination with ritonavir. Fosamprenavir did not impair mating or fertility of male or female rats and did not affect the development and maturation of sperm from treated rats.

14 CLINICAL STUDIES

14.1 Therapy-Naive Adult Trials

APV30001

A randomized, open-label trial evaluated treatment with fosamprenavir calcium tablets (1,400 mg twice daily) versus nelfinavir (1,250 mg twice daily) in 249 antiretroviral treatment-naive subjects. Both groups of subjects also received abacavir (300 mg twice daily) and lamivudine (150 mg twice daily).

The mean age of the subjects in this trial was 37 years (range: 17 to 70 years); 69% of the subjects were male, 20% were CDC Class C (AIDS), 24% were white, 32% were black, and 44% were Hispanic. At baseline, the median CD4+ cell count was 212 cells per mm³ (range: 2 to 1,136 cells per mm³; 18% of subjects had a CD4+ cell count of less than 50 cells per mm³ and 30% were in the range of 50 to less than 200 cells per mm³). Baseline median HIV-1 RNA was 4.83 log₁₀ copies per mL (range: 1.69 to 7.41 log₁₀ copies per mL; 45% of subjects had greater than 100,000 copies per mL).

The outcomes of randomized treatment are provided in Table 15.

Table 15. Outcomes of Randomized Treatment through Week 48 (APV30001)

Outcome (Rebound or discontinuation = failure)	Fosamprenavir Calcium 1,400 mg Twice Daily (n = 166)	Nelfinavir 1,250 mg Twice Daily (n = 83)
Responder ^a	66% (57%)	52% (42%)
Virologic failure	19%	32%
Rebound	16%	19%
Never suppressed through Week 48	3%	13%
Clinical progression	1%	1%
Death	0%	1%
Discontinued due to adverse reactions	4%	2%
Discontinued due to other reasons ^b	10%	10%

1. Subjects achieved and maintained confirmed HIV-1 RNA less than 400 copies per mL (less than 50 copies per mL) through Week 48 (Roche AMPLICOR HIV-1 MONITOR Assay Version 1.5).
2. Includes consent withdrawn, lost to follow up, protocol violations, those with missing data, and other reasons.

Treatment response by viral load strata is shown in Table 16.

Table 16. Proportions of Responders through Week 48 by Screening Viral Load (APV30001)

Screening Viral Load HIV-1 RNA (copies/mL)	Fosamprenavir Calcium 1,400 mg Twice Daily		Nelfinavir 1,250 mg Twice Daily	
	< 400 copies/mL	n	< 400 copies/mL	n
≤ 100,000	65%	93	65%	46
> 100,000	67%	73	36%	37

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 201 cells per mm³ in the group receiving fosamprenavir calcium and 216 cells per mm³ in the nelfinavir group.

APV30002

A randomized, open-label trial evaluated treatment with fosamprenavir calcium tablets (1,400 mg once daily) plus ritonavir (200 mg once daily) versus nelfinavir (1,250 mg twice daily) in 649 treatment-naïve subjects. Both treatment groups also received abacavir (300 mg twice daily) and lamivudine (150 mg twice daily).

The mean age of the subjects in this trial was 37 years (range: 18 to 69 years); 73% of the subjects were male, 22% were CDC Class C, 53% were white, 36% were black, and 8% were Hispanic. At baseline, the median CD4+ cell count was 170 cells per mm³ (range: 1 to 1,055 cells per mm³; 20% of subjects had a CD4+ cell count of less than 50 cells per mm³ and 35% were in the range of 50 to less than 200 cells per mm³). Baseline median HIV-1 RNA was 4.81 log₁₀ copies per mL (range: 2.65 to 7.29 log₁₀ copies per mL; 43% of subjects had greater than 100,000 copies per mL).

The outcomes of randomized treatment are provided in Table 17.

Table 17. Outcomes of Randomized Treatment through Week 48 (APV30002)

Outcome (Rebound or discontinuation = failure)	Fosamprenavir Calcium 1,400 mg/Ritonavir 200 mg Once Daily (n = 322)	Nelfinavir 1,250 mg Twice Daily (n = 327)
Responder ^a	69% (58%)	68% (55%)
Virologic failure	6%	16%
Rebound	5%	8%
Never suppressed through Week 48	1%	8%
Death	1%	0%
Discontinued due to adverse reactions	9%	6%
Discontinued due to other	15%	10%

reasons ^b		
----------------------	--	--

1. Subjects achieved and maintained confirmed HIV-1 RNA less than 400 copies per mL (less than 50 copies per mL) through Week 48 (Roche AMPLICOR HIV-1 MONITOR Assay Version 1.5).
2. Includes consent withdrawn, lost to follow up, protocol violations, those with missing data, and other reasons.

Treatment response by viral load strata is shown in Table 18.

Table 18. Proportions of Responders through Week 48 by Screening Viral Load (APV30002)

Screening Viral Load HIV-1 RNA (copies/mL)	Fosamprenavir Calcium 1,400 mg/Ritonavir 200 mg Once Daily		Nelfinavir 1,250 mg Twice Daily	
	< 400 copies/mL	n	< 400 copies/mL	n
≤ 100,000	72%	197	73%	194
> 100,000	66%	125	64%	133

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 203 cells per mm³ in the group receiving fosamprenavir calcium and 207 cells per mm³ in the nelfinavir group.

14.2 Protease Inhibitor-Experienced Adult Trials

APV30003

A randomized, open-label, multicenter trial evaluated 2 different regimens of fosamprenavir calcium plus ritonavir (fosamprenavir calcium tablets 700 mg twice daily plus ritonavir 100 mg twice daily or fosamprenavir calcium tablets 1,400 mg once daily plus ritonavir 200 mg once daily) versus lopinavir/ritonavir (400 mg/100 mg twice daily) in 315 subjects who had experienced virologic failure to 1 or 2 prior protease inhibitor-containing regimens.

The mean age of the subjects in this trial was 42 years (range: 24 to 72 years); 85% were male, 33% were CDC Class C, 67% were white, 24% were black, and 9% were Hispanic. The median CD4+ cell count at baseline was 263 cells per mm³ (range: 2 to 1,171 cells per mm³). Baseline median plasma HIV-1 RNA level was 4.14 log₁₀ copies per mL (range: 1.69 to 6.41 log₁₀ copies per mL).

The median durations of prior exposure to NRTIs were 257 weeks for subjects receiving fosamprenavir calcium/ritonavir twice daily (79% had greater than or equal to 3 prior NRTIs) and 210 weeks for subjects receiving lopinavir/ritonavir (64% had greater than or equal to 3 prior NRTIs). The median durations of prior exposure to protease inhibitors were 149 weeks for subjects receiving fosamprenavir calcium/ritonavir twice daily (49% received greater than or equal to 2 prior protease inhibitors) and 130 weeks for subjects receiving lopinavir/ritonavir (40% received greater than or equal to 2 prior protease inhibitors).

The time-averaged changes in plasma HIV-1 RNA from baseline (AAUCMB) at 48 weeks (the endpoint on which the trial was powered) were -1.4 log₁₀ copies per mL for twice-daily fosamprenavir calcium/ritonavir and -1.67 log₁₀ copies per mL for the lopinavir/ritonavir group.

The proportions of subjects who achieved and maintained confirmed HIV-1 RNA less than 400 copies per mL (secondary efficacy endpoint) were 58% with twice-daily fosamprenavir calcium/ritonavir and 61% with lopinavir/ritonavir (95% CI for the difference: -16.6, 10.1). The proportions of subjects with HIV-1 RNA less than 50 copies per mL with twice-daily fosamprenavir calcium/ritonavir and with lopinavir/ritonavir were 46% and 50%, respectively (95% CI for the difference: -18.3, 8.9). The proportions of subjects who were virologic failures were 29% with twice-daily fosamprenavir calcium/ritonavir and 27% with lopinavir/ritonavir.

The frequency of discontinuations due to adverse events and other reasons, and deaths were similar between treatment arms.

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 81 cells per mm³ with twice-daily fosamprenavir calcium/ritonavir and 91 cells per mm³ with lopinavir/ritonavir.

This trial was not large enough to reach a definitive conclusion that fosamprenavir calcium/ritonavir and lopinavir/ritonavir are clinically equivalent.

Once-daily administration of fosamprenavir calcium plus ritonavir is not recommended for protease inhibitor-experienced patients. Through Week 48, 50% and 37% of subjects receiving fosamprenavir calcium 1,400 mg plus ritonavir 200 mg once daily had plasma HIV-1 RNA less than 400 copies per mL and less than 50 copies per mL, respectively.

14.3 Pediatric Trials

Three open-label trials in pediatric subjects aged at least 4 weeks to 18 years were conducted. In one trial (APV29005), twice-daily dosing regimens (fosamprenavir calcium with or without ritonavir) were evaluated in combination with other antiretroviral agents in pediatric subjects aged 2 to 18 years. In a second trial (APV20002), twice-daily dosing regimens (fosamprenavir calcium with ritonavir) were evaluated in combination with other antiretroviral agents in pediatric subjects aged at least 4 weeks to younger than 2 years. A third trial (APV20003) evaluated once-daily dosing of fosamprenavir calcium with ritonavir; the pharmacokinetic data from this trial did not support a once-daily dosing regimen in any pediatric patient population.

APV29005

Fosamprenavir calcium: Twenty (18 therapy-naive and 2 therapy-experienced) pediatric subjects received fosamprenavir calcium oral suspension without ritonavir twice daily. At Week 24, 65% (13 of 20) achieved HIV-1 RNA less than 400 copies per mL, and the median increase from baseline in CD4+ cell count was 350 cells per mm³.

Fosamprenavir calcium plus Ritonavir: Forty-nine protease inhibitor-naive and 40 protease inhibitor-experienced pediatric subjects received fosamprenavir calcium oral suspension or tablets with ritonavir twice daily. At Week 24, 71% of protease inhibitor-naive (35 of 49) and 55% of protease inhibitor-experienced (22 of 40) subjects achieved HIV-1 RNA less than 400 copies per mL; median increases from baseline in CD4+ cell counts were 184 cells per mm³ and 150 cells per mm³ in protease inhibitor-naive and experienced subjects, respectively.

APV20002

Fifty-four pediatric subjects (49 protease inhibitor-naive and 5 protease inhibitor-experienced) received fosamprenavir calcium oral suspension with ritonavir twice daily. At Week 24, 72% of subjects achieved HIV-1 RNA less than 400 copies per mL. The median increases from baseline in CD4+ cell counts were 400 cells per mm³ in subjects aged at least 4 weeks to younger than 6 months and 278 cells per mm³ in subjects aged 6 months to 2 years.

16 HOW SUPPLIED/STORAGE AND HANDLING

Fosamprenavir calcium tablets, 700 mg, are pink colored, coated, oval-shaped tablets with “**RJ47**” imprinted on one side with black ink and plain on the other side. They are supplied as follows-

NDC 63304-583-30 Bottles of 30 with child-resistant closure

NDC 63304-583-60 Bottles of 60 with child-resistant closure

NDC 63304-583-01 Bottles of 100

Store at 20° C to 25° C (68° F to 77° F) [See USP Controlled Room Temperature].

Keep bottle tightly closed. Protect from moisture.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Drug Interactions

A statement to patients and healthcare providers is included on the product's bottle label: **ALERT: Find out about medicines that should NOT be taken with fosamprenavir calcium tablets.**

Fosamprenavir calcium tablets may interact with many drugs; therefore, advise patients to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products, particularly St. John's wort.

Advise patients receiving PDE5 inhibitors that they may be at an increased risk of PDE5 inhibitor-associated adverse events, including hypotension, visual changes, and priapism, and should promptly report any symptoms to their healthcare provider [see *Contraindications (4), Warnings and Precautions (5.1), Drug Interactions (7)*].

Contraception

Instruct patients receiving combined hormonal contraception to use an effective alternative contraceptive method or an additional barrier method during therapy with fosamprenavir calcium tablets because hormonal levels may decrease, and if used in combination with fosamprenavir calcium tablets and ritonavir, liver enzyme elevations may occur [see *Drug Interactions (7.2), Use in Specific Populations (8.3)*].

Sulfa Allergy

Advise patients to inform their healthcare provider if they have a sulfa allergy. The potential for cross-sensitivity between drugs in the sulfonamide class and fosamprenavir is unknown [see *Warnings and Precautions (5.3)*].

Severe Skin Reactions

Advise patients that skin reactions ranging from mild to severe, including Stevens-Johnson Syndrome, have been reported with fosamprenavir calcium tablets. Advise patients to discontinue fosamprenavir calcium tablets immediately for severe or life-threatening skin reactions or for moderate rashes accompanied by systemic symptoms [see *Warnings and Precautions (5.2), Adverse Reactions (6)*].

Hepatic Toxicity

Advise patients that it is recommended to have laboratory testing before and during therapy as patients with underlying hepatitis B or C or marked elevations of transaminases prior to treatment may be at increased risk for developing or worsening transaminase elevations with use of fosamprenavir calcium tablets, particularly at higher than recommended doses which should not be used. [see *Warnings and Precautions (5.4)*].

Immune Reconstitution Syndrome

Advise patients to inform their healthcare provider immediately of any signs or symptoms of infection as inflammation from previous infection may occur soon after combination antiretroviral therapy, including when fosamprenavir calcium tablets is started [see *Warnings and Precautions (5.6)*].

Increase in Body Fat

Inform patients that an increase of body fat may occur in patients receiving protease inhibitors, including fosamprenavir calcium tablets, and that the cause and long-term health effects of these conditions are not known at this time [see *Warnings and Precautions (5.7)*].

Lipid Elevations

Advise patients that it is recommended to have laboratory testing before and during therapy as increases in the concentration of triglycerides and cholesterol have been reported with use of fosamprenavir calcium tablets [see *Warnings and Precautions (5.8), Adverse Reactions (6)*].

Pregnancy Registry

Advise patients that there is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to fosamprenavir calcium tablets during pregnancy [see *Use in Specific Populations (8.1)*].

Lactation

Instruct women with HIV-1 infection not to breastfeed because HIV-1 can be passed to the baby in the breast milk [see *Use in Specific Populations (8.2)*].

Missed Dose

Instruct patients that if they miss a dose of fosamprenavir calcium tablets, to take it as soon as they remember. Advise patients not to double their next dose or take more than the prescribed dose [see *Dosage and Administration (2)*].

The brands listed are trademarks of their respective owners.

Manufactured by:

Ohm Laboratories Inc.

New Brunswick, NJ 08901

Distributed by:

Sun Pharmaceutical Industries, Inc.

Cranbury, NJ 08512

May 2019 FDA-05

Patient Medication Information

PATIENT INFORMATION

FOSAMPRENAVIR (foss-am-PREN-ah-ver) CALCIUM TABLETS

Rx only

What is the most important information I should know about fosamprenavir calcium tablets?

Fosamprenavir calcium tablets can interact with other medicines and cause serious side effects. It is important to know the medicines that should not be taken with fosamprenavir calcium tablets. See the section “Who should not take fosamprenavir calcium tablets?”

fosamprenavir calcium tablets can cause serious side effects, including:

- **Severe skin reactions.** fosamprenavir calcium tablets may cause severe or life-threatening skin reactions or rash.

If you get a rash with any of the following symptoms, stop taking fosamprenavir calcium tablets and call your healthcare provider or get medical help right away:

-
- hives or sores in your mouth, or your skin or blisters and peels
 - swelling of your face, eyes, lips, tongue, throat

- trouble swallowing or breathing
-

For more information about side effects, see “What are the possible side effects of fosamprenavir calcium tablets?”

What are fosamprenavir calcium tablets?

Fosamprenavir calcium tablets are prescription medicines that are used together with other antiretroviral medicines to treat human immunodeficiency virus 1 (HIV-1) infection.

HIV-1 is the virus that causes Acquired Immune Deficiency Syndrome (AIDS).

It is not known if fosamprenavir calcium tablets are safe and effective in children younger than 4 weeks of age.

Do not take fosamprenavir calcium tablets if you:

- **are allergic** to amprenavir, fosamprenavir calcium, or any of the ingredients in fosamprenavir calcium tablets. See the end of this leaflet for a complete list of ingredients in fosamprenavir calcium tablets.
- **take any of the following medicines:**
 - alfuzosin
 - rifampin
 - ergot including:
 - dihydroergotamine mesylate
 - ergonovine
 - ergotamine tartrate
 - methylergonovine
 - cisapride
 - St. John’s wort (*Hypericum perforatum*)
 - lomitapide
 - lovastatin
 - simvastatin
 - pimozide
 - delavirdine mesylate
 - sildenafil (REVATIO), for treatment of pulmonary arterial hypertension
 - triazolam
 - midazolam, when taken by mouth

Serious problems can happen if you or your child take any of the medicines listed above with fosamprenavir calcium tablets.

If you are taking fosamprenavir calcium tablets with ritonavir, do not take the following medicines:

- flecainide
- propafenone
- lurasidone

Before taking fosamprenavir calcium tablets, tell your healthcare provider about all of your medical conditions, including if you:

- are allergic to medicines that contain sulfa.

- have or have had liver problems, including hepatitis B or C virus infection.
 - have kidney problems.
 - have high blood sugar (diabetes).
 - have hemophilia.
 - have high cholesterol.
 - are pregnant or plan to become pregnant. It is not known if fosamprenavir calcium tablets will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant during treatment with fosamprenavir calcium tablets.
 - fosamprenavir calcium tablets may reduce how well hormonal contraceptives (birth control pills) work. Females who may become pregnant should use a different form of birth control or an additional barrier method of birth control during treatment with fosamprenavir calcium tablets.
- Pregnancy Registry.** There is a pregnancy registry for women who take fosamprenavir calcium tablets during pregnancy. The purpose of the registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.
- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take fosamprenavir calcium tablets.**
 - You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.
 - It is not known if fosamprenavir calcium tablets can pass to your baby in your breast milk.
 - Talk with your healthcare provider about the best way to feed your baby.
- Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines interact with fosamprenavir calcium tablets. **Keep a list of your medicines to show your healthcare provider and pharmacist.**
- You can ask your healthcare provider or pharmacist for a list of medicines that interact with fosamprenavir calcium tablets.
 - **Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take fosamprenavir calcium tablets with other medicines.

How should I take fosamprenavir calcium tablets?

- **Take fosamprenavir calcium tablets exactly as your healthcare provider tells you to take it.**
- If you miss a dose of fosamprenavir calcium tablets, take it as soon as you remember. Do not take 2 doses at the same time or take more than your healthcare provider tells you to take.
- Stay under the care of a healthcare provider during treatment with fosamprenavir calcium tablets.
- If your child is taking fosamprenavir calcium tablets, your child's healthcare provider will decide the right dose based on your child's weight.
- fosamprenavir calcium tablets may be taken with or without food.
- Do not run out of fosamprenavir calcium tablets. The virus in your blood may increase and the virus may become harder to treat. When your supply starts to run low, get more from your healthcare provider or pharmacy.
- If you take too much fosamprenavir calcium tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

What are the possible side effects of fosamprenavir calcium tablets?

Fosamprenavir calcium tablets may cause serious side effects including:

- **See "What is the most important information I should know about fosamprenavir calcium tablets?"**
- **Liver problems.** Your healthcare provider should do blood tests before and during your

treatment with fosamprenavir calcium tablets to check your liver function. Some people with liver problems, including hepatitis B or C, may have an increased risk of developing worsening liver problems during treatment with fosamprenavir calcium tablets.

- **Diabetes and high blood sugar (hyperglycemia).** Some people who take protease inhibitors, including fosamprenavir calcium tablets, can get high blood sugar, develop diabetes, or your diabetes can get worse. Tell your healthcare provider if you notice an increase in thirst or urinate often during treatment with fosamprenavir calcium tablets.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Call your healthcare provider right away if you start having new symptoms after starting your HIV-1 medicine.
- **Increase in body fat.** An increase in body fat can happen in people who take protease inhibitors, including fosamprenavir calcium tablets. The exact cause and long-term health effects of these conditions are not known.
- **Changes in blood tests.** Some people have changes in blood tests while taking fosamprenavir calcium tablets. These include an increase in liver function tests, blood fat levels (cholesterol and triglycerides) and decrease in red blood cells. Your healthcare provider should do regular blood tests before and during your treatment with fosamprenavir calcium tablets.
- **Increased bleeding problems in some people with hemophilia.** Some people with hemophilia have increased bleeding with protease inhibitors, including fosamprenavir calcium tablets.
- **Kidney stones.** Some people have developed kidney stones during treatment with fosamprenavir calcium tablets. Tell your healthcare provider right away if you develop any of the following signs or symptoms of kidney stones:
 - pain in your side
 - blood in your urine
 - pain when you urinate

The most common side effects of fosamprenavir calcium tablets in adults include:

- nausea
 - vomiting
 - rash
 - diarrhea
 - headache
-

The most common side effects of fosamprenavir calcium tablets in children include vomiting and decrease in white blood cells.

These are not all the possible side effects of fosamprenavir calcium tablets.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store fosamprenavir calcium tablets?

- Store fosamprenavir calcium tablets at room temperature between 68° F to 77° F (20° C to 25° C).
- Keep the bottle of fosamprenavir calcium tablets tightly closed. Protect from moisture.

Keep fosamprenavir calcium tablets and all medicines out of the reach of children.

General information about the safe and effective use of fosamprenavir calcium tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet.

Do not use fosamprenavir calcium tablets for a condition for which it was not prescribed. Do not give fosamprenavir calcium tablets to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about fosamprenavir calcium tablets that is written for health professionals.

What are the ingredients in fosamprenavir calcium tablets?

Active ingredient: fosamprenavir calcium (amorphous).

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, and povidone. The tablet coating contains the inactive ingredients ethylcellulose, ferric oxide red, hypromellose, talc, and titanium dioxide. The imprinting ink contains the inactive ingredients ferric oxide black, propylene glycol, and shellac.

This Patient Information has been approved by the U.S. Food and Drug Administration.

The brands listed are trademarks of their respective owners.

Manufactured by:

Ohm Laboratories Inc.

New Brunswick, NJ 08901

Distributed by:

Sun Pharmaceutical Industries, Inc.

Cranbury, NJ 08512

May 2019 FDA-05

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

NDC 63304-583-60

Fosamprenavir Calcium Tablets

700 mg*

ALERT: Find out about medicines that should NOT be taken with Fosamprenavir Calcium Tablets.

Note to Pharmacist: Do not cover ALERT box with pharmacy label.

Rx only

60 Tablets ohm®

Manufactured by:
Ohm Laboratories, Inc.
New Brunswick, NJ 08901
Distributed by:
Sun Pharmaceutical Industries, Inc.
Cranbury, NJ 08512

NDC 63304-583-60

Fosamprenavir Calcium Tablets

700 mg*

ALERT: Find out about medicines that should NOT be taken with Fosamprenavir Calcium Tablets.

Note to Pharmacist: Do not cover ALERT box with pharmacy label.

Rx only
60 Tablets

ohm®

1119



5204166

For use in combination regimens with or without ritonavir.
*Each coated tablet contains 700 mg of fosamprenavir as fosamprenavir calcium (amorphous).
USUAL DOSAGE: See prescribing information for dosage information.
Store at 20° C to 25° C (68° F to 77° F) (See USP Controlled Room Temperature). Keep bottle tightly closed. Protect from moisture.



N 3

SPACE FOR SERIALIZATION

FOSAMPRENAVIR CALCIUM

fosamprenavir calcium tablet, coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63304-583
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FOSAMPRENAVIR CALCIUM (UNII: ID1GU2627N) (AMPRENAVIR - UNII:5S0W860XNR)	FOSAMPRENAVIR	700 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SHELLAC (UNII: 46N107B710)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	pink	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	RJ47
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63304-583-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/28/2019	
2	NDC:63304-583-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/28/2019	
3	NDC:63304-583-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	11/28/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204024	11/28/2019	

Labeler - Sun Pharmaceutical Industries, Inc. (146974886)

Registrant - Sun Pharmaceutical Industries, Inc. (146974886)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	MANUFACTURE(63304-583)

Revised: 12/2019

Sun Pharmaceutical Industries, Inc.