

CONRX COLD- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet
Eagle Distributors, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ConRx Cold

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325mg

Dextromethorphan HBr 10 mg

Guaifenesin 200mg

Phenylephrine HCl 5mg

Purpose

Pain reliever/ fever reducer

Cough Suppressant

Expectorant

Nasal decongestant

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

- **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: more than 12 tablets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a persistent chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts for more than 7 days
- fever gets more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.
These could be signs of serious conditions.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Do not take more than directed

Adults and Children 12 years and over	<ul style="list-style-type: none">• Take 2 tablets with water every 6-8 hours as needed• Do not take more than 8 tablets in 24 hour
Children under 12 years of age	Consult a doctor

Other Information

- Do not use if pouch is torn or damaged.
- Store between 20-25°C (68-77°F). Avoid excessive heat and humidity.
- See side panel for lot number and expiration date.

Inactive ingredients

Colloidal silicon dioxide, Sodium Lauryl Sulphate, Edetaet disodium, dioctyl sodium sulphosuccinate, polyvinyl pyrrollidone, gelatin, corn starch', magnesium Stearate, microcrystalline cellulose, polyethylene glycol 6000,Sodium Benzoate ,Sodium Starch Glycolate, Stearic acid, Purified Talc, D & C yellow #10 aluminium lake ,Titanium Dioxide.

Product manufactured for:

Eagle Distributors, Inc.

Los Angeles, CA 90011

PRINCIPAL DISPLAY PANEL - 50 Pouch Carton

NDC: 68737-227-11

**Compare to the Active Ingredients in
Tylenol® Cold
Head Congestion***

ConRx

COLD

SEVERE Sore throat -Cough - Headache

- | | |
|------------------------|------------------|
| ▪ Acetaminophen | Sore throat |
| ▪ Dextromethorphan HBr | Cough |
| ▪ Guaifenesin | Chest Congestion |
| ▪ Phenylephrine HCl | Nasal Congestion |

TO OPEN

PUSH IN TAB AND PULL OUT

**Compare to the Active Ingredients in
Tylenol® Cold
Head Congestion***

50 Pouches of 2 Caplets Each

Compare to the Active Ingredients in

NDC 68737-227-11

Tylenol® Cold Head Congestion*

Con-Ra COLD

SEVERE Sore throat - Cough - Headache

- | | |
|------------------------|------------------|
| ■ Acetaminophen | Sore throat |
| ■ Dextromethorphan HBr | Cough |
| ■ Guaifenesin | Chest Congestion |
| ■ Phenylephrine HCl | Nasal Congestion |

Compare to the Active Ingredients in

Tylenol® Cold Head Congestion*

Con-Ra COLD

SEVERE Sore throat - Cough - Headache

- | | |
|------------------------|------------------|
| ■ Acetaminophen | Sore throat |
| ■ Dextromethorphan HBr | Cough |
| ■ Guaifenesin | Chest Congestion |
| ■ Phenylephrine HCl | Nasal Congestion |

TO OPEN



PUSH IN TAB AND PULL OUT

Compare to the Active Ingredients in

Tylenol® Cold Head Congestion*

50 Pouches of 2 Caplets Each





50 Pouches of 2 Caplets Each

Sore throat
Cough
Chest Congestion
Nasal Congestion

- Acetaminophen
- Dextromethorphan HBr
- Guaifenesin
- Phenylephrine HCl

SEVERE Sore throat - Cough - Headache

Con-Pro COLD

Tylenol® Cold
Head Congestion*

Compare to the Active Ingredients in



Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)

adults and children 12 years and over	■ take 2 tablets with water every 6-8 hours as needed
	■ do not take more than 8 tablets in 24 hours
children under 12 years	■ consult a doctor

Other information

- store between 70°-75°F (21°-24°C)

Compare to the Active Ingredients in

Tylenol® Cold

Head Congestion*

Con-Pro COLD

SEVERE Sore throat - Cough - Headache

- Acetaminophen
- Dextromethorphan HBr
- Guaifenesin
- Phenylephrine HCl

Sore throat
Cough
Chest Congestion
Nasal Congestion

DO NOT USE WITH OTHER
MEDICINES CONTAINING
ACETAMINOPHEN

*This product is not manufactured or distributed by: McNeil Consumer Specialty Pharmaceuticals Division of McNeil PPC, Inc. Eagle Distributors, Inc., does not own the Tylenol® Cold Head Congestion trademark.

- do not use if pouch is torn or open
- see side panel for lot number and expiration date
- avoid excessive heat and humidity

Inactive ingredients: colloidal silicon dioxide, sodium lauryl sulfate, edetate disodium, disodium sodium sulfosuccinate, polyvinyl pyrrolidone, gelatin, corn starch, magnesium stearate, microcrystalline cellulose, polyethylene glycol 6000, sodium benzoate, sodium starch glycolate, stearic acid, purified talc, D&C yellow #10 aluminum lake, titanium dioxide.

Questions or comments? call 1-800-570-8850 (M-F 9 am to 5 pm PST)

Part No: 8884_2012



Product manufactured for:
Eagle Distributors, Inc.
Los Angeles, CA 90011

CONRX COLD

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68737-227
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	325 mg
Dextromethorphan Hydrobromide (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	10 mg
Guaifenesin (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	200 mg
Phenylephrine Hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
BIS(1-METHYLAMYL) SODIUM SULFO SUCCINATE (UNII: 772Y8KZU65)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
GELATIN (UNII: 2G86QN327L)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

TALC (UNII: 7SEV7J4R1U)	
D&C YELLOW No. 10 (UNII: 35SW5USQ3G)	
ALUMINUM OXIDE (UNII: LM26O6933)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	12mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68737-227-11	50 in 1 BOX		
1		2 in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	07/10/2012	

Labeler - Eagle Distributors, Inc. (929837425)

Revised: 7/2012

Eagle Distributors, Inc.