

**MEIJER MAXIMUM STRENGTH SEVERE CONGESTION AND COUGH- dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid**

**MEIJER, INC.**

**Reference Label Set Id: 7f4ccd7c-9b17-428b-8cdf-42237728f9b2**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Meijer Maximum Strength Severe Congestion and Cough**

**Drug Facts**

<b>Active ingredients (in each 20 mL)</b>	<b>Purposes</b>
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

## When using this product

- **do not use more than directed**

## Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- **adults and children 12 years of age and older:** 20 mL in dosing cup provided every 4 hours.
- **children under 12 years of age:** Do not use.

## Other information

- each 20 mL contains: **sodium 8 mg**
- low sodium
- store at room temperature
- do not refrigerate

## Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavors, potassium citrate, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

## Questions or comments?

**1-866-467-2748**

## PRINCIPAL DISPLAY PANEL

NDC 41250-446-06

**Compare to Mucinex® Fast- Max® Maximum Strength Severe Congestion & Cough Active Ingredients\***

**Maximum Strength**

**Severe Congestion & Cough**  
 Dextromethorphan HBr - Cough Suppressant  
 Guaifenesin - Expectorant  
 Phenylephrine HCl - Nasal Decongestant



- **Control Cough**
- **Relieves Nasal and Chest Congestion**
- **Thins & loosens mucus**

For Ages 12+

6 FL OZ (180 mL)

**Tamper evident: do not use if printed seal under cap is broken or missing.**

\*This product is not manufactured or distributed by Reckitt Benckiser, the distributor of Mucinex® Fast-Max® Maximum Strength Severe Congestion & Cough.



**MEIJER MAXIMUM STRENGTH SEVERE CONGESTION AND COUGH**

dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-446
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>dextromethorphan hydrobromide</b> (UNII: 9D2RTI9KYH) (dextromethorphan - UNII:7355X3ROTS)	dextromethorphan hydrobromide	20 mg in 20 mL
<b>guaifenesin</b> (UNII: 495W7451VQ) (guaifenesin - UNII:495W7451VQ)	guaifenesin	400 mg in 20 mL
<b>phenylephrine hydrochloride</b> (UNII: 04JA59TNSJ) (phenylephrine - UNII:1WS297W6MV)	phenylephrine hydrochloride	10 mg in 20 mL

## Inactive Ingredients

Ingredient Name	Strength
anhydrous citric acid (UNII: XF417D3PSL)	
edetate disodium (UNII: 7FLD91C86K)	
FD&C Blue NO. 1 (UNII: H3R47K3TBD)	
FD&C Red NO. 40 (UNII: WZB9127XOA)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
propylene glycol (UNII: 6DC9Q167V3)	
propyl gallate (UNII: 8D4SNN7V92)	
water (UNII: 059QF0KO0R)	
sodium benzoate (UNII: OJ245FE5EU)	
sorbitol (UNII: 506T60A25R)	
sucralose (UNII: 96K6UQ3ZD4)	
xanthan gum (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-446-06	180 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/20/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	04/20/2020	

**Labeler** - MEIJER, INC. (006959555)

Revised: 5/2020

MEIJER, INC.