

**IBUPROFEN PM- ibuprofen, diphenhydramine hcl capsule, liquid filled**  
**TARGET Corporation**

-----  
**Drug Facts**

**Active ingredient (in each capsule)**

Diphenhydramine hydrochloride 25 mg

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\***  
**(present as the free acid and potassium salt)**

**\*non-steroidal anti-inflammatory drug**

**Purpose**

Nighttime sleep-aid

**Pain Reliever**

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke

- chest pain
- trouble breathing
- leg swelling
- slurred speech
- weakness in one part or side of body
- pain gets worse or last more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed**
- adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours

### **Other information**

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F)
- swallow whole; do not crush, chew, or dissolve
- protect from light

### **Inactive ingredients**

D&C red #33, FD&C blue #1, gelatin, medium chain triglycerides, polyethylene glycol 600, potassium hydroxide, purified water, sorbitan, sorbitol

### **Questions or comments?**

Call **1-800-910-6874**

### **Principal Display Panel**

Compare to active ingredients in Advil® PM\*\*\*

Ibuprofen PM capsules

Ibuprofen 200 mg

pain reliever (NSAID)

diphenhydramine HCl, 25 mg

nighttime sleep-Aid

**SOFTGELS\*\***

(\*\*LIQUID-FILLED CAPSULES)

\*\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® PM.

**TAMPER-EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Distributed by Target Corporation

Minneapolis, MN 55403

TM & ©2018 Target Brands, Inc.

**Product Packaging**

Product of:  
Lot / Exp.:  
FD005033  
PLD-AS31A



Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2018 Target Brands, Inc.  
094 01 0431 R01  
C-000703-01-039

**Drug Facts**

**Active ingredients (in each capsule)**

Diphenhydramine hydrochloride 25 mg..... Nighttime sleep-aid  
Solubilized ibuprofen (equal to 200 mg ibuprofen NSAID)..... Pain reliever  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Uses**

■ for relief of occasional sleeplessness when associated with minor aches and pains  
■ helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing)  
■ shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinner (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ unless you have time for a full night's sleep  
■ in children under 12 years of age  
■ right before or after heart surgery

**Drug Facts (continued)**

■ with any other product containing diphenhydramine, even one used on skin  
■ if you have sleeplessness without pain

**Ask a doctor before use if**

■ stomach bleeding warning applies along with taking pain relievers or fever reducers  
■ you have problems or serious side effects not mentioned here  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic  
■ you have a breathing problem such as emphysema or chronic bronchitis

**When using this product**

■ drowsiness will occur  
■ avoid alcoholic drinks  
■ do not drive a motor vehicle or operate machinery  
■ take with food or milk, if stomach upset occurs  
■ taking any other antihistamines  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

**Stop use and ask a doctor if**

■ you experience any of the following signs of stomach bleeding  
■ have bloody or black stools  
■ feel faint  
■ vomit blood  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke  
■ chest pain ■ trouble breathing  
■ leg swelling ■ slurred speech  
■ weakness in one part or side of body  
■ pain gets worse or lasts more than 10 days

**Drug Facts (continued)**

■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

■ do not take more than directed  
■ adults and children 12 years and over: take 2 capsules at bedtime  
■ do not take more than 2 capsules in 24 hours

**Other information**

■ each capsule contains: potassium 20 mg  
■ read all warnings and directions before use. Keep carton.  
■ store at 20° to 25°C (68° to 77°F)  
■ avoid excessive heat above 40°C (104°F)  
■ protect from light  
■ swallow whole; do not crush, chew, or dissolve

**Inactive ingredients**

androsorb, D&C red #33, FD&C blue #1, gelatin, hydrolyzed gelatin, medium chain triglycerides, opacode white ink, polyethylene glycol 600, potassium hydroxide, purified water

**Questions or comments?**

Call 1-800-910-8874  
\*\*\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® PM.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Compare to active ingredients in Advil® PM\*\*\*

**ibuprofen PM capsules**



ibuprofen 200 mg  
pain reliever (NSAID)  
diphenhydramine HCl 25 mg  
nighttime sleep-aid



ACTUAL SIZE

**40**

SOFTGELS

**40 SOFTGELS\*\*** (\*\*LIQUID-FILLED CAPSULES)

NDC 11673-756-40

## TARGET Ibuprofen PM Capsules

### IBUPROFEN PM

ibuprofen, diphenhydramine hcl capsule, liquid filled

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-756
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

#### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	1007
<b>Contains</b>			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-756-40	1 in 1 BOX	03/31/2018	03/31/2025
1		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

#### Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
ANDA	ANDA200888	03/31/2018	03/31/2025

**Labeler** - TARGET Corporation (006961700)

Revised: 11/2022

TARGET Corporation