# NICOTINE- nicotine polacrilex gum, chewing Dr. Reddy's Laboratories Inc.

-----

#### Nicotine Polacrilex Gum Coated Ice Mint 4 mg ANDA204833 - Equate

## **Drug Facts**

#### Active ingredient (in each chewing piece)

Nicotine Polacrilex (equal to 4 mg nicotine)

#### **Purpose**

Stop smoking aid

#### Use

 reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

#### Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

# Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

# Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

# Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

## Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### **Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1	Weeks 7	Weeks 10
to 6	to 9	to 12
1 piece		1 piece
every	every	every
1 to 2	2 to 4	4 to 8
hours	hours	hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

#### Other information

• each piece contains: calcium 115 mg, sodium 8 mg

- store at 20 25°C (68 77°F)
- protect from light

## Inactive ingredients

acacia, acesulfame potassium, carnauba wax, D&C yellow #10, FD&C blue #2, FD&C red #40, FD&C yellow #6, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol.

#### Questions or comments?

Call toll-free 1-800-585-8682 weekdays (8:00am - 8:00 pm EST

## **Principal Display Panel**

\*Compare to the active ingredient in Nicorette® Gum

Habitrol

Nicotine Polacrilex Gum

USP, 4 mg (nicotine)

#### STOP SMOKING AID

Gum

4mg

# FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP

If you smoke your first cigarette <u>MORE THAN</u> 30 MINUTES after waking up, use Habitrol gum, 2 mg Gum

Sugar Free coated nicotine gum

COATED FRUIT FLAVOR

# TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivate to quit
- 2. Use Enough Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks
- 3. Use long enough Use Nicotine Polacrilex Gum for the full 12 weeks
- 4. Use with a Support Program as directed in the enclosed User's Guide

To remove the gum, tear off single unit

Peel off backing starting at corner with loose edge

Push gum through foil

- not for sale to those under 18 years of age
- proof of age required

• not for sale in vending machines or from any source where proof of age cannot be verified

\*This product is not manufactured or distributed by Haleon group of companies, distributor of Nicorette® Gum

# BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

#### **DISTRIBUTED BY:**

Dr. Reddy's Laboratories, Inc.

Princeton, NJ 08540

## **Package Label**

#### TO INCREASE YOUR SUCCESS IN QUITTING:

- You must be motivated to quit.
  Use Enough Chew at least 9 pieces of Nicotine Polacriex Gum per day during the first six weeks.
  Use Long Enough Use Nicotine Polacriex Gum for the full 12 weeks.
  Use With a Support Program as directed in the enclosed User's Guide.





NDC 75907-300-15 \*Compare to the active incredient in Nicoratte® Gum

# Habitrol

**Nicotine Polacrilex Gum** USP, 4 mg (nicotine)

STOP SMOKING AID

Gum



#### Sugar-free

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Habitrol 2 mg Gum.

150

4 mg EACH



Fruit Flavor

# Habitrol

# **Nicotine Polacrilex Gum** USP, 4 mg (nicotine)

STOP SMOKING AID

Mithor innformation e such place contains calcium 15 mg, souther 8 mg ff 77 - 55 g 25 - 55 is east ff 77 - 50 g 25 - 55 is east protect them by 2

If you expedience along or frequent coembay, you may use a second please within the hour harmonic or of community, use one gives after enther within the hour harmonic or of community, use one gives after enther m to not one more than it's pleases at day.
 If it is reported to complete brackers. If you had you need to use the garn if it is reported to complete brackers. If you had you need to use the garn if it is reported to complete brackers.

t mg EACH

OST

If you smoke your first cigarette MORE THAN 30 MINUTES after much per Stortided east on politice

FOR THOSE WHO SMOKE THEIR FIRST CHORRETTE WITHIN 30 MINUTES OF WAKING UP.

Sugar-free



wng

IN, recent heart stack, or irregular heartbeat. Machine can increase STOP SMOKING AID

PLD-AB21C FC009487

Princeton, NJ 08540 SULED BY:

BLISTER PACKAGED FOR YOUR PROTECTION. Do not use if individual seals are open or torm.

not for ealst for those under 18 years of age
 proof of age required
 not for eals in vooding machines or from any source
 most for eals in vooding machines

		599
bjecoe be, qoë pv. pje pust p	e iseal in eau galifing to	publices April chances rewing a place
the notice gam, or while	Research autom somethic	
	anny a styled art to bloom	
	pathe Buswarp uplied toods	
	d it stegles. Then park it b	
hee simen year mounts	ditides saff select it is	
		spre
hand ent) may at yew mach	no a banu od žeum bra en	colon e a medico solos solos
sworl 8 at 4 tand ext lag at yew night	2 to 4 booms of 2 to 4 cond a con	enod 5 of 1 cabon s ai mug cafaca afluer
hand ent) may at yew mach	no a banu od žeum bra en	colon e a medico solos solos
sworl 8 at 4 tand ext lag at yew night	2 to 4 booms of 2 to 4 cond a con	subside a medico subside a medico

we make the interesting and the property constraint converge converge, and the property constraint converge con





#### **HABITROL** Nicotine Polacrilex Gum Coated Fruit Flavor

#### **NICOTINE**

nicotine polacrilex gum, chewing

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:75907-300

**Route of Administration** ORAL

## **Active Ingredient/Active Moiety**

Ingredient Name Basis of Strength Strength

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) NICOTINE 4 mg

# **Inactive Ingredients**

Ingredient Name	Strength
<b>ACACIA</b> (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

#### **Product Characteristics**

Color	yellow (light tan)	Score	no score
Shape	SQUARE	Size	14mm
Flavor	FRUIT	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75907-300- 80	80 in 1 CARTON	06/12/2025	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:75907-300- 15	150 in 1 CARTON	06/12/2025	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079038	06/12/2025	

# Labeler - Dr. Reddy's Laboratories Inc. (802315887)

Revised: 6/2025 Dr. Reddy's Laboratories Inc.