

**BACTOSHIELD CHG- chlorhexidine gluconate liquid**  
**SC Johnson Professional USA, Inc.**

-----  
**Bactoshield® CHG 4%**

***Drug Facts***

**Active ingredient**

Chlorhexidine gluconate solution, 4%

**Purposes**

Surgical hand scrub

Healthcare personnel handwash

Skin wound and general skin cleanser

**Uses**

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** handwash to help reduce bacteria that potentially can cause disease
- **skin wound and general skin cleanser:** helps reduce bacteria on the skin

**Warnings**

**For external use only**

**Allergy alert**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- if you are allergic to chlorhexidine gluconate or any other ingredient in this preparation
- as a patient preoperative skin preparation (especially on the head or face)
- in contact with meninges
- in the genital area

## **When using this product**

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.
- if contact occurs, rinse with cold water right away
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when an underlying condition makes it necessary to reduce the bacterial population of the skin

**Stop use and ask a doctor if** irritation, sensitization or allergic reaction occurs and lasts for 72 hours. These may be signs of a serious condition.

## **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use full strength
- do not dilute

## **Surgical hand scrub**

- remove jewelry
- wet hands and forearms with water and apply 5 mL of the product
- wash/scrub hands and forearms for 3 minutes paying particular attention to the nails, cuticles, and interdigital spaces
- rinse thoroughly with water
- wash for an additional 3 minutes with 5 mL of the product and rinse under running water
- dry thoroughly

## **Healthcare personnel handwash**

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds
- rinse and dry thoroughly

## **Skin wound and general skin cleanser**

- thoroughly rinse the area to be cleansed with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently. Rinse thoroughly with water.
- rinse again thoroughly

## **Other information**

- store between 20-25°C (68-77°F)

- avoid excessive heat (above 104°F/40°C)
- for additional information, see Safety Data Sheet (SDS)
- if swallowed get medical help or contact a Poison Control Center immediately

### **Inactive ingredients**

cocamide DEA, cocamine oxide, fragrance, gluconic acid<sup>1</sup>, gluconolactone<sup>1</sup>, hydroxyethyl cellulose, isopropyl alcohol 4%, PEG-75 lanolin, PEG-150 distearate, propylene glycol, quaternium-60, water

---

1 contains one or more of these ingredients

### **Questions or comments?**

Call 1-866-783-0422 Monday through Friday 8:00 AM to 5:00 PM EST

### **PRINCIPAL DISPLAY PANEL - 946 mL Bottle Label**

SCJ PROFESSIONAL™  
HEALTHCARE

Bactoshield®  
CHG 4%

Surgical Scrub  
Chlorhexidine Gluconate 4%  
Solution Antiseptic

Surgical Scrub

4% CHG

Antimicrobial

Net Contents: 946mL (32 fl oz)

REORDER #  
1344-24

1344-21J(321)  
SAP # 4000000058

deb  
SKIN CARE

**SCJ PROFESSIONAL™**  
HEALTHCARE

**Bactoshield®**  
CHG 4%

**Surgical Scrub**  
Chlorhexidine Gluconate 4%  
Solution Antiseptic

 **Surgical Scrub**

 **4% CHG**

 **Antimicrobial**

Net Contents: 946mL (32 fl oz)

 **9124006203**

REBUDICE # 134424  
1344911323  
SAP # 4000000058



**Drug Facts**

**Active ingredient** Chlorhexidine gluconate solution, 4%..... Surgical hand scrub  
Healthcare personnel handwash  
Skin wound and general skin cleanser

**Purposes**

**Uses**

- **surgical hand scrub:** significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel handwash:** handwash to help reduce bacteria that potentially can cause disease
- **skin wound and general skin cleanser:** helps reduce bacteria on the skin

**Warnings**  
For external use only

**Allergy alert**  
This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- facial swelling
- shock
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- if you are allergic to chlorhexidine gluconate or any other ingredient in this preparation
- as a patient preoperative skin preparation (especially on the head or face)
- in contact with meningies
- in the genital area

**Drug Facts (continued)**

**When using this product**

- Keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums.
- if contact occurs, rinse with cold water right away
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when an underlying condition makes it necessary to reduce the bacterial population of the skin

**Stop use and ask a doctor if** irritation, sensitization or allergic reaction occurs and lasts for 72 hours. These may be signs of a serious condition.

**Keep out of reach of children.**  
If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use full strength
- do not dilute

**Surgical hand scrub:**

- remove jewelry
- wet hands and forearms with water and apply 5 mL of the product
- wash/scrub hands and forearms for 3 minutes paying particular attention to the nails, cuticles, and interdigital spaces

**Drug Facts (continued)**

- rinse thoroughly with water
- wash for an additional 3 minutes with 5 mL of the product and rinse under running water
- dry thoroughly

**Healthcare personnel handwash:**

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds
- rinse and dry thoroughly

**Skin wound and general skin cleanser:**

- thoroughly rinse the area to be cleansed with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently. Rinse thoroughly with water.
- rinse again thoroughly

**Other information**

- store between 20-25°C (68-77°F)
- avoid excessive heat (above 104°F/40°C)
- for additional information, see Safety Data Sheet (SDS)
- if swallowed get medical help or contact a Poison Control Center immediately

**Inactive ingredients**  
cocamide DEA, cocamine oxide, fragrance, gluconic acid\*, gluconolactone\*, hydroxyethyl cellulose, isopropyl alcohol 4%, PEG-75 lanolin, PEG-150 distearate, propylene glycol, quaternium-60, water \*contains one or more of these ingredients

**Questions or comments?**  
☎ Call 1-866-783-0422 Monday through Friday 8:00 AM to 5:00 PM EST

Manufactured for: SC Johnson Professional USA, Inc.  
Charlotte, NC 28217 1-866-783-0422 www.scjcp.com

**WARNING:** Cancer - www.P65Warnings.ca.gov.  
Keep from freezing.

## BACTOSHIELD CHG

chlorhexidine gluconate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11084-128
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORHEXIDINE GLUCONATE</b> (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	40 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>COCO DIETHANOLAMIDE</b> (UNII: 92005F972D)	
<b>COCAMINE OXIDE</b> (UNII: QWA2IZI6FI)	
<b>GLUCONIC ACID</b> (UNII: R4R8J0Q44B)	
<b>GLUCONOLACTONE</b> (UNII: WQ29KQ9POT)	
<b>HYDROXYETHYL CELLULOSE (5000 MP.A.S AT 1%)</b> (UNII: X70SE62ZAR)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>PEG-75 LANOLIN</b> (UNII: 09179OX7TB)	
<b>PEG-150 DISTEARATE</b> (UNII: 6F36Q0I0AC)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>Quaternium-33</b> (UNII: XPS4174QZJ)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11084-128-03	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/15/2019	
2	NDC:11084-128-34	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/15/2019	12/31/2023
3	NDC:11084-128-24	946 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/15/2019	
4	NDC:11084-128-08	3785 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/15/2019	12/31/2023

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019258	08/15/2019	

**Labeler** - SC Johnson Professional USA, Inc. (607378015)

## Establishment

Name	Address	ID/FEI	Business Operations
AVA, Inc.		615317075	MANUFACTURE(11084-128)

Revised: 7/2025

SC Johnson Professional USA, Inc.