SEVERE COLD AND FLU DAYTIME NON DROWSY NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride / acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride Walgreens

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DAYTIME NON DROWSY Severe Cold & Flu NIGHTTIME Severe Cold & Flu

SEVERE COLD AND FLU

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains headache
- fever sore throat reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Allergy alert Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease
 diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Drug Facts (continued)

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7

days ■ fever gets worse or lasts more than 3 days

- redness or swelling is present
 new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed see Overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 years & over	2 softgels with water every 4 hours	
children 4 to under 12 years	ask a doctor	
children under 4 years	do not use	

■ when using other Nighttime or Daytime products, carefully read each label to ensure correct dosing

Other information

■ store at room temperature

Inactive ingredients

FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

NIGHTTIME SEVERE COLD & FLU SOFTGELS

Drug Facts

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant

Doxylamine succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep minor aches & pains headache
- fever sore throat runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

NIGHTTIME SEVERE COLD & FLU SOFTGELS

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease diabetes glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

NIGHTTIME SEVERE COLD & FLU SOFTGELS

Drug Facts (continued)

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7

days ■ fever gets worse or lasts more than 3 days

- redness or swelling is present
 new symptoms occur
- cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed see Overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 years & over	2 softgels with water every 4 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

Other information

store at room temperature

Inactive ingredients

D&C Yellow #10, FD&C Blue #1, gelatin, glycerin, polyethylene glycol 400, povidone K30, propylene glycol, purified water, shellac, sorbitol sorbitan, sodium hydroxide, titanium dioxide

Questions or comments?

Call toll free: 1-888-333-9792

PRINCIPAL DISPLAY PANEL - Kit Carton DAY & NIGHT PACK

NDC 0363-9896-24

Walgreens

Compare to Vicks® DayQuil® Severe Cold & Flu & Vicks® NyQuil® Severe Cold & Flu active ingredients††

DAYTIME • NON-DROWSY

Severe
Cold & Flu
ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCI / NASAL DECONGESTANT

DISSOLVES QUICKLY

MAXIMUM STRENGTH

ACTUAL SIZE

16SOFTGELS

NIGHTTIME

Severe
Cold & Flu
ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE / ANTIHISTAMINE
PHENYLEPHRINE HCI / NASAL DECONGESTANT

MAXIMUM STRENGTH

ACTUAL SIZE

8SOFTGELS

24 SOFTGELS

SEVERE COLD AND FLU DAYTIME NON DROWSY NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride / acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0363-9896

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9896-24	1 in 1 PACKAGE, COMBINATION; Type 0: Not a Combination Product	07/05/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	16
Part 2	1 BLISTER PACK	8

Part 1 of 2

SEVERE COLD AND FLU DAYTIME NON-DROWSY

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	

Inactive Ingredients			
Ingredient Name	Strength		
FD&C RED NO. 40 (UNII: WZB9127XOA)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
GELATIN (UNII: 2G86QN327L)			
GLYCERIN (UNII: PDC6A3C0OX)			
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
POVIDONE K30 (UNII: U725QWY32X)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SORBITOL (UNII: 506T60A25R)			
SORBITAN (UNII: 6092ICV9RU)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL (OBLONG)	Size	20mm
Flavor		Imprint Code	341
Contains			

Pa	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category Application Number or Monograph Citation Marketing Start Date Marketing End Date				
OTC monograph final	part341	07/05/2018		

Part 2 of 2

SEVERE COLD AND FLU NIGHTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Ingredient Name Strength D&C YELLOW NO. 10 (UNII: 355W5USQ3G) FD&C BLUE NO. 1 (UNII: H3R47K3TBD) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C0OX) POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) POVIDONE K30 (UNII: U725QWY32X) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) WATER (UNII: 059QF0K0OR) SHELLAC (UNII: 46N107B71O) SODIUM HYDROXIDE (UNII: 55X04QC32I) TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics			
Color	green	Score	no score
Shape	OVAL (OBLONG)	Size	20mm
Flavor		Imprint Code	116
Contains			

Pa	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/05/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/05/2018	

Labeler - Walgreens (008965063)

Registrant - Spirit Pharmaceuticals LLC (179621011)

Establishment			
Name	Address	ID/FEI	Business Operations
MEDGEL PVT LTD		677385498	manufacture(0363-9896)

Revised: 6/2022 Walgreens