

**GLYCERIN - glycerin suppository**  
**Rite Aid Corporation**

-----  
**Rite Aid Pediatric Glycerin Supp**

**Active ingredient (in each suppository)**

Glycerin 1 g

**Purpose**

Hyperosmotic laxative

**Uses**

- for relief of occasional constipation
- this product generally produces a bowel movement in 1/4 to 1 hour

**Warnings**

**For rectal use only**

May cause rectal discomfort of burning sensation

**Ask a Doctor before using laxative if child has**

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits lasting more than two weeks
- already used a laxative for more than 1 week

**Stop use and consult a doctor if child has**

- rectal bleeding
- no bowel movement within 1 hour of using this product

These symptoms may indicate a serious condition.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

**Single daily dosage**

---

children 2 to under 6 years	1 suppository or as directed by a doctor
children under 2 years	ask a doctor

---

Insert suppository well up into rectum.

Suppository need not melt completely to produce laxative action.

### Other information

- Store container tightly closed.
- Keep away from excessive heat.

### Inactive ingredients

purified water, sodium hydroxide, stearic acid

### Pediatric Glycerin Suppositories, 25 count

Compare to the active ingredient in Fleet® Glycerin Suppositories\*

**CHILDREN'S LAXATIVE**  
**GLYCERIN SUPPOSITORIES**

**GLYCERIN 1g**

for fast, gentle relief of occasional constipation

**25**  
CHILD SIZE SUPPOSITORIES

**RITE AID PHARMACY**

**Drug Facts**

**Active ingredient (in each suppository)** Purpose  
Glycerin 1g .....Hyperosmotic laxative

**Uses**  
■ for relief of occasional constipation  
■ this product generally produces bowel movement in 1/4 to 1 hour

**Warnings**  
For rectal use only  
■ May cause rectal discomfort or a burning sensation.  
■ Ask a doctor before using any laxative if child has  
■ abdominal pain, nausea or vomiting  
■ a sudden change in bowel habits lasting more than 2 weeks  
■ already used a laxative for more than 1 week  
■ Stop use and consult a doctor if child has  
■ rectal bleeding  
■ no bowel movement within 1 hour of using this product  
■ These symptoms may indicate a serious condition.  
■ Keep out of reach of children.  
■ If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** Single daily dosage  
children 2 to 6 years 1 suppository, or as directed by a doctor  
children under 2 years ask a doctor

Insert suppository well up into rectum.  
Suppository need not melt completely to produce laxative action.

**Other information**  
■ Store container tightly closed.  
■ Keep away from excessive heat.

**Inactive ingredients**  
purified water, sodium hydroxide, stearic acid

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

11822 33218 7

NO VARNISH AREA

\*This product is not the same as the active ingredient in Fleet® Glycerin Suppositories. Rite Aid, the Rite Aid logo, and Rite Aid Pharmacy are registered trademarks of Rite Aid Inc. © 2011 Rite Aid Inc.

The product package shown above represents a sample of that currently in use. Additional packaging may also be available.

RAPedGlycerinSupp25ct.jpg

Distributed by

Rite Aid

30 Hunter Lane

Camp Hill, PA 17011

## GLYCERIN

glycerin suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-3321
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C00X) (GLYCERIN - UNII:PDC6A3C00X)	GLYCERIN	1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3321-8	25 in 1 JAR; Type 0: Not a Combination Product	02/16/2012	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	02/16/2012	

**Labeler** - Rite Aid Corporation (014578892)

**Registrant** - DSC Laboratories Inc. (097807374)

### Establishment

Name	Address	ID/FEI	Business Operations
DSC Laboratories, Inc.		097807374	manufacture(11822-3321)

Revised: 12/2023

Rite Aid Corporation