

**AMITRIPTYLINE HYDROCHLORIDE- amitriptyline hydrochloride tablet
REMEDYREPACK INC.**

**Amitriptyline Hydrochloride Tablets, USP
Rx only**


Suicidality and Antidepressant Drugs:

Antidepressants increased the risk compared to placebo of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults in short-term studies of major depressive disorder (MDD) and other psychiatric disorders. Anyone considering the use of amitriptyline hydrochloride tablets or any other antidepressant in a child, adolescent, or young adult must balance this risk with the clinical need. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction in risk with antidepressants compared to placebo in adults aged 65 and older. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior. Families and caregivers should be advised of the need for close observation and communication with the prescriber. Amitriptyline hydrochloride is not approved for use in pediatric patients. (See WARNINGS: Clinical Worsening and Suicide Risk, PRECAUTIONS: Information for Patients, and PRECAUTIONS: Pediatric Use)

DESCRIPTION

Amitriptyline Hydrochloride, USP a dibenzocycloheptadiene derivative, is a white or practically white, odorless or partially odorless, crystalline powder or small crystals which is Freely soluble in water, in alcohol, in chloroform and in methanol, insoluble in ether.

It is designated chemically as 10,11-Dihydro- *N,N*-dimethyl-5 *H*-dibenzo[*a,d*]cycloheptene- Δ^5 - γ -propylamine hydrochloride. It has the following structural formula:



$C_{20}H_{23}N \cdot HCl$

M.W. 313.86

Each tablet for oral administration contains 10, 25, 50, 75, 100, or 150 mg amitriptyline hydrochloride USP. Inactive ingredients include colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl cellulose, isopropyl alcohol, lactose monohydrate, magnesium stearate, microcrystalline cellulose, pregelatinized starch. The film coating contains Opadry[®] Pink (carmine, FD&C Blue #1, FD&C Yellow #6, Hypromellose 2910, polyethylene glycol, titanium dioxide); Opadry[®] Green (FD&C Blue #1, FD&C Blue #2, FD&C Red #40, Hypromellose 2910, iron oxide yellow, polyethylene glycol, titanium dioxide); Opadry[®] Brown (FD&C Blue #2, FD&C Red #40, Hypromellose 2910, iron oxide red, polyethylene glycol, titanium dioxide); Opadry[®] Purple (carmine, FD&C Blue #2, ferrousferrous oxide, Hypromellose 2910, polyethylene glycol, titanium dioxide); Opadry[®] Orange (FD&C Yellow #6, Hypromellose 2910, iron oxide red, polyethylene glycol, titanium dioxide); Opadry[®] Green (FD&C Blue #1, FD&C Blue #2, FD&C Red #40, Hypromellose 2910, iron oxide yellow, polyethylene glycol, titanium dioxide).

FDA approved dissolution specifications differs from the USP.

CLINICAL PHARMACOLOGY

Amitriptyline hydrochloride is an antidepressant with sedative effects. Its mechanism of action in man is not known. It is not a monoamine oxidase inhibitor and it does not act primarily by stimulation of the central nervous system.

Amitriptyline hydrochloride inhibits the membrane pump mechanism responsible for uptake of norepinephrine and serotonin in adrenergic and serotonergic neurons. Pharmacologically, this action may potentiate or prolong neuronal activity since reuptake of these biogenic amines is important physiologically in terminating transmitting activity. This interference with reuptake of norepinephrine and/or serotonin is believed by some to underlie the antidepressant activity of amitriptyline hydrochloride.

INDICATIONS AND USAGE

For the relief of symptoms of depression. Endogenous depression is more likely to be alleviated than are other depressive states.

CONTRAINDICATIONS

Amitriptyline hydrochloride is contraindicated in patients who have shown prior

hypersensitivity to it.

It should not be given concomitantly with monoamine oxidase inhibitors. Hyperpyretic crises, severe convulsions, and deaths have occurred in patients receiving tricyclic antidepressant and monoamine oxidase inhibiting drugs simultaneously. When it is desired to replace a monoamine oxidase inhibitor with amitriptyline hydrochloride, a minimum of 14 days should be allowed to elapse after the former is discontinued. Amitriptyline hydrochloride should then be initiated cautiously with gradual increase in dosage until optimum response is achieved.

Amitriptyline hydrochloride should not be given with cisapride due to the potential for increased QT interval and increased risk for arrhythmia.

This drug is not recommended for use during the acute recovery phase following myocardial infarction.

WARNINGS

Clinical Worsening and Suicide Risk

Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a long-standing concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment. Pooled analyses of short-term placebo-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18 to 24) with major depressive disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction with antidepressants compared to placebo in adults aged 65 and older.

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, obsessive compulsive disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of 9 antidepressant drugs in over 4,400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients. There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger patients for almost all drugs studied. There were differences in absolute risk of suicidality across the different indications, with the highest incidence in MDD. The risk differences (drug vs placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number of cases of suicidality per 1,000 patients treated) are provided in Table 1.

Table 1

	Drug-Placebo Difference in
--	-----------------------------------

Age Range	Number of Cases of Suicidality per 1,000 Patients Treated
	Increases Compared to Placebo
<18	14 additional cases
18 to 24	5 additional cases
	Decreases Compared to Placebo
25 to 64	1 fewer case
≥65	6 fewer cases

No suicides occurred in any of the pediatric trials. There were suicides in the adult trials, but the number was not sufficient to reach any conclusion about drug effect on suicide.

It is unknown whether the suicidality risk extends to longerterm use, i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression.

All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases.

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for major depressive disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

Families and caregivers of patients being treated with antidepressants for major depressive disorder or other indications, both psychiatric and nonpsychiatric, should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to health care providers. Such monitoring should include daily observation by families and caregivers.

Prescriptions for amitriptyline hydrochloride tablets should be written for the smallest quantity of tablets consistent with good patient management, in order to reduce the risk of overdose.

Serotonin Syndrome:

The development of a potentially life-threatening serotonin syndrome has been reported

with tricyclic antidepressants, including amitriptyline hydrochloride tablets, alone but particularly with concomitant use of other serotonergic drugs (including triptans, tricyclic antidepressants, SSRI/SNRI, fentanyl, lithium, tramadol, tryptophan, buspirone, and St. John's Wort) and with drugs that impair metabolism of serotonin (in particular, MAOIs, both those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue). Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular changes (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Patients should be monitored for the emergence of serotonin syndrome.

The concomitant use of amitriptyline hydrochloride tablets with MAOIs intended to treat psychiatric disorders is contraindicated. Amitriptyline hydrochloride tablets should also not be started in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue. All reports with methylene blue that provided information on the route of administration involved intravenous administration in the dose range of 1 mg/kg to 8 mg/kg. No reports involved the administration of methylene blue by other routes (such as oral tablets or local tissue injection) or at lower doses. There may be circumstances when it is necessary to initiate treatment with an MAOI such as linezolid or intravenous methylene blue in a patient taking amitriptyline hydrochloride tablets. Amitriptyline hydrochloride tablets should be discontinued before initiating treatment with the MAOI (see CONTRAINDICATIONS and DOSAGE AND ADMINISTRATION). If concomitant use of amitriptyline hydrochloride tablets with other serotonergic drugs, including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, buspirone, tryptophan, and St. John's Wort is clinically warranted, patients should be made aware of a potential increased risk for serotonin syndrome, particularly during treatment initiation and dose increases. Treatment with amitriptyline hydrochloride tablets and any concomitant serotonergic agents should be discontinued immediately if the above events occur and supportive symptomatic treatment should be initiated.

Screening Patients for Bipolar Disorder

A major depressive episode may be the initial presentation of bipolar disorder. It is generally believed (though not established in controlled trials) that treating such an episode with an antidepressant alone may increase the likelihood of precipitation of a mixed/manic episode in patients at risk for bipolar disorder. Whether any of the symptoms described above represent such a conversion is unknown. However, prior to initiating treatment with an antidepressant, patients with depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression. It should be noted that amitriptyline hydrochloride tablets are not approved for use in treating bipolar depression.

Amitriptyline hydrochloride tablets may block the antihypertensive action of guanethidine or similarly acting compounds.

It should be used with caution in patients with a history of seizures and, because of its atropine-like action, in patients with a history of urinary retention or angle-closure glaucoma. In patients with angle-closure glaucoma, even average doses may precipitate an attack.

Patients with cardiovascular disorders should be watched closely. Tricyclic

antidepressant drugs, including amitriptyline hydrochloride tablets, particularly when given in high doses, have been reported to produce arrhythmias, sinus tachycardia, and prolongation of the conduction time. Myocardial infarction and stroke have been reported with drugs of this class.

Close supervision is required when amitriptyline hydrochloride tablets are given to hyperthyroid patients or those receiving thyroid medication.

Amitriptyline hydrochloride tablets may enhance the response to alcohol and the effects of barbiturates and other CNS depressants. In patients who may use alcohol excessively, it should be borne in mind that the potentiation may increase the danger inherent in any suicide attempt or overdose. Delirium has been reported with concurrent administration of amitriptyline hydrochloride tablets and disulfiram.

Hyponatremia

Hyponatremia has occurred as a result of treatment with amitriptyline hydrochloride tablets. In many cases, hyponatremia appears to be the result of the syndrome of inappropriate antidiuretic hormone secretion (SIADH). Signs and symptoms of hyponatremia include headache, difficulty concentrating, memory impairment, confusion, weakness, and unsteadiness, which may lead to falls. Signs and symptoms associated with more severe and/or acute cases have included syncope, seizure, coma, respiratory arrest, and death.

In patients with symptomatic hyponatremia, discontinue amitriptyline hydrochloride tablets and institute appropriate medical intervention. Elderly patients, patients taking diuretics, and those who are volume-depleted may be at greater risk of developing hyponatremia with amitriptyline hydrochloride tablets.

Angle-Closure Glaucoma

The pupillary dilation that occurs following use of many antidepressant drugs including amitriptyline hydrochloride tablets may trigger an angle closure attack in a patient with anatomically narrow angles who does not have a patent iridectomy.

Usage in Pregnancy

Teratogenic effects were not observed in mice, rats, or rabbits when amitriptyline was given orally at doses of 2 to 40 mg/kg/day (up to 13 times the maximum recommended human dose¹). Studies in literature have shown amitriptyline to be teratogenic in mice and hamsters when given by various routes of administration at doses of 28 to 100 mg/kg/day (9 to 33 times the maximum recommended human dose), producing multiple malformations. Another study in the rat reported that an oral dose of 25 mg/kg/day (8 times the maximum recommended human dose) produced delays in ossification of fetal vertebral bodies without other signs of embryotoxicity. In rabbits, an oral dose of 60 mg/kg/day (20 times the maximum recommended human dose) was reported to cause incomplete ossification of cranial bones.

Amitriptyline has been shown to cross the placenta. Although a causal relationship has not been established, there have been a few reports of adverse events, including CNS effects, limb deformities, or developmental delay, in infants whose mothers had taken amitriptyline during pregnancy.

There are no adequate and well-controlled studies in pregnant women. Amitriptyline hydrochloride should be used during pregnancy only if the potential benefit to the

mother justifies the potential risk to the fetus.

¹Based on a maximum recommended amitriptyline dose of 150 mg/day or 3 mg/kg/day for a 50 kg patient.

Nursing Mothers

Amitriptyline hydrochloride is excreted into breast milk. In one report in which a patient received amitriptyline hydrochloride tablet 100 mg/day while nursing her infant, levels of 83 to 141 ng/mL were detected in the mother's serum. Levels of 135 to 151 ng/mL were found in the breast milk, but no trace of the drug could be detected in the infant's serum.

Because of the potential for serious adverse reactions in nursing infants from amitriptyline hydrochloride tablets, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Usage in Pediatric Patients

In view of the lack of experience with the use of this drug in pediatric patients, it is not recommended at the present time for patients under 12 years of age.

PRECAUTIONS

Schizophrenic patients may develop increased symptoms of psychosis; patients with paranoid symptomatology may have an exaggeration of such symptoms. Depressed patients, particularly those with known manic-depressive illness, may experience a shift to mania or hypomania. In these circumstances the dose of amitriptyline hydrochloride may be reduced or a major tranquilizer such as perphenazine may be administered concurrently.

The possibility of suicide in depressed patients remains until significant remission occurs. Potentially suicidal patients should not have access to large quantities of this drug. Prescriptions should be written for the smallest amount feasible.

Concurrent administration of amitriptyline hydrochloride and electroshock therapy may increase the hazards associated with such therapy. Such treatment should be limited to patients for whom it is essential.

When possible, the drug should be discontinued several days before elective surgery.

Both elevation and lowering of blood sugar levels have been reported.

Amitriptyline hydrochloride should be used with caution in patients with impaired liver function.

Information for Patients

Prescribers or other health professionals should inform patients, their families, and their caregivers about the benefits and risks associated with treatment with amitriptyline hydrochloride tablets and should counsel them in its appropriate use. A patient Medication Guide about "Antidepressant Medicines, Depression and other Serious Mental Illnesses, and Suicidal Thoughts or Actions" is available for amitriptyline hydrochloride tablets. The prescriber or health professional should instruct patients, their families, and their caregivers to read the Medication Guide and should assist them in understanding

its contents. Patients should be given the opportunity to discuss the contents of the Medication Guide and to obtain answers to any questions they may have. The complete text of the Medication Guide is reprinted at the end of this document.

Patients should be advised of the following issues and asked to alert their prescriber if these occur while taking amitriptyline hydrochloride tablets.

Patients should be advised that taking amitriptyline hydrochloride tablets can cause mild pupillary dilation, which in susceptible individuals, can lead to an episode of angle-closure glaucoma. Pre-existing glaucoma is almost always open-angle glaucoma because angle-closure glaucoma, when diagnosed, can be treated definitively with iridectomy. Open-angle glaucoma is not a risk factor for angle-closure glaucoma. Patients may wish to be examined to determine whether they are susceptible to angle closure, and have a prophylactic procedure (e.g., iridectomy), if they are susceptible.

Clinical Worsening and Suicide Risk

Patients, their families, and their caregivers should be encouraged to be alert to the emergence of anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, mania, other unusual changes in behavior, worsening of depression, and suicidal ideation, especially early during antidepressant treatment and when the dose is adjusted up or down. Families and caregivers of patients should be advised to look for the emergence of such symptoms on a day-to-day basis, since changes may be abrupt. Such symptoms should be reported to the patient's prescriber or health professional, especially if they are severe, abrupt in onset, or were not part of the patient's presenting symptoms. Symptoms such as these may be associated with an increased risk for suicidal thinking and behavior and indicate a need for very close monitoring and possibly changes in the medication.

While on therapy with amitriptyline hydrochloride, patients should be advised as to the possible impairment of mental and/or physical abilities required for performance of hazardous tasks, such as operating machinery or driving a motor vehicle.

DRUG INTERACTIONS

Topiramate

Some patients may experience a large increase in amitriptyline hydrochloride concentration in the presence of topiramate and any adjustments in amitriptyline hydrochloride dose should be made according to the patient's clinical response and not on the basis of plasma levels.

Drugs Metabolized by P450 2D6

The biochemical activity of the drug metabolizing isozyme cytochrome P450 2D6 (debrisoquin hydroxylase) is reduced in a subset of the caucasian population (about 7 to 10% of Caucasians are so called "poor metabolizers"); reliable estimates of the prevalence of reduced P450 2D6 isozyme activity among Asian, African and other populations are not yet available. Poor metabolizers have higher than expected plasma concentrations of tricyclic antidepressants (TCAs) when given usual doses. Depending on the fraction of drug metabolized by P450 2D6, the increase in plasma concentration may be small, or quite large (8 fold increase in plasma AUC of the TCA).

In addition, certain drugs inhibit the activity of this isozyme and make normal metabolizers resemble poor metabolizers. An individual who is stable on a given dose of TCA may become abruptly toxic when given one of these inhibiting drugs as concomitant therapy. The drugs that inhibit cytochrome P450 2D6 include some that are not metabolized by the enzyme (quinidine; cimetidine) and many that are substrates for P450 2D6 (many other antidepressants, phenothiazines, and the Type 1C antiarrhythmics propafenone and flecainide). While all the selective serotonin reuptake inhibitors (SSRIs), e.g., fluoxetine, sertraline, and paroxetine, inhibit P450 2D6, they may vary in the extent of inhibition. The extent to which SSRI-TCA interactions may pose clinical problems will depend on the degree of inhibition and the pharmacokinetics of the SSRI involved. Nevertheless, caution is indicated in the coadministration of TCAs with any of the SSRIs and also in switching from one class to the other. Of particular importance, sufficient time must elapse before initiating TCA treatment in a patient being withdrawn from fluoxetine, given the long half-life of the parent and active metabolite (at least 5 weeks may be necessary).

Concomitant use of tricyclic antidepressants with drugs that can inhibit cytochrome P450 2D6 may require lower doses than usually prescribed for either the tricyclic antidepressant or the other drug. Furthermore, whenever one of these other drugs is withdrawn from co-therapy, an increased dose of tricyclic antidepressant may be required. It is desirable to monitor TCA plasma levels whenever a TCA is going to be coadministered with another drug known to be an inhibitor of P450 2D6.

Monoamine oxidase inhibitors – see **CONTRAINDICATIONS** section. Guanethidine or similarly acting compounds; thyroid medication; alcohol, barbiturates and other CNS depressants; and disulfiram – see **WARNINGS** section.

When amitriptyline hydrochloride is given with anticholinergic agents or sympathomimetic drugs, including epinephrine combined with local anesthetics, close supervision and careful adjustment of dosages are required.

Hyperpyrexia has been reported when amitriptyline hydrochloride is administered with anticholinergic agents or with neuroleptic drugs, particularly during hot weather.

Paralytic ileus may occur in patients taking tricyclic antidepressants in combination with anticholinergic-type drugs.

Cimetidine is reported to reduce hepatic metabolism of certain tricyclic antidepressants, thereby delaying elimination and increasing steady-state concentrations of these drugs. Clinically significant effects have been reported with the tricyclic antidepressants when used concomitantly with cimetidine. Increases in plasma levels of tricyclic antidepressants, and in the frequency and severity of side effects, particularly anticholinergic, have been reported when cimetidine was added to the drug regimen. Discontinuation of cimetidine in well-controlled patients receiving tricyclic antidepressants and cimetidine may decrease the plasma levels and efficacy of the antidepressants.

Caution is advised if patients receive large doses of ethchlorvynol concurrently. Transient delirium has been reported in patients who were treated with one gram of ethchlorvynol and 75 to 150 mg of amitriptyline hydrochloride.

Geriatric Use

Clinical experience has not identified differences in responses between elderly and

younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic function, concomitant disease and other drug therapy in elderly patients.

Geriatric patients are particularly sensitive to the anticholinergic side effects of tricyclic antidepressants including amitriptyline hydrochloride. Peripheral anticholinergic effects include tachycardia, urinary retention, constipation, dry mouth, blurred vision, and exacerbation of narrow-angle glaucoma. Central nervous system anticholinergic effects include cognitive impairment, psychomotor slowing, confusion, sedation, and delirium. Elderly patients taking amitriptyline hydrochloride may be at increased risk for falls. Elderly patients should be started on low doses of amitriptyline hydrochloride and observed closely (see **DOSAGE AND ADMINISTRATION**).

Pediatric Use

Safety and effectiveness in the pediatric population have not been established (see **BOX WARNING** and **WARNINGS: Clinical Worsening and Suicide Risk**). Anyone considering the use of amitriptyline hydrochloride in a child or adolescent must balance the potential risks with the clinical need.

ADVERSE REACTIONS

Within each category the following adverse reactions are listed in order of decreasing severity. Included in the listing are a few adverse reactions which have not been reported with this specific drug. However, pharmacological similarities among the tricyclic antidepressant drugs require that each of the reactions be considered when amitriptyline hydrochloride is administered.

Cardiovascular: Myocardial infarction; stroke; nonspecific ECG changes and changes in AV conduction; heart block; arrhythmias; hypotension, particularly orthostatic hypotension; syncope; hypertension; tachycardia; palpitation.

CNS and Neuromuscular: Coma; seizures; hallucinations; delusions; confusional states; disorientation; incoordination; ataxia; tremors; peripheral neuropathy; numbness, tingling and paresthesias of the extremities; extrapyramidal symptoms including abnormal involuntary movements and tardive dyskinesia; dysarthria; disturbed concentration; excitement; anxiety; insomnia; restlessness; nightmares; drowsiness; dizziness; weakness; fatigue; headache; syndrome of inappropriate ADH (antidiuretic hormone) secretion; tinnitus; alteration in EEG patterns.

Anticholinergic: Paralytic ileus, hyperpyrexia; urinary retention, dilatation of the urinary tract; constipation; blurred vision, disturbance of accommodation, increased ocular pressure, mydriasis; dry mouth.

Allergic: Skin rash; urticaria; photosensitization; edema of face and tongue.

Hematologic: Bone marrow depression including agranulocytosis, leukopenia, thrombocytopenia; purpura; eosinophilia.

Gastrointestinal: Rarely hepatitis (including altered liver function and jaundice); nausea; epigastric distress; vomiting; anorexia; stomatitis; peculiar taste; diarrhea; parotid swelling; black tongue.

Endocrine: Testicular swelling and gynecomastia in the male; breast enlargement and galactorrhea in the female; increased or decreased libido; impotence; elevation and lowering of blood sugar levels.

Other: Alopecia; edema; weight gain or loss; urinary frequency; increased perspiration, hyponatremia.

Withdrawal Symptoms

After prolonged administration, abrupt cessation of treatment may produce nausea, headache, and malaise. Gradual dosage reduction has been reported to produce, within two weeks, transient symptoms including irritability, restlessness, and dream and sleep disturbance.

These symptoms are not indicative of addiction. Rare instances have been reported of mania or hypomania occurring within 2 to 7 days following cessation of chronic therapy with tricyclic antidepressants.

Causal Relationship Unknown

Other reactions, reported under circumstances where a causal relationship could not be established, are listed to serve as alerting information to physicians:

Body as a Whole: Lupus-like syndrome (migratory arthritis, positive ANA and rheumatoid factor).

Digestive: Hepatic failure, ageusia.

Postmarketing Adverse Events

A syndrome resembling neuroleptic malignant syndrome (NMS) has been very rarely reported after starting or increasing the dose of amitriptyline hydrochloride, with and without concomitant medications known to cause NMS. Symptoms have included muscle rigidity, fever, mental status changes, diaphoresis, tachycardia, and tremor.

Very rare cases of serotonin syndrome (SS) have been reported with amitriptyline hydrochloride in combination with other drugs that have a recognized association with SS.

To report SUSPECTED ADVERSE REACTIONS, contact Rising Pharma Holdings, Inc., at 1-844-874-7464 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Deaths may occur from overdosage with this class of drugs. Multiple drug ingestion (including alcohol) is common in deliberate tricyclic antidepressant overdose. As the management is complex and changing, it is recommended that the physician contact a poison control center for current information on treatment. Signs and symptoms of toxicity develop rapidly after tricyclic antidepressant overdose, therefore, hospital monitoring is required as soon as possible.

Manifestations

Critical manifestations of overdose include: cardiac dysrhythmias, severe hypotension, convulsions, and CNS depression, including coma. Changes in the electrocardiogram

particularly in QRS axis or width, are clinically significant indicators of tricyclic antidepressant toxicity. In addition, a rightward axis shift in the terminal QRS complex together with a prolonged QT interval and sinus tachycardia are specific and sensitive indicators of first generation tricyclic overdose. The absence of these findings is not exclusionary. Prolonged PR interval, ST-T wave changes, ventricular tachycardia and fibrillation may also occur.

Other signs of overdose may include: impaired myocardial contractility, confusion, disturbed concentration, transient visual hallucinations, dilated pupils, disorders of ocular motility, agitation, hyperactive reflexes polyradiculoneuropathy, stupor, drowsiness, muscle rigidity, vomiting, hypothermia, hyperpyrexia, or any of the symptoms listed under **ADVERSE REACTIONS**.

Management

General

Obtain an ECG and immediately initiate cardiac monitoring. Protect the patient's airway, establish an intravenous line and initiate gastric decontamination. A minimum of six hours of observation with cardiac monitoring and observation for signs of CNS or respiratory depression, hypotension, cardiac dysrhythmias and/or conduction blocks, and seizures is necessary. If signs of toxicity occur at any time during the period extended monitoring is required. There are case reports of patients succumbing to fatal dysrhythmias late after overdose; these patients had clinical evidence of significant poisoning prior to death and most received inadequate gastrointestinal decontamination. Monitoring of plasma drug levels should not guide management of the patient.

Gastrointestinal Decontamination

All patients suspected of tricyclic antidepressant overdose should receive gastrointestinal decontamination. This should include, large volume gastric lavage followed by activated charcoal. If consciousness is impaired, the airway should be secured prior to lavage. EMESIS IS CONTRAINDICATED.

Cardiovascular

A maximal limb-lead QRS duration of ≥ 0.10 seconds may be the best indication of the severity of the overdose. Intravenous sodium bicarbonate should be used to maintain the serum pH in the range of 7.45 to 7.55. If the pH response is inadequate, hyperventilation may also be used. Concomitant use of hyperventilation and sodium bicarbonate should be done with extreme caution, with frequent pH monitoring. A pH > 7.60 or a $pCO_2 < 20$ mm Hg is undesirable. Dysrhythmias unresponsive to sodium bicarbonate therapy/hyperventilation may respond to lidocaine, bretylium or phenytoin. Type 1 A and 1 C antiarrhythmics are generally contraindicated (e.g., quinidine, disopyramide, and procainamide).

In rare instances, hemoperfusion may be beneficial in acute refractory cardiovascular instability in patients with acute toxicity. However, hemodialysis, peritoneal dialysis, exchange transfusions, and forced diuresis generally have been reported as ineffective in tricyclic antidepressant poisoning.

CNS

In patients with CNS depression early intubation is advised because of the potential for abrupt deterioration. Seizures should be controlled with benzodiazepines, or if these are

ineffective, other anticonvulsants (e.g., phenobarbital, phenytoin).

Physostigmine is not recommended except to treat lifethreatening symptoms that have been unresponsive to other therapies, and then only in consultation with a poison control center.

Psychiatric Follow-up

Since overdose is often deliberate, patients may attempt suicide by other means during the recovery phase. Psychiatric referral may be appropriate.

Pediatric Management

The principles of management of pediatric and adult overdosages are similar. It is strongly recommended that the physician contact the local poison control center for specific pediatric treatment.

DOSAGE AND ADMINISTRATION

Oral Dosage

Dosage should be initiated at a low level and increased gradually, noting carefully the clinical response and any evidence of intolerance.

Initial Dosage for Adults

For outpatients, 75 mg of amitriptyline hydrochloride a day in divided doses is usually satisfactory. If necessary, this may be increased to a total of 150 mg per day. Increases are made preferably in the late afternoon and/or bedtime doses. A sedative effect may be apparent before the antidepressant effect is noted, but an adequate therapeutic effect may take as long as 30 days to develop.

An alternate method of initiating therapy in outpatients is to begin with 50 to 100 mg amitriptyline hydrochloride at bedtime. This may be increased by 25 or 50 mg as necessary in the bedtime dose to a total of 150 mg per day.

Hospitalized patients may require 100 mg a day initially. This can be increased gradually to 200 mg a day if necessary. A small number of hospitalized patients may need as much as 300 mg a day.

Adolescent and Elderly Patients

In general, lower dosages are recommended for these patients. Ten mg 3 times a day with 20 mg at bedtime may be satisfactory in adolescent and elderly patients who do not tolerate higher dosages.

Maintenance

The usual maintenance dosage of amitriptyline hydrochloride is 50 to 100 mg per day. In some patients, 40 mg per day is sufficient. For maintenance therapy, the total daily dosage may be given in a single dose, preferably at bedtime. When satisfactory improvement has been reached, dosage should be reduced to the lowest amount that will maintain relief of symptoms. It is appropriate to continue maintenance therapy 3 months or longer to lessen the possibility of relapse.

Usage in Pediatric Patients

In view of the lack of experience with the use of this drug in pediatric patients, it is not recommended at the present time for patients under 12 years of age.

Plasma Levels

Because of the wide variation in the absorption and distribution of tricyclic antidepressants in body fluids, it is difficult to directly correlate plasma levels and therapeutic effect. However, determination of plasma levels may be useful in identifying patients who appear to have toxic effects and may have excessively high levels, or those in whom lack of absorption or noncompliance is suspected. Because of increased intestinal transit time and decreased hepatic metabolism in elderly patients, plasma levels are generally higher for a given oral dose of amitriptyline hydrochloride than in younger patients. Elderly patients should be monitored carefully and quantitative serum levels obtained as clinically appropriate. Adjustments in dosage should be made according to the patient's clinical response and not on the basis of plasma levels. ²

²Hollister, L.E.; Monitoring Tricyclic Antidepressant Plasma Concentrations. JAMA 1979; 241(23):2530-2533.

HOW SUPPLIED

Amitriptyline hydrochloride tablets, USP for oral administration are available as:

The 50 mg tablets are Brown colored, round tablets, debossed with "32" on one side and plain on the other side. They are supplied as:

NDC: 70518-4048-00

NDC: 70518-4048-01

NDC: 70518-4048-02

PACKAGING: 90 in 1 BOTTLE PLASTIC

PACKAGING: 30 in 1 BLISTER PACK

PACKAGING: 30 in 1 BOTTLE PLASTIC

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container.

Repackaged and Distributed By:

Remedy Repack, Inc.

625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762

METABOLISM

Studies in man following oral administration of ¹⁴C-labeled drug indicated that amitriptyline hydrochloride is rapidly absorbed and metabolized. Radioactivity of the plasma was practically negligible, although significant amounts of radioactivity appeared in the urine by 4 to 6 hours and one-half to one-third of the drug was excreted within 24 hours.

Amitriptyline hydrochloride is metabolized by N-demethylation and bridge hydroxylation in man, rabbit, and rat. Virtually the entire dose is excreted as glucuronide or sulfate conjugate of metabolites, with little unchanged drug appearing in the urine. Other metabolic pathways may be involved.

REFERENCES

- Ayd, F.J., Jr.: Amitriptyline therapy for depressive reactions, *Psychosom.* 1: 320-325, Nov.-Dec. 1960.
- Diamond, S.: Human metabolism of amitriptyline tagged with carbon 14, *Curr. Therap. Res.* 7: 170-175, Mar. 1965.
- Dorfman, W.: Clinical experiences with amitriptyline (A preliminary report), *Psychosom.* 1: 153- 155, May-June 1960.
- Fallete, J.M.; Stasney, C.R.; Mintz, A.A.: Amitriptyline poisoning treated with physostigmine, *S. Med. J.* 63: 1492-1493, Dec. 1970 (in Soc. Proc.).
- Hollister, L.E.; Overall, J.E.; Johnson, M.; Pennington, V.; Katz, G.; Shelton, J.: Controlled comparison of amitriptyline, imipramine and placebo in hospitalized depressed patients, *J. Nerv. and Ment. Dis.* 139: 370-375, Oct. 1964.
- Hordern, A.; Burt, C.G.; Holt, N.F.: Depressive states. A pharmacotherapeutic study, Springfield, Ill., Charles C. Thomas, 1965.
- Jenike, M.A.: Treatment of Affective Illness in the Elderly with Drugs and Electroconvulsive Therapy, *J. Geriatr. Psychiatry* 1989;22(1).77-112.
- Klerman, G.L.; Cole, J.O.: Clinical pharmacology of imipramine and related antidepressant compounds, *Int. J. Psychiat.* 3: 267-304, Apr. 1976.
- Liu, B.; Anderson, C.; Mittman, N. et al: Use of selective serotonin-reuptake inhibitors or tricyclic antidepressants and risk of hip fractures in elderly people. *Lancet* 1998; 351 (9112):1303-1307.
- McConaghy, N.; Joffe, A.D.; Kingston, W.R.; Stevenson, H.G.; Atkinson, I.; Cole, E.; Fennessy, L.A.: Correlation of clinical features of depressed outpatients with response to amitriptyline and protriptyline, *Brit. J. Psychiat.* 114: 103-106, Jan. 1968.
- McDonald, I.M.; Perkins, M.; Marjerrison, G.; Podilsky, M.: A controlled comparison of amitriptyline and electroconvulsive therapy in the treatment of depression, *Amer. J. Psychiat.* 122: 1427-1431. June 1966 (in Brief Communications).
- Slovis, T.; Ott, J.; Teitelbaum, D.; Lipscomb, W.: Physostigmine therapy in acute tricyclic antidepressant poisoning, *Clin. Toxicol.* 4: 451-459, Sept. 1971.
- Symposium on depression with special studies of a new antidepressant, amitriptyline, *Dis. Nerv. Syst.* 22: 5-56, May 1961 (Sect. 2).

Dispense with Medication Guide available at:

<https://www.risingpharma.com/Medguides/AmitriptylineTabletsUSPMG.pdf>

Repackaged and Distributed By:

Remedy Repack, Inc.

MEDICATION GUIDE

Amitriptyline Hydrochloride

("am" i trip' tileen hye" droe klor' ide")

Tablets, USP

Dispense with Medication Guide available at:

<https://www.risingpharma.com/Medguides/AmitriptylineTabletsUSPMG.pdf>

Antidepressant Medicines, Depression and other Serious Mental Illnesses, and Suicidal Thoughts or Actions

Read the Medication Guide that comes with you or your family member's antidepressant medicine. This Medication Guide is only about the risk of suicidal thoughts and actions with antidepressant medicines. **Talk to your, or your family member's, healthcare provider about:**

- all risks and benefits of treatment with antidepressant medicines
- all treatment choices for depression or other serious mental illness

What is the most important information I should know about antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions?

1. **Antidepressant medicines may increase suicidal thoughts or actions in some children, teenagers, and young adults within the first few months of treatment.**
2. **Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions.** These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions.
3. **How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?**
 - Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed.
 - Call the healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.
 - Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

Call a healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety

- feeling very agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood
- **Low salt (sodium) levels in the blood. Elderly people may be at greater risk for this. Symptoms may include:**
 - Headache
 - weakness or feeling unsteady
 - confusion, problems concentrating or thinking or memory problems
 - **Visual problems:** eye pain, changes in vision, swelling or redness in or around the eye
- **Serotonin Syndrome.** This condition can be life-threatening and symptoms may include:
 - agitation, hallucinations, coma, or other changes in mental status
 - racing heartbeat, high or low blood pressure
 - coordination problems or muscle twitching (overactive reflexes)
 - nausea, vomiting, or diarrhea
 - sweating or fever
 - muscle rigidity

What else do I need to know about antidepressant medicines?

- **Never stop an antidepressant medicine without first talking to a healthcare provider.** Stopping an antidepressant medicine suddenly can cause other symptoms.
- **Visual problems:** Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.
- **Antidepressants are medicines used to treat depression and other illnesses.** It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.
- **Antidepressant medicines have other side effects.** Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member.
- **Antidepressant medicines can interact with other medicines.** Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking with your healthcare provider.
- **Not all antidepressant medicines prescribed for children are FDA approved for use in children.** Talk to your child's healthcare provider for more information.
- Tell your healthcare provider if you have low sodium levels in your blood during treatment with amitriptyline hydrochloride tablets

Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088.

This Medication Guide has been approved by the U.S. Food and Drug Administration for all antidepressants.

For more information, call Rising Pharma Holdings, Inc., at 1-844-874-7464.

Repackaged By / Distributed By: RemedyRepack Inc.

625 Kolter Drive, Indiana, PA 15701

(724) 465-8762

DRUG: Amitriptyline Hydrochloride

GENERIC: Amitriptyline Hydrochloride

DOSAGE: TABLET

ADMINISTRATION: ORAL

NDC: 70518-4048-0

NDC: 70518-4048-1

NDC: 70518-4048-2

COLOR: brown

SHAPE: ROUND

SCORE: No score

SIZE: 8 mm

IMPRINT: 32

PACKAGING: 90 in 1 BOTTLE PLASTIC

PACKAGING: 30 in 1 BLISTER PACK

PACKAGING: 30 in 1 BOTTLE PLASTIC

ACTIVE INGREDIENT(S):

- AMITRIPTYLINE HYDROCHLORIDE 50mg in 1

INACTIVE INGREDIENT(S):

- SILICON DIOXIDE
- CROSCARMELLOSE SODIUM
- HYDROXYPROPYL CELLULOSE (90000 WAMW)
- ISOPROPYL ALCOHOL
- LACTOSE MONOHYDRATE
- MAGNESIUM STEARATE
- MICROCRYSTALLINE CELLULOSE 101
- MICROCRYSTALLINE CELLULOSE 102
- STARCH, CORN
- HYPROMELLOSE 2910 (6 MPA.S)
- POLYETHYLENE GLYCOL 8000
- TITANIUM DIOXIDE

- FD&C BLUE NO. 2
- FD&C RED NO. 40
- FERRIC OXIDE RED

Amitriptyline HCl Tablet

NDC #: **70518-4048-01**
Expires:

LOT #:
Org NDC: 16571-0107-10
MFG: Rising Pharma, Inc.,
Allendale, NJ 07401

50 mg

QTY: 30 Tablets

Round BROWN 32

Keep this and all medication out
of the reach of children
Store at 20-25°C (68-77°F);
excursions permitted to 15-30°C
(59-86°F) [See USP]

Usual Dosage: See Insert



RX ONLY

Repackaged By:
RemedyRepack Inc.,
Indiana, PA 15701, 724.465.8762

LOT #:
NDC #:
MFG #:
EXP #:
MFG #:
EXP #:
MFG #:
EXP #:
MFG #:
EXP #:

Amitriptyline HCl Tablet

NDC #: **70518-4048-02**
Expires:

LOT #:
Org NDC: 16571-0107-10
MFG: Rising Pharma, Inc.,
Allendale, NJ 07401

50 mg

QTY: 30 Tablets

Round BROWN 32

Keep this and all medication out
of the reach of children
Store at 20-25°C (68-77°F);
excursions permitted to 15-30°C
(59-86°F) [See USP]

Usual Dosage: See Insert



RX ONLY

Repackaged By:
RemedyRepack Inc.,
Indiana, PA 15701, 724.465.8762

LOT #:
NDC #:
MFG #:
EXP #:
MFG #:
EXP #:
MFG #:
EXP #:
MFG #:
EXP #:

AMITRIPTYLINE HYDROCHLORIDE

amitriptyline hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70518-4048(NDC:16571- 107)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMITRIPTYLINE HYDROCHLORIDE (UNII: 26LUD4J09K) (AMITRIPTYLINE -	AMITRIPTYLINE	50 mg

UNII:1806D8D52K)	HYDROCHLORIDE	50 mg
------------------	---------------	-------

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYDROXYPROPYL CELLULOSE (90000 WAMW) (UNII: UKE75GEA7F)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE 101 (UNII: 7T9FYH5QMK)	
MICROCRYSTALLINE CELLULOSE 102 (UNII: PNR0YF693Y)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	32
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70518-4048-0	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/10/2024	04/11/2024
2	NDC:70518-4048-1	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	04/10/2024	
3	NDC:70518-4048-2	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/04/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA217411	04/10/2024	

Labeler - REMEDYREPACK INC. (829572556)