

**ICON 5000 II- iodine liquid
IBA**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ICON 5000 II - NPE FREE

ACTIVE INGREDIENTS:

Iodine Complex:.....25.00%

(Provide a full .5% available iodine when used at the recommended dilution ratio)

INERT INGREDIENTS:

Triple Emollient Syste (As Humectants And Skin
Conditioners):.....20.00%

(Provides a total of 2.0% emollients at the recommended dilution ratio)

Other
Inerts.....55.00%

(Includes Buffering Agents, Stabilizing Agents, And Surfactants)

TOTAL:.....100.00%

GENERAL RECOMMENDATIONS

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When soluiton in the teat dip cup becomes dirty and/or noticeably loses color. discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

CAUTION

KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID

FIRST AID: IF TAKEN INTERNALLY, DRINK LARGE QUANTITIES OF WATER AND GET IMMEDIATE MEDICAL ATTENTION. IN CASE OF CONTACT WITH EYES, FLUSH LIBERALLY WITH TEPID WATER AND GET MEDICAL ATTENTION. NEVER GIVE ANYHTING BY MOUTH TO AN UNCONSCIOUS PERSON.

FOR MEDICAL EMERGENCIES CALL 1-303-623-5716

RECOMMENDED PROCEDURES FOR TEAT SANITATION

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Onserve foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray teh cow's teats full length with diluted **ICON 5000 II** Teat Dip.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats and teat orifices with a clean, single service paper towel(s).
6. Attach milking unit

B.POST -MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with diluted **ICON 5000 II** Teat Dip. **Ensure good coverage of all sides of the teats and teat orifices.**

Dilute 1 part ICON 5000 II TEAT DIP per 9 parts potable , tepid water.

WARNING

WARNING: ICON 5000 II TEAT DIP CONCENTRATE MUST BE DILUTED BEFORE USE. DO NOT MIX WITH ANYTHING BUT POTABLE WATER. DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER. ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMGE TO THE TEATS.

Icon 5000 II NPE Free

NDC 29983-1018-9



A HIGHLY EFFECTIVE SANITIZING TEAT DIP

AN AID IN THE PREVENTION OF NEW INTRAMAMMARY INFECTIONS



IODINE TEAT DIP CONCENTRATE

Especially Formulated For Dairies Where **NPE FREE** Teat Dips Are Preferred Or Required

ACTIVE INGREDIENT:

Iodine Complex..... 25.00%
(Provides A Full .5% Available Iodine When Used At The Recommended Dilution Ratio)

INERT INGREDIENTS:

Triple Emollients (As Humectants And Skin Conditioners)..... 20.00%
(Provides A Total Of 2.0% Emollients At The Recommended Dilution Ratio)

Other Inerts..... 55.00%
(Includes Buffering Agents, Stabilizing Agents And Surfactants)

TOTAL:..... 100.00%

When mixed according to label directions, this product provides highly effective levels of free iodine for fast kill of mastitis causing organisms.

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR CONTACT WITH CONCENTRATED PRODUCTS, FLUSH THOROUGHLY WITH FRESH, TEPID WATER. DISCARD CONTAMINATED CLOTHING AND FOOTWEAR OR WASH BEFORE REUSE. FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, spray or foam the cow's teats full length with **DILUTED** ICON 5000 II NPE FREE TEAT DIP.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats and teat orifices with a clean, single service paper towel(s).
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with **DILUTED** ICON 5000 II NPE FREE TEAT DIP. Ensure good coverage of all sides of the teats and teat orifices.

CAUTION: KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

WARNING

ICON 5000 II NPE FREE TEAT DIP CONCENTRATE **MUST BE DILUTED** BEFORE USE. DO NOT MIX WITH ANYTHING BUT POTABLE WATER.

PROTECT THIS PRODUCT FROM FREEZING.

DO NOT RETURN DILUTED TEAT DIP TO STORAGE CONTAINER.

ALWAYS ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

Dilute 1 Part
ICON 5000 II NPE FREE
TEAT DIP
Per 9 Parts Potable,
Tepid Water.

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE: LOT NO.: NET CONTENTS: 275 GALLONS (1,040.875 LITERS) IBA STOCK # 261040 1121W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

ICON 5000 II

iodine liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1018
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.5 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1018-9	1188.75 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/27/2014	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 12/2025

IBA