


METHAZOLAMIDE- methazolamide tablet
Padagis US LLC

METHAZOLAMIDE TABLETS USP 25 mg and 50 mg
Rx Only

DESCRIPTION:

Methazolamide, a sulfonamide derivative, is a white crystalline powder, weakly acidic, slightly soluble in water, alcohol and acetone. The chemical name for methazolamide is: N-[5-(aminosulfonyl)- 3-methyl-1,3,4-thiadiazol- 2(3H)-ylidene]-acetamide and it has the following structural formula:



Each tablet, for oral administration, contains 25 mg or 50 mg methazolamide. In addition, each tablet contains the following inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate (powder), and microcrystalline cellulose.

CLINICAL PHARMACOLOGY:

Methazolamide is a potent inhibitor of carbonic anhydrase.

Methazolamide is well absorbed from the gastrointestinal tract. Peak plasma concentrations are observed 1 to 2 hours after dosing. In a multiple-dose, pharmacokinetic study, administration of methazolamide 25 mg bid, 50 mg bid, and 100 mg bid demonstrated a linear relationship between plasma methazolamide levels and methazolamide dose. Peak plasma concentrations (C_{max}) for the 25 mg, 50 mg and 100 mg bid regimens were 2.5 mcg/mL, 5.1 mcg/mL, and 10.7 mcg/mL, respectively. The area under the plasma concentration time curves (AUC) was 1130 mcg.min/mL, 2571 mcg.min/mL, and 5418 mcg.min/mL for the 25 mg, 50 mg, and 100 mg dosage regimens, respectively.

Methazolamide is distributed throughout the body including the plasma, cerebrospinal fluid, aqueous humor of the eye, red blood cells, bile and extracellular fluid. The mean apparent volume of distribution (V_{area}/F) ranges from 17 L to 23 L. Approximately 55% is bound to plasma proteins. The steady-state methazolamide red blood cell: plasma ratio varies with dose and was found to be 27:1, 16:1, and 10:1 following the

administration of methazolamide 25 mg bid, 50 mg bid, and 100 mg bid, respectively.

The mean steady-state plasma elimination half-life for methazolamide is approximately 14 hours. At steady-state, approximately 25% of the dose is recovered unchanged in the urine over the dosing interval. Renal clearance accounts for 20% to 25% of the total clearance of drug. After repeated bid-tid dosing, methazolamide accumulates to steady-state concentrations in 7 days.

Methazolamide's inhibitory action on carbonic anhydrase decreases the secretion of aqueous humor and results in a decrease in intraocular pressure. The onset of the decrease in intraocular pressure generally occurs within 2 to 4 hours, has a peak effect in 6 to 8 hours and a total duration of 10 to 18 hours.

Methazolamide is a sulfonamide derivative; however, it does not have any clinically significant antimicrobial properties. Although methazolamide achieves a high concentration in the cerebrospinal fluid, it is not considered an effective anticonvulsant.

Methazolamide has a weak and transient diuretic effect; therefore, use results in an increase in urinary volume, with excretion of sodium, potassium, and chloride. The drug should not be used as a diuretic. Inhibition of renal bicarbonate reabsorption produces an alkaline urine. Plasma bicarbonate decreases, and a relative, transient metabolic acidosis may occur due to a disequilibrium in carbon dioxide transport in the red blood cell. Urinary citrate excretion is decreased by approximately 40% after doses of 100 mg every 8 hours. Uric acid output has been shown to decrease 36% in the first 24 hour period.

INDICATIONS AND USAGE:

Methazolamide Tablets USP is indicated in the treatment of ocular conditions where lowering intraocular pressure is likely to be of therapeutic benefit, such as chronic open-angle glaucoma, secondary glaucoma, and preoperatively in acute angle-closure glaucoma where lowering the intraocular pressure is desired before surgery.

CONTRAINDICATIONS:

Methazolamide therapy is contraindicated in situations in which sodium and/or potassium serum levels are depressed, in cases of marked kidney or liver disease or dysfunction, in adrenal gland failure, and in hyperchloremic acidosis. In patients with cirrhosis, use may precipitate the development of hepatic encephalopathy.

Long-term administration of methazolamide is contraindicated in patients with angle-closure glaucoma, since organic closure of the angle may occur in spite of lowered intraocular pressure.

WARNINGS:

Fatalities have occurred, although rarely, due to severe reactions to sulfonamides including Stevens-Johnson syndrome, toxic epidermal necrolysis, fulminant hepatic necrosis, agranulocytosis, aplastic anemia, and other blood dyscrasias. Hypersensitivity reactions may recur when a sulfonamide is readministered, irrespective of the route of administration.

If hypersensitivity or other serious reactions occur, the use of this drug should be discontinued.

Caution is advised for patients receiving high-dose aspirin and methazolamide concomitantly, as anorexia, tachypnea, lethargy, coma, and death have been reported with concomitant use of high-dose aspirin and carbonic anhydrase inhibitors.

PRECAUTIONS:

General:

Potassium excretion is increased initially upon administration of methazolamide and in patients with cirrhosis or hepatic insufficiency could precipitate a hepatic coma.

In patients with pulmonary obstruction or emphysema, where alveolar ventilation may be impaired, methazolamide should be used with caution because it may precipitate or aggravate acidosis.

Information for Patients:

Adverse reactions common to all sulfonamide derivatives may occur: anaphylaxis, fever, rash (including erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis), crystalluria, renal calculus, bone marrow depression, thrombocytopenic purpura, hemolytic anemia, leukopenia, pancytopenia, and agranulocytosis. Precaution is advised for early detection of such reactions, and the drug should be discontinued and appropriate therapy instituted.

Caution is advised for patients receiving high-dose aspirin and methazolamide concomitantly.

Laboratory Tests:

To monitor for hematologic reactions common to all sulfonamides, it is recommended that a baseline CBC and platelet count be obtained on patients prior to initiating methazolamide therapy and at regular intervals during therapy. If significant changes occur, early discontinuance and institution of appropriate therapy are important. Periodic monitoring of serum electrolytes is also recommended.

Drug Interactions:

Methazolamide should be used with caution in patients on steroid therapy because of the potential for developing hypokalemia.

Caution is advised for patients receiving high-dose aspirin and methazolamide concomitantly, as anorexia, tachypnea, lethargy, coma and death have been reported with concomitant use of high-dose aspirin and carbonic anhydrase inhibitors (see **WARNINGS**).

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Long-term studies in animals to evaluate the carcinogenic potential of methazolamide and its effect on fertility have not been conducted. Methazolamide was not mutagenic in the Ames bacterial test.

Pregnancy:

Teratogenic effects.

Pregnancy Category C.

Methazolamide has been shown to be teratogenic (skeletal anomalies) in rats when given in doses approximately 40 times the human dose. There are no adequate and well controlled studies in pregnant women. Methazolamide should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers:

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from methazolamide, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use:

The safety and effectiveness of methazolamide in pediatric patients have not been established.

ADVERSE REACTIONS:

Adverse reactions, occurring most often early in therapy, include paresthesias, particularly a "tingling" feeling in the extremities; hearing dysfunction or tinnitus; fatigue; malaise; loss of appetite; taste alteration; gastrointestinal disturbances such as nausea, vomiting, and diarrhea; polyuria; and occasional instances of drowsiness and confusion.

Metabolic acidosis and electrolyte imbalance may occur.

Transient myopia has been reported. This condition invariably subsides upon diminution or discontinuance of the medication.

Other occasional adverse reactions include urticaria, melena, hematuria, glycosuria, hepatic insufficiency, flaccid paralysis, photosensitivity, convulsions, and, rarely, crystalluria and renal calculi. Also see **PRECAUTIONS: Information for Patients** for possible reactions common to sulfonamide derivatives. Fatalities have occurred, although rarely, due to severe reactions to sulfonamides including Stevens-Johnson syndrome, toxic epidermal necrolysis, fulminant hepatic necrosis, agranulocytosis, aplastic anemia, and other blood dyscrasias (see **WARNINGS**).

OVERDOSAGE:

No data is available regarding methazolamide overdosage in humans as no cases of acute poisoning with this drug have been reported. Animal data suggest that even a high dose of methazolamide is nontoxic. No specific antidote is known. Treatment should be symptomatic and supportive.

Electrolyte imbalance, development of an acidotic state, and central nervous system

effects might be expected to occur. Serum electrolyte levels (particularly potassium) and blood pH levels should be monitored. Supportive measures may be required to restore electrolyte and pH balance.

DOSAGE AND ADMINISTRATION:

The effective therapeutic dose administered varies from 50 mg to 100 mg two or three times daily. The drug may be used concomitantly with miotic and osmotic agents.

HOW SUPPLIED:

Methazolamide Tablets USP, 25 mg are round, white tablets, debossed "EFF" on one side and "21" on the other side and are supplied in bottles of 100, NDC 0574-**0790**-01.

Methazolamide Tablets USP, 50 mg are round, white, scored tablets debossed "EFF" on one side and "20" on the other side and are supplied in bottles of 100, NDC 0574-**0791**-01.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

Rx Only

Manufactured for Perrigo

Minneapolis, MN 55427

Code 617M00

Rev 04-14 A

5P800 RC J1

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL - 25 mg Label

Rx Only

NDC 0574-**0790**-01

Methazolamide Tablets, USP

25 mg

100 Tablets

<p>EACH TABLET CONTAINS: Methazolamide.....25 mg</p> <p>USUAL DOSAGE: 50 mg to 100 mg 2 or 3 times a day. See package insert.</p> <p>DISPENSE in a tight container as defined in the USP, with a child-resistant closure (as required).</p>	<p>STORE at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].</p> <p>WARNING: Keep this and all drugs out of the reach of children.</p> <p>Mfd for Perrigo Minneapolis, MN 55427 628L10 7F278 RC F2 Rev 10-16 B</p>	<p style="text-align: right;">Rx Only</p> <p>NDC 0574-0790-01</p> <p>Methazolamide Tablets, USP</p> <p>25 mg</p> <p>100 Tablets Perrigo®</p>	
<p>Non Varnish Area</p>			

The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]
 Lot [insert product's lot number]
 Exp [insert product's expiration date]

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL - 50 mg Label

Rx Only

NDC 0574-**0791**-01

Methazolamide Tablets, USP

50 mg

100 Tablets

<p>EACH TABLET CONTAINS: Methazolamide.....50 mg</p> <p>USUAL DOSAGE: 50 mg to 100 mg 2 or 3 times a day. See package insert.</p> <p>DISPENSE in a tight container as defined in the USP, with a child-resistant closure (as required).</p>	<p>STORE at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].</p> <p>WARNING: Keep this and all drugs out of the reach of children.</p> <p>Mfd for Perrigo Minneapolis, MN 55427 617M10 5P878 RC F2 Rev 10-16 B</p>	<p style="text-align: right;">Rx Only</p> <p>NDC 0574-0791-01</p> <p>Methazolamide Tablets, USP</p> <p>50 mg</p> <p>100 Tablets Perrigo®</p>	
<p>Non Varnish Area</p>			

The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]
 Lot [insert product's lot number]
 Exp [insert product's expiration date]

METHAZOLAMIDE				
methazolamide tablet				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0574-0790	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
METHAZOLAMIDE (UNII: W733B0S9SD) (METHAZOLAMIDE - UNII:W733B0S9SD)		METHAZOLAMIDE	25 mg	
Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
Product Characteristics				
Color	WHITE	Score	no score	
Shape	ROUND	Size	6mm	
Flavor		Imprint Code	EFF;21	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0574-0790-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040062	09/30/2014	

METHAZOLAMIDE

methazolamide tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0574-0791
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METHAZOLAMIDE (UNII: W733B0S9SD) (METHAZOLAMIDE - UNII:W733B0S9SD)	METHAZOLAMIDE	50 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	7mm
Flavor		Imprint Code	EFF;20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0574-0791-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040062	09/30/2014	

Labeler - Padagis US LLC (967694121)

Revised: 12/2021

Padagis US LLC