

**TOPCARE TRIPLE ANTIBIOTIC- bacitracin zinc, neomycin, polymyxin b,
pramoxine hydrochloride ointment
Topco Associates LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Triple Antibiotic Ointment Drug Facts

Active ingredients (in each gram)

Bacitracin zinc 500 units

Neomycin sulfate equivalent to 3.5 mg of neomycin base

Polymyxin B 10,000 units as polymyxin B sulfate

Pramoxine HCl 10 mg

Purpose

First aid antibiotic

External analgesic

Uses

first aid to help prevent infection and for temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 2 years of age or older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger)
on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredient

white petrolatum

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

COMPARE TO NEOSPORIN® + PAIN RELIEF

ACTIVE INGREDIENTS

First Aid Antibiotic & Pain Relieving Ointment

Helps Prevent Infection in Minor Cuts, Scrapes & Burns

Maximum Strength

TopCare® health

MAXIMUM STRENGTH

Triple Antibiotic Ointment + Pain Relief

BACITRACIN ZINC, NEOMYCIN SULFATE, POLYMYXIN B SULFATE & PRAMOXINE HCl

First Aid Antibiotic / Pain Relieving Ointment

NET WT 1 OZ (28 g)

COMPARE TO NEOSPORIN®
+ PAIN RELIEF
ACTIVE INGREDIENTS*

First Aid Antibiotic & Pain Relieving Ointment

Helps Prevent Infection in Minor Cuts,
Scrapes & Burns

Maximum Strength



MAXIMUM STRENGTH

Triple Antibiotic Ointment

+ Pain Relief

BACITRACIN ZINC, NEOMYCIN SULFATE, POLYMYXIN B SULFATE & PRAMOXINE HCl

First Aid Antibiotic / Pain Relieving Ointment

NET WT 1 OZ (28 g)

NDC 36800-790-64

DISTRIBUTED BY
TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE
IL 60007
©TOPCO PERA0521
QUESTIONS?
1-888-423-0139
topcare@topco.com
www.topcarebrand.com



Scan here for more
information or
call 1-888-423-0139



This TopCare® product is laboratory
tested to guarantee its highest quality.
Your total satisfaction is guaranteed.

*This product is not affiliated with,
manufactured by, or produced by
the makers of Neosporin®.



0 36800 24874 8
67964 88 C2



Triple Antibiotic Ointment + Pain Relief

Drug Facts

Active Ingredients (in each gram)	Purpose
Bacitracin zinc 500 units	First aid antibiotic
Neomycin sulfate equivalent to 35 mg of neomycin base	First aid antibiotic
Polymyxin B 10,000 units as polymyxin B sulfate	First aid antibiotic
Pramoxine HCl 10 mg	External analgesic

Uses First aid to help prevent infection and for temporary relief of pain or discomfort in minor:
 cuts scrapes burns

Warnings

For external use only.

- Do not use
- if you are allergic to any of the ingredients
 - in the eyes
 - over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Drug Facts (continued)

- Stop use and ask a doctor if
- you need to use longer than 1 week
 - condition persists or gets worse
 - symptoms persist for more than 1 week, or disappear again within a few days
 - rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 2 years of age or older
- clean the affected area
- apply as small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information Store at 20-25°C (68-77°F)

Inactive ingredient white petrolatum

Questions or comments? 1-888-423-0139

TOPCARE TRIPLE ANTIBIOTIC

bacitracin zinc, neomycin, polymyxin b, pramoxine hydrochloride ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-780	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)		BACITRACIN	500 [USP'U] in 1 g	
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)		NEOMYCIN	3.5 mg in 1 g	
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ.07J96K)		POLYMYXIN B	10000 [USP'U] in 1 g	
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)		PRAMOXINE HYDROCHLORIDE	10 mg in 1 g	
Inactive Ingredients				
Ingredient Name			Strength	
PETROLATUM (UNII: 4T6H12BN9U)				
Product Characteristics				
Color	YELLOW (Translucent)		Score	
Shape			Size	
Flavor			Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-780-64	1 in 1 CARTON	03/15/2020	
1		28 g in 1 TUBE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph final	part333B		03/15/2020	

Labeler - Topco Associates LLC (006935977)

Revised: 2/2022

Topco Associates LLC