

# **BUDESONIDE- budesonide capsule, coated pellets**

## **REMEDYREPACK INC.**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use BUDESONIDE DELAYED-RELEASE CAPSULES safely and effectively. See full prescribing information for BUDESONIDE DELAYED-RELEASE CAPSULES.**

**BUDESONIDE delayed-release capsules, for oral use**

**Initial U.S. Approval:1997**

### **RECENT MAJOR CHANGES**

Warnings and Precautions

Immunosuppression and Increased Risk of Infection ( 5.3) 06/2024

Kaposi's Sarcoma ( 5.4) 06/2024

### **INDICATIONS AND USAGE**

Budesonide delayed-release capsules are corticosteroid indicated for:

- Treatment of mild to moderate active Crohn's disease involving the ileum and/or the ascending colon, in patients 8 years and older. ( 1.1)
- Maintenance of clinical remission of mild to moderate Crohn's disease involving the ileum and/or the ascending colon for up to 3 months in adults. ( 1.2)

### **DOSAGE AND ADMINISTRATION**

#### **Administration Instructions ( 2.1)**

- Take once daily in the morning.
- Swallow whole. Do not chew or crush.
- For patients unable to swallow an intact capsule, open the capsules and empty the granules onto one tablespoonful of applesauce. Mix and consume the entire contents within 30 minutes. Do not chew or crush. Follow with 8 ounces of water.
- Avoid consumption of grapefruit juice for the duration of therapy.

#### **Recommended Dosage**

*Mild to moderate active Crohn's disease ( 2.2)*

- Adults: 9 mg once daily for up to 8 weeks; repeat 8 week treatment courses for recurring episodes of active disease.
- Pediatrics 8 to 17 years who weigh more than 25 kg: 9 mg once daily for up to 8 weeks, followed by 6 mg once daily in the morning for 2 weeks.

*Maintenance of clinical remission of mild to moderate Crohn's disease ( 2.3)*

- Adults: 6 mg once daily for up to 3 months; taper to complete cessation after 3 months. Continued treatment for more than 3 months has not been shown to provide substantial clinical benefit.
- When switching from oral prednisolone, begin tapering prednisolone concomitantly with initiating budesonide delayed-release capsules.

*Hepatic Impairment*

- Consider reducing the dosage to 3 mg once daily in adult patients with moderate hepatic impairment (Child-Pugh Class B). ( 2.4, 5.1, 8.6)

### **DOSAGE FORMS AND STRENGTHS**

Delayed-Release Capsules: 3 mg ( 3)

### **CONTRAINDICATIONS**

Hypersensitivity to budesonide or any of the ingredients in budesonide delayed-release capsules. ( 4)

### **WARNINGS AND PRECAUTIONS**

- *Hypercorticism and Adrenal Axis Suppression:* May occur with treatment; monitor for signs and symptoms; pediatrics and patients with hepatic impairment may be at increased risk. ( 2.4, 5.1, 8.4, 8.6)
- *Symptoms of Steroid Withdrawal in Patients Transferred from Other Systemic Corticosteroids:* Taper slowly from corticosteroids with high systemic effects; monitor for withdrawal symptoms and unmasking of allergies (rhinitis, eczema). ( 5.2)

- *Immunosuppression and Increased Risk of Infection*: Increased risk of viral, bacterial, fungal, protozoal and helminthic infections, including potentially fatal varicella and measles infection. Monitor patients for new or worsening infection and consider drug discontinuation. Avoid use in patients with fungal infections, *Strongyloides* infestation, cerebral malaria and ocular herpes simplex. Screen for hepatitis B infection. ( 5.3)
- *Kaposi's Sarcoma*: Reported to occur in patients receiving corticosteroid therapy, most often for chronic conditions. ( 5.4)
- *Other Corticosteroid Effects*: Monitor patients with concomitant conditions where corticosteroids may have unwanted effects (e.g., hypertension, diabetes mellitus). ( 5.5)

#### ----- ADVERSE REACTIONS -----

Most common adverse reactions ( $\geq 5\%$ ) in adults are: headache, respiratory infection, nausea, back pain, dyspepsia, dizziness, abdominal pain, flatulence, vomiting, fatigue and pain. ( 6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Northstar Rx LLC at 1-800-206-7821 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### ----- DRUG INTERACTIONS -----

**CYP3A4 Inhibitors (e.g., ketoconazole, grapefruit juice)**: Can increase systemic budesonide concentrations: avoid use. ( 2.1, 7.1)

#### ----- USE IN SPECIFIC POPULATIONS -----

**Pregnancy**: Based on animal data, may cause fetal harm. ( 8.1)

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 4/2026**

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

- 1.1 Treatment of Mild to Moderate Active Crohn's Disease
- 1.2 Maintenance of Clinical Remission of Mild to Moderate Crohn's Disease

### **2 DOSAGE AND ADMINISTRATION**

- 2.1 Administration Instructions
- 2.2 Treatment of Mild to Moderate Active Crohn's Disease
- 2.3 Maintenance of Clinical Remission of Mild to Moderate Crohn's Disease
- 2.4 Dosage Adjustment in Adult Patients with Hepatic Impairment

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Hypercorticism and Adrenal Axis Suppression
- 5.2 Symptoms of Steroid Withdrawal in Patients Transferred from Other Systemic Corticosteroids
- 5.3 Immunosuppression and Increased Risk of Infection
- 5.4 Kaposi's Sarcoma
- 5.5 Other Corticosteroid Effects

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

### **7 DRUG INTERACTIONS**

- 7.1 CYP3A4 Inhibitors

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

8.6 Hepatic Impairment

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

14.1 Treatment of Mild to Moderate Active Crohn's Disease

14.2 Maintenance of Clinical Remission of Mild to Moderate Crohn's Disease

## **15 REFERENCES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Treatment of Mild to Moderate Active Crohn's Disease**

Budesonide delayed-release capsules are indicated for the treatment of mild to moderate active Crohn's disease involving the ileum and/or the ascending colon in patients 8 years of age and older.

#### **1.2 Maintenance of Clinical Remission of Mild to Moderate Crohn's Disease**

Budesonide delayed-release capsules are indicated for the maintenance of clinical remission of mild to moderate Crohn's disease involving the ileum and/or the ascending colon for up to 3 months in adults.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Administration Instructions**

- Take budesonide delayed-release capsules once daily in the morning.
  - Swallow budesonide delayed-release capsules whole. Do not chew or crush.
  - For patients unable to swallow an intact capsule, budesonide delayed-release capsules can be opened and administered as follows:
1. Place one tablespoonful of applesauce into a clean container (e.g., empty bowl). The applesauce used should not be hot and should be soft enough to be swallowed without chewing.
  2. Open the capsule(s).
  3. Carefully empty all the granules inside the capsule(s) on the applesauce.
  4. Mix the granules with the applesauce.

5. Consume the entire contents within 30 minutes of mixing. Do not chew or crush the granules. Do not save the applesauce and granules for future use.
  6. Follow the applesauce and granules immediately with a glass (8 ounces) of cool water to ensure complete swallowing of the granules.
- Avoid consumption of grapefruit juice for the duration of budesonide delayed-release capsules therapy [see *Drug Interactions ( 7.1)*].

## **2.2 Treatment of Mild to Moderate Active Crohn's Disease**

The recommended dosage of budesonide delayed-release capsules is:

### *Adults*

9 mg orally once daily for up to 8 weeks. Repeated 8 week courses of budesonide delayed-release capsules can be given for recurring episodes of active disease.

### *Pediatric patients 8 to 17 years who weigh more than 25 kg*

9 mg orally once daily for up to 8 weeks, followed by 6 mg once daily for 2 weeks.

## **2.3 Maintenance of Clinical Remission of Mild to Moderate Crohn's Disease**

The recommended dosage in adults, following an 8 week course(s) of treatment for active disease and once the patient's symptoms are controlled (CDAI less than 150), is budesonide delayed-release capsules 6 mg orally once daily for maintenance of clinical remission up to 3 months. If symptom control is still maintained at 3 months an attempt to taper to complete cessation is recommended. Continued treatment with budesonide delayed-release capsules 6 mg for more than 3 months has not been shown to provide substantial clinical benefit.

Patients with mild to moderate active Crohn's disease involving the ileum and/or ascending colon have been switched from oral prednisolone to budesonide delayed-release capsules with no reported episodes of adrenal insufficiency. Since prednisolone should not be stopped abruptly, tapering should begin concomitantly with initiating budesonide delayed-release capsules treatment.

## **2.4 Dosage Adjustment in Adult Patients with Hepatic Impairment**

Consider reducing the dosage of budesonide delayed-release capsules to 3 mg once daily for adult patients with moderate hepatic impairment (Child-Pugh Class B). Avoid use in patients with severe hepatic impairment (Child-Pugh Class C) [see *Warnings and Precautions ( 5.1)*, *Use in Specific Populations ( 8.6)*].

## **3 DOSAGE FORMS AND STRENGTHS**

Budesonide Delayed-Release Capsules, 3 mg are white to off-white, free flowing pellets, filled in size '1' hard gelatin capsules having opaque light-orange colored cap printed with "720" in black ink and opaque white body.

## **4 CONTRAINDICATIONS**

Budesonide delayed-release capsules are contraindicated in patients with

hypersensitivity to budesonide or any of the ingredients of budesonide delayed-release capsules. Serious hypersensitivity reactions, including anaphylaxis have occurred [see *Adverse Reactions ( 6.2)*].

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hypercorticism and Adrenal Axis Suppression**

Systemic effects such as hypercorticism and adrenal axis suppression may occur with use of corticosteroids, including budesonide delayed-release capsules [see *Adverse Reactions ( 6.1)*, *Clinical Pharmacology ( 12.2)*]. Pediatric patients with Crohn's disease have a slightly higher systemic exposure of budesonide and increased cortisol suppression than adults with Crohn's disease [see *Use in Specific Populations ( 8.4)*, *Clinical Pharmacology ( 12.2)*]. Monitor patients for signs and symptoms of hypercorticism and adrenal axis suppression during treatment with budesonide delayed-release capsules.

Patients with moderate to severe hepatic impairment (Child-Pugh Class B and C respectively) could be at an increased risk of hypercorticism and adrenal axis suppression due to an increased systemic exposure of oral budesonide. Avoid use in patients with severe hepatic impairment (Child-Pugh Class C). Monitor for increased signs and/or symptoms of hypercorticism and consider reducing the dosage in patients with moderate hepatic impairment (Child-Pugh Class B) [see *Dosage and Administration ( 2.4)*, *Use in Specific Populations ( 8.6)*, *Clinical Pharmacology ( 12.3)*].

Corticosteroids, including budesonide delayed-release capsules, can reduce the response of the hypothalamus-pituitary-adrenal (HPA) axis to stress. In situations where patients are subject to surgery or other stress situations, supplementation with a systemic corticosteroid is recommended.

### **5.2 Symptoms of Steroid Withdrawal in Patients Transferred from Other Systemic Corticosteroids**

Monitor patients who are transferred from corticosteroid treatment with high systemic effects to corticosteroids with lower systemic availability, such as budesonide delayed-release capsules, since symptoms attributed to withdrawal of steroid therapy, including those of acute adrenal axis suppression or benign intracranial hypertension, may develop. Adrenocortical function monitoring may be required in these patients and the dose of corticosteroid treatment with high systemic effects should be reduced cautiously.

Replacement of systemic corticosteroids with budesonide delayed-release capsules may unmask allergies (e.g., rhinitis and eczema), which were previously controlled by the systemic drug.

### **5.3 Immunosuppression and Increased Risk of Infection**

Corticosteroids, including budesonide delayed-release capsules, suppress the immune system and increase the risk of infection with any pathogen, including viral, bacterial, fungal, protozoan, or helminthic pathogens. Corticosteroids can:

- Reduce resistance to new infections
- Exacerbate existing infections
- Increase the risk of disseminated infections

- Increase the risk of reactivation or exacerbation of latent infections
- Mask some signs of infection

Corticosteroid-associated infections can be mild but can be severe and at times fatal. The rate of infectious complications increases with increasing corticosteroid dosages.

Monitor patients for the development of infection and consider discontinuation of budesonide delayed-release capsules if the patient develops an infection while on treatment.

### ***Tuberculosis***

If budesonide delayed-release capsules is used in patients with latent tuberculosis or tuberculin reactivity, reactivation of tuberculosis may occur. Closely monitor such patients for reactivation. During prolonged budesonide delayed-release capsules therapy, patients with latent tuberculosis or tuberculin reactivity should receive chemoprophylaxis.

### ***Varicella Zoster and Measles Viral Infections***

Varicella and measles can have a serious or even fatal course in non-immune pediatric and adult patients taking corticosteroids, including budesonide delayed-release capsules. In corticosteroid-treated patients who have not had these diseases or are non-immune, particular care should be taken to avoid exposure to varicella and measles:

- If a budesonide delayed-release capsules-treated patient is exposed to varicella, prophylaxis with varicella zoster immune globulin may be indicated. If varicella develops, treatment with antiviral agents may be considered.
- If a budesonide delayed-release capsules-treated patient is exposed to measles, prophylaxis with immunoglobulin may be indicated.

### ***Hepatitis B Virus Reactivation***

Hepatitis B virus reactivation can occur in patients who are hepatitis B carriers treated with immunosuppressive dosages of corticosteroids, including budesonide delayed-release capsules. Reactivation can also occur infrequently in corticosteroid-treated patients who appear to have resolved hepatitis B infection.

Screen patients for hepatitis B infection before initiating immunosuppressive (e.g., prolonged) treatment with budesonide delayed-release capsules. For patients who show evidence of hepatitis B infection, recommend consultation with physicians with expertise in managing hepatitis B regarding monitoring and consideration for hepatitis B antiviral therapy.

### ***Fungal Infections***

Corticosteroids, including budesonide delayed-release capsules, may exacerbate systemic fungal infections; therefore, avoid budesonide delayed-release capsules use in the presence of such infections. For patients on chronic budesonide delayed-release capsules therapy who develop systemic fungal infections, budesonide delayed-release capsules withdrawal or dosage reduction is recommended.

### ***Amebiasis***

Corticosteroids, including budesonide delayed-release capsules, may activate latent amebiasis. Therefore, it is recommended that latent amebiasis or active amebiasis be ruled out before initiating budesonide delayed-release capsules in patients who have spent time in the tropics or patients with unexplained diarrhea.

### ***Strongyloides Infestation***

Avoid budesonide delayed-release capsules in patients with known or suspected *Strongyloides* (threadworm) infection. Corticosteroid-induced immunosuppression may lead to *Strongyloides* superinfection and dissemination with widespread larval migration, often accompanied by severe enterocolitis and potentially fatal gram-negative septicemia.

### ***Cerebral Malaria***

Avoid corticosteroids, including budesonide delayed-release capsules, in patients with cerebral malaria.

### ***Ocular Herpes Simplex***

Avoid corticosteroids, including budesonide delayed-release capsules, in patients with active ocular herpes simplex.

### **5.4 Kaposi's Sarcoma**

Kaposi's sarcoma has been reported to occur in patients receiving corticosteroid therapy, most often for chronic conditions. Discontinuation of corticosteroids may result in clinical improvement of Kaposi's sarcoma.

### **5.5 Other Corticosteroid Effects**

Monitor patients with hypertension, diabetes mellitus, osteoporosis, peptic ulcer, glaucoma or cataracts, or with a family history of diabetes or glaucoma, or with any other condition where corticosteroids may have unwanted effects.

## **6 ADVERSE REACTIONS**

The following clinically significant adverse reactions are described elsewhere in labeling:

- Hypercorticism and adrenal axis suppression [*see Warnings and Precautions (5.1)*]
- Symptoms of steroid withdrawal in those patients transferred from other systemic corticosteroids [*see Warnings and Precautions ( 5.2)*]
- Immunosuppression and increased risk of infection [*see Warnings and Precautions ( 5.3)*]
- Kaposi's sarcoma [*see Warnings and Precautions ( 5.4)*]
- Other corticosteroid effects [*see Warnings and Precautions ( 5.5)*]

### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### ***Adults***

The data described below reflect exposure to budesonide delayed-release capsules in 520 patients with Crohn's disease, including 520 exposed to 9 mg per day (total daily dose) for 8 weeks and 145 exposed to 6 mg per day for one year in placebo controlled clinical trials. Of the 520 patients, 38% were males and the age range was 17 to 74 years.

## Treatment of Mild to Moderate Active Crohn's Disease

The safety of budesonide delayed-release capsules was evaluated in 651 adult patients in five clinical trials of 8 weeks duration in patients with active mild to moderate Crohn's disease. The most common adverse reactions, occurring in greater than or equal to 5% of the patients, are listed in Table 1.

**Table 1 Common Adverse Reactions <sup>1</sup>in 8-Week Treatment Clinical Trials**

<b>Adverse Reaction</b>	<b>Budesonide Delayed-Release Capsules 9 mg n=520 Number (%)</b>	<b>Placebo n=107 Number (%)</b>	<b>Prednisolone <sup>2</sup> 40 mg n=145 Number (%)</b>	<b>Comparator <sup>3</sup> n=88 Number (%)</b>
Headache	107(21)	19(18)	31(21)	11(13)
Respiratory Infection	55(11)	7(7)	20(14)	5(6)
Nausea	57(11)	10(9)	18(12)	7(8)
Back Pain	36(7)	10(9)	17(12)	5(6)
Dyspepsia	31(6)	4(4)	17(12)	3(3)
Dizziness	38(7)	5(5)	18(12)	5(6)
Abdominal Pain	32(6)	18(17)	6(4)	10(11)
Flatulence	30(6)	6(6)	12(8)	5(6)
Vomiting	29(6)	6(6)	6(4)	6(7)
Fatigue	25(5)	8(7)	11(8)	0(0)
Pain	24(5)	8(7)	17(12)	2(2)

<sup>1</sup>Occurring in greater than or equal to 5% of the patients in any treated group.

<sup>2</sup>Prednisolone tapering scheme: either 40 mg in week 1 to 2, thereafter tapering with 5 mg per week; or 40 mg in week 1 to 2, 30 mg in week 3 to 4, thereafter tapering with 5 mg per week.

<sup>3</sup>This drug is not approved for the treatment of Crohn's disease in the United States.

The incidence of signs and symptoms of hypercorticism reported by active questioning of patients in 4 of the 5 short-term clinical trials are displayed in Table 2.

**Table 2 Summary and Incidence of Signs/Symptoms of Hypercorticism in 8-Week Treatment Clinical Trials**

<b>Signs/Symptom</b>	<b>Budesonide Delayed-Release Capsules 9 mg n=427 Number (%)</b>	<b>Placebo n=107 Number (%)</b>	<b>Prednisolone <sup>1</sup> 40 mg n=145 Number (%)</b>
Total	145 (34%)	29 (27%)	69 (48%)
Acne	63(15)	14(13)	33(23) <sup>2</sup>
Bruising Easily	63(15)	12(11)	13(9)
Moon Face	46(11)	4(4)	53(37) <sup>2</sup>
Swollen Ankles	32(7)	6(6)	13(9)

Hirsutism <sup>3</sup>	22(5)	2(2)	5(3)
Buffalo hump	6(1)	2(2)	5(3)
Skin Striae	4(1)	2(2)	0(0)

<sup>1</sup> Prednisolone tapering scheme: either 40 mg in week 1 to 2, thereafter tapering with 5 mg/week; or 40 mg in week 1 to 2, 30 mg in week 3 to 4, thereafter tapering with 5 mg/week.

<sup>2</sup> Statistically significantly different from budesonide delayed-release capsules 9 mg.

<sup>3</sup> including hair growth increased, local and hair growth increased, general.

### *Maintenance of Clinical Remission of Mild to Moderate Crohn's Disease*

The safety of budesonide delayed-release capsules was evaluated in 233 adult patients in four long-term clinical trials (52 weeks) of maintenance of clinical remission in patients with mild to moderate Crohn's disease. A total of 145 patients were treated with budesonide delayed-release capsules 6 mg once daily.

The adverse reaction profile of budesonide delayed-release capsules 6 mg once daily in maintenance of Crohn's disease was similar to that of short-term treatment with budesonide delayed-release capsules 9 mg once daily in active Crohn's disease. In the long-term clinical trials, the following adverse reactions occurred in greater than or equal to 5% and are not listed in Table 1: diarrhea (10%); sinusitis (8%); infection viral (6%); and arthralgia (5%).

Signs/symptoms of hypercorticism reported by active questioning of patients in the long-term maintenance clinical trials are displayed in Table 3.

**Table 3 Summary and Incidence of Signs/Symptoms of Hypercorticism in Long-Term Clinical Trials**

	<b>Budesonide Delayed-Release Capsules 3 mg n=88</b>	<b>Budesonide Delayed-release Capsules 6 mg n=145</b>	<b>Placebo n=143</b>
<b>Signs/Symptom</b>	<b>Number (%)</b>	<b>Number (%)</b>	<b>Number (%)</b>
Bruising easily	4(5)	15(10)	5(4)
Acne	4(5)	14(10)	3(2)
Moon face	3(3)	6(4)	0
Hirsutism	2(2)	5(3)	1(1)
Swollen ankles	2(2)	3(2)	3(2)
Buffalo hump	1(1)	1(1)	0
Skin striae	2(2)	0	0

The incidence of signs/symptoms of hypercorticism as described above in long-term maintenance clinical trials was similar to that seen in the short-term treatment clinical trials.

### *Less Common Adverse Reactions in Treatment and Maintenance Clinical Trials*

Less common adverse reactions (less than 5%), occurring in adult patients treated with budesonide delayed-release capsules 9 mg (total daily dose) in short-term treatment clinical studies and/or budesonide delayed-release capsules 6 mg (total daily dose) in long-term maintenance clinical trials, with an incidence are listed below by system organ

class:

*Cardiac disorders:* palpitation, tachycardia

*Eye disorders:* eye abnormality, vision abnormal

*General disorders and administration site conditions:* asthenia, chest pain, dependent edema, face edema, flu-like disorder, malaise, fever

*Gastrointestinal disorders:* anus disorder, enteritis, epigastric pain, gastrointestinal fistula, glossitis, hemorrhoids, intestinal obstruction, tongue edema, tooth disorder

*Infections and infestations:* Ear infection - not otherwise specified, bronchitis, abscess, rhinitis, urinary tract infection, thrush

*Investigations:* weight increased

*Metabolism and nutrition disorders:* appetite increased

*Musculoskeletal and connective tissue disorders:* arthritis, cramps, myalgia

*Nervous system disorders:* hyperkinesia, paresthesia, tremor, vertigo, somnolence, amnesia

*Psychiatric disorders:* agitation, confusion, insomnia, nervousness, sleep disorder

*Renal and urinary disorders:* dysuria, micturition frequency, nocturia

*Reproductive system and breast disorders:* intermenstrual bleeding, menstrual disorder

*Respiratory, thoracic and mediastinal disorders:* dyspnea, pharynx disorder

*Skin and subcutaneous tissue disorders:* alopecia, dermatitis, eczema, skin disorder, sweating increased, purpura

*Vascular disorders:* flushing, hypertension

### *Bone Mineral Density*

A randomized, open, parallel-group multicenter safety clinical trial specifically compared the effect of budesonide delayed-release capsules (less than 9 mg per day) and prednisolone (less than 40 mg per day) on bone mineral density over 2 years when used at doses adjusted to disease severity. Bone mineral density decreased significantly less with budesonide delayed-release capsules than with prednisolone in steroid-naïve patients, whereas no difference could be detected between treatment groups for steroid-dependent patients and previous steroid users. The incidence of symptoms associated with hypercorticism was significantly higher with prednisolone treatment.

### *Clinical Laboratory Test Findings*

The following potentially clinically significant laboratory changes in clinical trials, irrespective of relationship to budesonide delayed-release capsules, were reported in greater than or equal to 1% of patients: hypokalemia, leukocytosis, anemia, hematuria, pyuria, erythrocyte sedimentation rate increased, alkaline phosphatase increased, atypical neutrophils, c-reactive protein increased and adrenal insufficiency.

### ***Pediatrics-Treatment of Mild to Moderate Active Crohn's Disease***

Adverse reactions reported in pediatric patients 8 to 17 years of age, who weigh more than 25 kg, were similar to those reactions described above in adult patients.

## 6.2 Postmarketing Experience

The following adverse reactions have been reported during post-approval use of budesonide delayed-release capsules. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Immune System Disorders:*Anaphylactic reactions

*Nervous System Disorders:*Benign intracranial hypertension

*Psychiatric Disorders:*Mood swings

## 7 DRUG INTERACTIONS

### 7.1 CYP3A4 Inhibitors

Budesonide is a substrate for CYP3A4. Avoid use with CYP3A4 inhibitors. Concomitant oral administration of a strong CYP3A4 inhibitor (ketoconazole) caused an eight-fold increase of the systemic exposure to oral budesonide. Inhibitors of CYP3A4 (e.g., ketoconazole, itraconazole, ritonavir, indinavir, saquinavir, erythromycin, and cyclosporine) can increase systemic budesonide concentrations [see *Clinical Pharmacology* ( 12.3)] .

#### **Grapefruit Juice**

Avoid ingestion of grapefruit juice with budesonide. Intake of grapefruit juice which inhibits CYP3A4 activity with budesonide can increase the systemic exposure for budesonide [see *Clinical Pharmacology* ( 12.3)] .

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### **Risk Summary**

Limited published studies report on the use of budesonide in pregnant women; however, the data are insufficient to inform a drug-associated risk for major birth defects and miscarriage. There are clinical considerations [see *Clinical Considerations*]. In animal reproduction studies with pregnant rats and rabbits, administration of subcutaneous budesonide during organogenesis at doses approximately 0.5 times or 0.05 times, respectively, the maximum recommended human dose, resulted in increased fetal loss, decreased pup weights, and skeletal abnormalities. Maternal toxicity was observed in both rats and rabbits at these dose levels [ see *Data*]. Based on animal data, advise pregnant women of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage of the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

#### **Clinical Considerations**

*Disease-Associated Maternal and/or Embryo/Fetal Risk*

Some published epidemiological studies show an association of adverse pregnancy outcomes in women with Crohn's disease, including preterm birth and low birth weight infants, during periods of increased disease activity (including increased stool frequency and abdominal pain). Pregnant women with Crohn's disease should be counseled regarding the importance of controlling disease.

#### *Fetal/Neonatal adverse reactions*

Hypoadrenalism may occur in infants born of mothers receiving corticosteroids during pregnancy. Infants should be carefully observed for signs of hypoadrenalism, such as poor feeding, irritability, weakness, and vomiting, and managed accordingly [see *Warnings and Precautions ( 5.1)*].

### **Data**

#### *Animal Data*

Budesonide was teratogenic and embryolethal in rabbits and rats.

In an embryo-fetal development study in pregnant rats dosed subcutaneously with budesonide during the period of organogenesis from gestation days 6 to 15 there were effects on fetal development and survival at subcutaneous doses up to approximately 500 mcg/kg in rats (approximately 0.5 times the maximum recommended human dose on a body surface area basis). In an embryo-fetal development study in pregnant rabbits dosed during the period of organogenesis from gestation days 6 to 18, there was an increase in maternal abortion, and effects on fetal development and reduction in litter weights at subcutaneous doses up to approximately 25 mcg/kg in rabbits (approximately 0.05 times the maximum recommended human dose on a body surface area basis). Maternal toxicity, including reduction in body weight gain, was observed at subcutaneous doses of 5 mcg/kg in rabbits (approximately 0.01 times the maximum recommended human dose on a body surface area basis) and 500 mcg/kg in rats (approximately 0.5 times the maximum recommended human dose on a body surface area basis).

In a peri- and post-natal development study, rats dosed subcutaneously with budesonide during the period of Day 15 post coitum to Day 21 postpartum, budesonide had no effects on delivery but did have an effect on growth and development of offspring. In addition, offspring survival was reduced and surviving offspring had decreased mean body weights at birth and during lactation at exposures 0.02 times the MRHD (on a mg/m<sup>2</sup> basis at maternal subcutaneous doses of 20 mcg/kg/day and higher). These findings occurred in the presence of maternal toxicity.

## **8.2 Lactation**

### ***Risk Summary***

Lactation studies have not been conducted with oral budesonide, including budesonide delayed-release capsules, and no information is available on the effects of the drug on the breastfed infant or the effects of the drug on milk production. One published study reports that budesonide is present in human milk following maternal inhalation of budesonide [ see *Data*]. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for budesonide delayed-release capsules and any potential adverse effects on the breastfed infant from budesonide delayed-release capsules, or from the underlying maternal condition.

### **Data**

One published study reports that budesonide is present in human milk following maternal inhalation of budesonide which resulted in infant doses approximately 0.3% to 1% of the maternal weight-adjusted dosage and a milk/plasma ratio ranging between 0.4 and 0.5. Budesonide plasma concentrations were not detected and no adverse events were noted in the breastfed infants following maternal use of inhaled budesonide. The recommended daily dose of budesonide delayed-release capsules is higher (up to 9 mg daily) compared with inhaled budesonide (up to 800 mcg daily) given to mothers in the above described study. The maximum budesonide plasma concentration following a 9 mg daily dose (in both single- and repeated-dose pharmacokinetic studies) of oral budesonide is approximately 2.15 to 4.31 ng/mL which is up to 10 times higher than the 0.43 to 0.86 ng/mL for a 800 mcg daily dose of inhaled budesonide at steady state in the above inhalation study. Assuming the coefficient of extrapolation between the inhaled and oral doses is constant across all dose levels, at therapeutic doses of budesonide delayed-release capsules, budesonide exposure to the nursing child may be up to 10 times higher than that by budesonide inhalation.

#### **8.4 Pediatric Use**

The safety and effectiveness of budesonide delayed-release capsules have been established in pediatric patients 8 to 17 years of age who weigh more than 25 kg for the treatment of mild to moderate active Crohn's disease involving the ileum and/or the ascending colon. Use of budesonide delayed-release capsules in this age group is supported by evidence from adequate and well controlled studies of budesonide delayed-release capsules in adults, with additional data from 2 clinical studies in 149 pediatric patients treated up to 8 weeks and one pharmacokinetic study in 8 pediatric patients [see *Adverse Reactions ( 6.1)*, *Clinical Pharmacology ( 12.3)*, and *Clinical Studies ( 14.1)*].

The observed safety profile of budesonide delayed-release capsules in pediatric patients is consistent with its known safety profile in adults and no new safety concerns were identified [see *Adverse Reactions ( 6.1)*].

The safety and effectiveness of budesonide delayed-release capsules have not been established in pediatric patients less than 8 years of age for the treatment of mild to moderate active Crohn's disease involving the ileum and/or the ascending colon.

The safety and effectiveness of budesonide delayed-release capsules have not been established in pediatric patients for the maintenance of clinical remission of mild to moderate Crohn's disease. An open-label study to evaluate the safety and tolerability of budesonide delayed-release capsules as maintenance treatment in pediatric patients aged 5 to 17 years was conducted, and did not establish the safety and efficacy of maintenance of clinical remission.

Systemic corticosteroids, including budesonide delayed-release capsules, may cause a reduction of growth velocity in pediatric patients. Pediatric patients with Crohn's disease have a 17% higher mean systemic exposure and cortisol suppression than adults with Crohn's disease [see *Warnings and Precautions ( 5.1)* and *Clinical Pharmacology ( 12.2)*].

#### **8.5 Geriatric Use**

Clinical studies of budesonide delayed-release capsules did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. Of the 651 patients treated with budesonide delayed-release

capsules in clinical studies, 17 (3%) were greater than or equal to 65 years of age and none were greater than 74 years of age. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## **8.6 Hepatic Impairment**

Patients with moderate to severe hepatic impairment (Child-Pugh Class B and C, respectively) could be at an increased risk of hypercorticism and adrenal axis suppression due to an increased systemic exposure to budesonide [see *Warnings and Precautions (5.1) and Clinical Pharmacology (12.3)*]. Avoid use in patients with severe hepatic impairment (Child-Pugh Class C). Monitor for increased signs and/or symptoms of hypercorticism and consider dosage reduction in patients with moderate hepatic impairment (Child-Pugh Class B) [see *Dosage and Administration (2.4)*]. No dosage adjustment is needed in patients with mild hepatic impairment (Child-Pugh Class A).

## **10 OVERDOSAGE**



Reports of acute toxicity and/or death following overdosage of glucocorticoids are rare. Treatment consists of immediate gastric lavage or emesis followed by supportive and symptomatic therapy.

If corticosteroids are used at excessive doses for prolonged periods, systemic corticosteroid effects such as hypercorticism and adrenal axis suppression may occur. For chronic overdosage in the case of severe disease requiring continuous steroid therapy, the dosage may be reduced temporarily.

Single oral doses of 200 and 400 mg/kg were lethal in female and male mice, respectively. The signs of acute toxicity were decreased motor activity, piloerection and generalized edema.

## **11 DESCRIPTION**

Budesonide USP, the active ingredient of budesonide delayed-release capsules, is a synthetic corticosteroid. Budesonide is designated chemically as (RS)-11 $\beta$ , 16 $\alpha$ , 17,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16,17-acetal with butyraldehyde. Budesonide is provided as a mixture of two epimers (22R and 22S). The molecular formula of budesonide is C<sub>25</sub>H<sub>34</sub>O<sub>6</sub> and its molecular weight is 430.5. Its structural formula is:



Budesonide, USP is a white to off-white, odorless, crystalline powder that is practically insoluble in water and heptane, sparingly soluble in ethanol, and freely soluble in chloroform. Its partition coefficient between octanol and water at pH 5 is  $1.6 \times 10^3$  ionic strength 0.01.

Each capsule for oral administration contains 3 mg of micronized budesonide with the following inactive ingredients: acetyltributyl citrate, ethylcellulose aqueous dispersion, gelatin, iron oxide red, iron oxide yellow, methacrylic acid copolymer dispersion, polysorbate 80, simethicone emulsion, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide and triethyl citrate. The capsule shell is printed with black pharmaceutical ink which contains following ingredients: iron oxide black, potassium hydroxide, propylene glycol and shellac.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Budesonide is an anti-inflammatory corticosteroid and has a high glucocorticoid effect and a weak mineralocorticoid effect, and the affinity of budesonide to glucocorticoid receptors, which reflects the intrinsic potency of the drug, is about 200-fold that of cortisol and 15-fold that of prednisolone.

### 12.2 Pharmacodynamics

Treatment with glucocorticoids, including budesonide delayed-release capsules is associated with a suppression of endogenous cortisol concentrations and an impairment of the hypothalamus-pituitary-adrenal (HPA) axis function. There was a positive correlation between the percent (%) reduction of  $AUC_{0-24}$  of plasma cortisol and systemic exposure to budesonide both in pediatric and adult patients.

#### **Adults**

Plasma cortisol suppression was compared following five days' administration of budesonide delayed-release capsules and prednisolone in a crossover study in healthy volunteers. The mean decrease in the area under the plasma cortisol concentration-time

curve over 24 hour (AUC<sub>0-24</sub>) was greater (78%) with prednisolone 20 mg per day compared to 45% with budesonide delayed-release capsules 9 mg per day.

### **Pediatrics**

The effect of budesonide on endogenous cortisol concentrations was compared between pediatrics (n=8, aged 9 to 14 years) and adults (n=6) with active Crohn's disease following administration of budesonide delayed-release capsules 9 mg once daily for 7 days. Compared to baseline values before treatment, the mean decrease in the AUC<sub>0-24</sub> of cortisol was 64% (±18%) in pediatrics and 50% (±27%) in adults after budesonide delayed-release capsules treatment [see *Warnings and Precautions ( 5.1)*, *Adverse Reactions ( 6.1)* and *Use in Specific Populations ( 8.4)*].

The responses to adrenocorticotropin challenge (i.e., ACTH stimulation test) was studied in pediatric patients aged 8 to 17 years, with mild to moderate active Crohn's disease in randomized, double-blind, active control study [see *Clinical Studies ( 14.1)*]. After 8 weeks of treatment with 9 mg once daily budesonide delayed-release capsules or with prednisolone, administered at tapering doses starting from 1 mg/kg, the proportion of patients with normal response to the ACTH challenge was 6% in the budesonide group compared to none in the prednisolone group; the proportion of patients with morning p-cortisol of greater than 5 mcg/dL was 50% in the budesonide group compared to 22% in the prednisolone group. The mean morning p-cortisol was 6.3 mcg/dL in the budesonide group and 2.6 mcg/dL in the prednisolone group (Table 4).

**Table 4 Proportion of Pediatric Patients 8 to 17 years old with Peak Endogenous Cortisol Levels (above 18 mcg/dL) after ACTH Stimulation and Normal Response\* to ACTH Challenge Following Administration of Budesonide Delayed-Release Capsules or Prednisolone for 8 weeks**

	Budesonide	Prednisolone
Peak plasma cortisol above 18 mcg/dL		
At baseline	91% (20/22)	91% (21/23)
At week 8	25% (4/16)	0% (0/18)
Normal response* to ACTH challenge		
At baseline	73% (16/22)	78% (18/23)
At week 8	6% (1/16)	0% (0/18)

\*The normal response to ACTH challenge included 3 criteria, as defined in the cosyntropin label: 1) morning cortisol level above 5 mcg/dL; 2) increase in cortisol level by at least 7 mcg/dL above the morning (pre-challenge) level following ACTH challenge; and cortisol level of above 18 mcg/dL following ACTH challenge. Cortisol concentration was measured at 30 min after intravenous or intramuscular injection of 0.25 mg cosyntropin at baseline and at week 8 after treatment.

## **12.3 Pharmacokinetics**

### **Absorption**

Mean oral bioavailability of budesonide ranged from 9% to 21% both in patients and in healthy subjects, demonstrating a high first-pass elimination of the drug.

Budesonide pharmacokinetics were dose-proportional following repeated administration in the dose range of 3 to 15 mg. No accumulation of budesonide was observed following repeated dosing.

Following oral administration of a single dose of 9 mg budesonide delayed-release

capsules in healthy subjects under fasting condition, the mean peak plasma concentration ( $C_{max}$ ) and the area under the plasma concentration time curve (AUC) for budesonide were  $1.50 \pm 0.79$  ng/mL and  $14.13 \pm 7.33$  ng•hr/mL, respectively. The time to peak concentration ( $T_{max}$ ) varied between 2 and 8 hours with a median value of 3.5 hours. In a different study, following oral administration of 9 mg budesonide delayed-release capsules for five days in healthy subjects, the mean  $C_{max}$  and the steady state AUC for budesonide were  $2.28 \pm 0.77$  ng/mL and  $15.93 \pm 6.29$  ng•hr/mL, respectively.

Following administration of 9 mg budesonide delayed-release capsules once daily in patients with active Crohn's disease, the mean  $C_{max}$  and AUC were  $1.7 \pm 0.9$  ng/mL and  $15.1 \pm 8.5$  ng•hr/mL, respectively. Following administration of budesonide delayed-release capsules, the  $T_{max}$  ranged in individual patients from 0.5 to 10 hours.

Concomitant administration of a high-fat meal delayed the  $T_{max}$  of budesonide by 2.3 hours compared to that under fasted conditions but did not significantly affect the AUC in healthy subjects. The mean  $C_{max}$  and AUC of budesonide were similar when single dose of budesonide delayed-release capsules (9 mg) was administered after opening the capsules and sprinkling the granules on applesauce versus as intact capsules in the fasted state (N=24) in healthy subjects. The  $T_{max}$  ranged from 3 to 10 hours with a median of 4 hours after administration of sprinkled granules on applesauce.

### **Distribution**

The mean volume of distribution ( $V_{ss}$ ) of budesonide varied between 2.2 and 3.9 L/kg in healthy subjects and in patients. Plasma protein binding was estimated to be 85% to 90% in the concentration range 0.43 to 99.02 ng/mL, independent of gender. The erythrocyte/plasma partition ratio at clinically relevant concentrations was about 0.8.

### **Elimination**

Budesonide had a plasma clearance, 0.9 to 1.8 L/min in healthy adults. Mean plasma clearance after intravenous administration of budesonide in patients with Crohn's disease was 1.0 L/min. These plasma clearance values approached the estimated liver blood flow, and, accordingly, suggest that budesonide is a high hepatic clearance drug. The plasma elimination half-life after administration of intravenous doses ranged between 2 and 3.6 hours, and did not differ between healthy adults and patients with Crohn's disease. The mean  $\pm$  SD plasma elimination half-life after a single dose of budesonide delayed-release capsules (9 mg) in the fasted state (N=24) in healthy subjects was  $6.3 \pm 1.6$  hours and ranged between 2 and 8 hours.

### **Metabolism**

Following absorption, budesonide is subject to high first pass metabolism (80% to 90%). *In vitro* experiments in human liver microsomes demonstrated that budesonide is rapidly and extensively biotransformed, mainly by CYP3A4, to its 2 major metabolites, 6 $\beta$ -hydroxy budesonide and 16 $\alpha$ -hydroxy prednisolone. The corticosteroid activity of these metabolites was negligible (less than 1/100) in relation to that of the parent compound. *In vivo* investigations with intravenous doses in healthy subjects were in agreement with the *in vitro* findings.

### **Excretion**

Budesonide was excreted in urine and feces in the form of metabolites. After oral as well as intravenous administration of micronized [ $^3$ H]-budesonide, approximately 60% of the recovered radioactivity was found in urine. The major metabolites, including 6 $\beta$ -hydroxy

budesonide and 16 $\alpha$ -hydroxy prednisolone, are mainly renally excreted, intact or in conjugated forms. No unchanged budesonide was detected in urine.

### **Specific Populations**

#### *Age: Pediatric Population (8 years and older)*

The pharmacokinetics of budesonide were investigated in pediatric patients aged 9 to 14 years (n=8) after oral administration of budesonide delayed-release capsules and intravenous administration of budesonide. Following administration of 9 mg budesonide delayed-release capsules once daily for 7 days, the median time to peak plasma concentration of budesonide was 5 hours and the mean peak plasma concentration was 2.58  $\pm$  1.51 ng/mL. The mean AUC was 17.78  $\pm$  5.25 ng•hr/mL and 17% higher than that in adult patients with Crohn's disease in the same study. The mean absolute oral availability was 9.2% (3 to 17%; n=4) in pediatric patients.

After single dose administration of intravenous budesonide (n=4), the mean volume of distribution ( $V_{SS}$ ) was 2.2  $\pm$  0.4 L/kg and mean clearance was 0.81  $\pm$  0.2 L/min. The mean elimination half-life was 1.9 hours in pediatric patients. The body-weight normalized clearance in pediatric patients was 20.5 mL/min/kg in comparison to 15.9 mL/min/kg in adult patients after intravenous administration [see *Warnings and Precautions ( 5.1)*, *Use in Specific Population ( 8.4)*].

#### *Hepatic Impairment*

In patients with mild (Child-Pugh Class A, n=4) or moderate (Child-Pugh Class B, n=4) hepatic impairment, budesonide 4 mg was administered orally as a single dose. The patients with moderate hepatic impairment had a 3.5-fold higher AUC compared to the healthy subjects with normal hepatic function while the patients with mild hepatic impairment had an approximately 1.4-fold higher AUC. The  $C_{max}$  values demonstrated similar increases [see *Dosage and Administration ( 2.4)*, *Warnings and Precautions ( 5.1)*]. The increased systemic exposure in patients with mild hepatic impairment was not considered to be clinically relevant. Patients with severe liver impairment (Child-Pugh Class C) were not studied.

### **Drug Interaction Studies**

Budesonide is metabolized via CYP3A4. Potent inhibitors of CYP3A4 can increase the plasma concentrations of budesonide several-fold. Conversely, induction of CYP3A4 potentially could result in the lowering of budesonide plasma concentrations.

#### *Effects of Other Drugs on Budesonide*

##### *Ketoconazole*

In an open, non-randomized, cross-over study, 6 healthy subjects were given budesonide 10 mg as a single dose, either alone or concomitantly with the last ketoconazole dose of 3 days treatment with ketoconazole 100 mg twice daily. Co-administration of ketoconazole resulted in an eight-fold increase in AUC of budesonide, compared to budesonide alone [see *Drug Interactions ( 7.1)*].

##### *Grapefruit Juice*

In an open, randomized, cross-over study, 8 healthy subjects were given budesonide delayed-release capsules 3 mg, either alone, or concomitantly with 600 mL concentrated grapefruit juice (which inhibits CYP3A4 activity predominantly in the intestinal mucosa), on the last of 4 daily administrations. Concomitant administration of grapefruit juice

resulted in a 2-fold increase of the bioavailability of budesonide compared to budesonide alone [see *Drug Interactions ( 7.1)*].

#### *Oral Contraceptives (CYP3A4 Substrates)*

In a parallel study, the pharmacokinetics of budesonide were not significantly different between healthy female subjects who received oral contraceptives containing desogestrel 0.15 mg and ethinyl estradiol 30 µg and healthy female subjects who did not receive oral contraceptives. Budesonide 4.5 mg once daily (one-half the recommended dose) for one week did not affect the plasma concentrations of ethinyl estradiol, a CYP3A4 substrate. The effect of budesonide 9 mg once daily on the plasma concentrations of ethinyl estradiol was not studied.

#### *Omeprazole*

In a study in 11 healthy subjects, performed in a double-blind, randomized, placebo controlled manner, the effect of 5 to 6 days treatment with omeprazole 20 mg once daily on the pharmacokinetics of budesonide administered as budesonide delayed-release capsules 9 mg as a single dose was investigated. Omeprazole 20 mg once daily did not affect the absorption or pharmacokinetics of budesonide.

#### *Cimetidine*

In an open, non-randomized, cross-over study, the potential effect of cimetidine on the pharmacokinetics of budesonide was studied. Six healthy subjects received cimetidine 1 gram daily (200 mg with meals and 400 mg at night) for 2 separate 3-day periods. Budesonide 4 mg was administered either alone or on the last day of one of the cimetidine treatment periods. Co-administration of cimetidine resulted in a 52% and 31% increase in the budesonide peak plasma concentration and the AUC of budesonide, respectively.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Carcinogenicity studies with budesonide were conducted in rats and mice. In a two-year study in Sprague-Dawley rats, budesonide caused a statistically significant increase in the incidence of gliomas in male rats at an oral dose of 50 mcg/kg (approximately 0.05 times the maximum recommended human dose on a body surface area basis). In addition, there were increased incidences of primary hepatocellular tumors in male rats at 25 mcg/kg (approximately 0.023 times the maximum recommended human dose on a body surface area basis) and above. No tumorigenicity was seen in female rats at oral doses up to 50 mcg/kg (approximately 0.05 times the maximum recommended human dose on a body surface area basis). In an additional two-year study in male Sprague-Dawley rats, budesonide caused no gliomas at an oral dose of 50 mcg/kg (approximately 0.05 times the maximum recommended human dose on a body surface area basis). However, it caused a statistically significant increase in the incidence of hepatocellular tumors at an oral dose of 50 mcg/kg (approximately 0.05 times the maximum recommended human dose on a body surface area basis). The concurrent reference corticosteroids (prednisolone and triamcinolone acetonide) showed similar findings. In a 91-week study in mice, budesonide caused no treatment-related carcinogenicity at oral doses up to 200 mcg/kg (approximately 0.1 times the maximum recommended human dose on a body surface area basis).

Budesonide was not genotoxic in the Ames test, the mouse lymphoma cell forward gene mutation (TK <sup>+/-</sup>) test, the human lymphocyte chromosome aberration test, the *Drosophila melanogaster* sex-linked recessive lethality test, the rat hepatocyte UDS test and the mouse micronucleus test.

In rats, budesonide had no effect on fertility at subcutaneous doses up to 80 mcg/kg (approximately 0.07 times the maximum recommended human dose on a body surface area basis). However, it caused a decrease in prenatal viability and viability in pups at birth and during lactation, along with a decrease in maternal body-weight gain, at subcutaneous doses of 20 mcg/kg (approximately 0.02 times the maximum recommended human dose on a body surface area basis) and above. No such effects were noted at 5 mcg/kg (approximately 0.005 times the maximum recommended human dose on a body surface area basis).

## 14 CLINICAL STUDIES

### 14.1 Treatment of Mild to Moderate Active Crohn's Disease

#### Adults

The efficacy of budesonide delayed-release capsules were evaluated in 994 patients with mild to moderate active Crohn's disease of the ileum and/or ascending colon in 5 randomized and double-blind studies of 8 weeks duration. The study patients ranged in age from 17 to 85 (mean 35), 40% were male and 97% were white. The Crohn's Disease Activity Index (CDAI) was the main clinical assessment used for determining efficacy in these 5 studies. <sup>1</sup>The CDAI is a validated index based on subjective aspects rated by the patient (frequency of liquid or very soft stools, abdominal pain rating and general well-being) and objective observations (number of extraintestinal symptoms, need for antidiarrheal drugs, presence of abdominal mass, body weight and hematocrit). Clinical improvement, defined as a CDAI score of less than or equal to 150 assessed after 8 weeks of treatment, was the primary efficacy variable in these 5 comparative efficacy studies of budesonide delayed-release capsules. Safety assessments in these studies included monitoring of adverse reactions. A checklist of potential symptoms of hypercorticism was used.

One study (Study 1) compared the efficacy of budesonide delayed-release capsules 9 mg daily in the morning to a comparator. At baseline, the median CDAI was 272. budesonide delayed-release capsules 9 mg daily resulted in a significantly higher clinical improvement rate at Week 8 than the comparator. See Table 5.

**Table 5 Clinical Improvement Rates (CDAI less than or equal to 150) After 8 weeks of Treatment**

Clinical Study	Budesonide Delayed-Release Capsules 9 mg Daily	Budesonide Delayed-Release Capsules 4.5 mg Twice Daily	Comparator <sup>3</sup>	Placebo	Prednisolone
1	62/91 (69%) <sup>1</sup>		37/83 (45%)		
2		31/61 (51%) <sup>2</sup>		13/64 (20%)	

3	38/79 (48%)	41/78 (53%)		13/40 (33%)	
4	35/58 (60%)	25/60 (42%)			35/58 (60%)
5	45/86 (52%)				56/85 (65%)

<sup>1</sup>. p=0.0004 compared to comparator.

<sup>2</sup>. p=0.001 compared to placebo.

<sup>3</sup>. This drug is not approved for the treatment of Crohn's disease in the United States.

Two placebo-controlled clinical trials (Studies 2 and 3) were conducted. Study 2 involved 258 patients and tested the effects of graded doses of budesonide delayed-release capsules (1.5 mg twice daily, 4.5 mg twice daily, or 7.5 mg twice daily) versus placebo. At baseline, the median CDAI was 290. The 1.5 mg twice daily arm (data not shown) could not be differentiated from placebo. The 4.5 mg twice daily arm was statistically different from placebo (Table 5), while no additional benefit was seen when the daily budesonide delayed-release capsules dose was increased to 15 mg per day (data not shown). Study 3 was a 3-armed parallel group study. The groups were treated with budesonide delayed-release capsules 9 mg once daily, budesonide delayed-release capsules 4.5 mg twice daily and placebo for 8 weeks, followed by a 2-week double-blind taper phase. The median CDAI at baseline was 263. Neither 9 mg daily nor 4.5 mg twice daily budesonide delayed-release capsules dose levels were statistically different from placebo (Table 5). The recommended dosage of budesonide delayed-release capsules for the treatment of mild to moderate active Crohn's disease involving the ileum and/or the ascending colon in adults is 9 mg once daily in the morning for up to 8 weeks [see *Dosage and Administration ( 2.1)*].

Two clinical trials (Studies 4 and 5) compared budesonide delayed-release capsules with oral prednisolone (initial dose 40 mg per day). Study 4 was a 3-armed parallel group study. The groups were treated with budesonide delayed-release capsules 9 mg once daily, budesonide delayed-release capsules 4.5 mg twice daily and prednisolone 40 mg (tapered dose) for 8 weeks, followed by a 4-week double blind taper phase. At baseline, the median CDAI was 277. Equal clinical improvement rates (60%) were seen in the budesonide delayed-release capsules 9 mg daily and the prednisolone groups in Study 4. In Study 5, 13% fewer patients in the budesonide delayed-release capsules group experienced clinical improvement than in the prednisolone group (no statistical difference) (Table 5).

The proportion of patients with normal plasma cortisol values (greater than 64.58 ng/mL) was significantly higher in the budesonide delayed-release capsules groups in both trials (60% to 66%) than in the prednisolone groups (26% to 28%) at Week 8.

### ***Pediatrics (8 to 17 Years of Age)***

The effectiveness of budesonide delayed-release capsules, in pediatric patients aged 8 to 17 years, who weigh more than 25 kg with mild to moderate active Crohn's disease (defined as Crohn's Disease Activity Index (CDAI)  $\geq$  200) involving the ileum and/or the ascending colon, was assessed in one randomized, double-blind, active control study. This study compared budesonide delayed-release capsules 9 mg once daily, with prednisolone, administered at tapering doses starting from 1 mg/kg. Twenty-two (22) patients were treated with budesonide delayed-release capsules and 24 patients were treated with prednisolone. After 8 weeks of treatment, 55% (95% CI: 32%, 77%) of

patients treated with budesonide delayed-release capsules reached the endpoint (CDAI  $\leq$ 150), as compared to 68% (95% CI: 47%, 89%) of patients treated with prednisolone. The average number of liquid or very soft stools per day (assessed over 7 days) decreased from 1.49 at baseline to 0.96 after treatment with budesonide delayed-release capsules and 2.00 at baseline to 0.52 after treatment with prednisolone. The average daily abdominal pain rating (where 0=none, 1=mild, 2=moderate, and 3=severe) decreased from 1.49 at baseline to 0.54 after treatment with budesonide delayed-release capsules and 1.64 at baseline to 0.38 after 8 weeks of treatment with prednisolone.

Use of budesonide delayed-release capsules in this age group is supported by evidence from adequate and well-controlled studies of budesonide delayed-release capsules in adults, and by safety and pharmacokinetic studies performed in pediatric patients.

## **14.2 Maintenance of Clinical Remission of Mild to Moderate Crohn's Disease**

### ***Adults***

The efficacy of budesonide delayed-release capsules for maintenance of clinical remission were evaluated in four double-blind, placebo-controlled, 12-month trials in which 380 patients were randomized and treated once daily with 3 mg or 6 mg budesonide delayed-release capsules or placebo. Patients ranged in age from 18 to 73 (mean 37) years. Sixty percent of the patients were female and 99% were Caucasian. The mean CDAI at entry was 96. Among the four clinical trials, approximately 75% of the patients enrolled had exclusively ileal disease. Colonoscopy was not performed following treatment. Budesonide delayed-release capsules 6 mg per day prolonged the time to relapse, defined as an increase in CDAI of at least 60 units to a total score greater than 150 or withdrawal due to disease deterioration. The median time to relapse in the pooled population of the 4 studies was 154 days for patients taking placebo, and 268 days for patients taking budesonide delayed-release capsules 6 mg per day. Budesonide delayed-release capsules 6 mg per day reduced the proportion of patients with loss of symptom control relative to placebo in the pooled population for the 4 studies at 3 months (28% versus 45% for placebo).

## **15 REFERENCES**

1. Best WR, Beckett JM, Singleton JW, Kern F: Development of a Crohn's Disease Activity Index, National Cooperative Crohn's Disease Study. *Gastroenterology* 1976; 70(3): 439-444.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Budesonide Delayed-Release Capsules 3 mg are white to off-white, free flowing pellets, filled in size '1' hard gelatin capsules having opaque light-orange colored cap printed with "720" in black ink and opaque white body and are supplied as follows:

NDC: 70518-4613-00

NDC: 70518-4613-01

OUTER PACKAGING: 50 in 1 BOX

PACKAGING: 1 in 1 POUCH

Store at 20°C to 25°C (68°F to 77°F) [See USP Controlled Room Temperature].

Keep container tightly closed.

Repackaged and Distributed By:

Remedy Repack, Inc.

625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762

## **17 PATIENT COUNSELING INFORMATION**

Advise Patients to read the FDA-Approved patient labeling (Patient Information).

### ***Hypercorticism and Adrenal Axis Suppression***

Advise patients that budesonide delayed-release capsules may cause hypercorticism and adrenal axis suppression and to follow a taper schedule, as instructed by their healthcare provider if transferring to budesonide delayed-release capsules from systemic corticosteroids [*see Warnings and Precautions ( 5.1, 5.2)*]. Advise patients that replacement of systemic corticosteroids with budesonide delayed-release capsules may unmask allergies (e.g., rhinitis and eczema), which were previously controlled by the systemic drug.

### ***Immunosuppression and Increased Risk of Infection***

Advise patients to avoid exposure to people with varicella (chicken pox) or measles. Advise patients inform their healthcare provider if they are exposed to varicella or measles or if they develop a new or worsening infection [*see Warnings and Precautions ( 5.3)*].

### ***Kaposi's Sarcoma***

Advise patients that Kaposi's sarcoma has been reported in patients receiving corticosteroids for chronic conditions and to inform their healthcare provider if they experience signs or symptoms of Kaposi's sarcoma [*see Warnings and Precautions ( 5.4)*].

### ***Pregnancy***

Advise female patients that budesonide delayed-release capsules may cause fetal harm and to inform their healthcare provider with a known or suspected pregnancy [*see Use in Specific Populations ( 8.1)*].

### ***Administration***

- Take budesonide delayed-release capsules once daily in the morning.
  - Swallow budesonide delayed-release capsules whole. Do not chew or crush.
  - For patients unable to swallow an intact capsule, budesonide delayed-release capsules can be opened and administered as follows:
1. Place one tablespoonful of applesauce into a clean container (e.g., empty bowl). The applesauce used should not be hot and should be soft enough to be swallowed without chewing.
  2. Open the capsule(s).
  3. Carefully empty all the granules inside the capsule(s) on the applesauce.
  4. Mix the granules with the applesauce.
  5. Consume the entire contents within 30 minutes of mixing. Do not chew or crush the granules. Do not save the applesauce and granules for future use.

6. Follow the applesauce and granules immediately with a glass (8 ounces) of cool water to ensure complete swallowing of the granules.
- Avoid consumption of grapefruit juice for the duration of their budesonide delayed-release capsules therapy [see *Drug Interactions* ( 7.1)].

**Repackaged and Distributed By:**

**Remedy Repack, Inc.**

**625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762**

**SPL PATIENT PACKAGE INSERT**

<b>PATIENT INFORMATION</b>
<b>Budesonide (bue des' oh nide) Delayed-Release Capsules</b>
Read this Patient Information before you start taking budesonide delayed-release capsules and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.
<b>What are budesonide delayed-release capsules ?</b> Budesonide delayed-release capsules are prescription corticosteroid medicine used to treat mild to moderate Crohn's disease that affects part of the small intestine (ileum) and part of the large intestine (ascending colon): <ul style="list-style-type: none"><li>• in people 8 years of age and older with active Crohn's disease</li><li>• in adults to help keep symptoms from coming back for up to 3 months</li></ul>
It is not known if budesonide delayed-release capsules are safe and effective in children under 8 years of age, or in children 8 to 17 years of age who weigh 55 pounds (25 kg) or less, for the treatment of mild to moderate active Crohn's disease that affects part of the small intestine (ileum) and part of the large intestine (ascending colon). It is not known if budesonide delayed-release capsules are safe and effective in children to help keep symptoms of mild to moderate Crohn's disease that affects part of the small intestine (ileum) and part of the large intestine (ascending colon) from coming back.
<b>Who should not take budesonide delayed-release capsules?</b> <b>Do not take budesonide delayed-release capsules if:</b> <ul style="list-style-type: none"><li>• you are allergic to budesonide or any of the ingredients in budesonide delayed-release capsules. See the end of this leaflet for a complete list of ingredients in budesonide delayed-release capsules.</li></ul>
<b>Before you take budesonide delayed-release capsules tell your healthcare provider if you have any other medical conditions including if you:</b> <ul style="list-style-type: none"><li>• have liver problems.</li><li>• are planning to have surgery.</li><li>• have chicken pox or measles or have recently been near anyone with chicken pox or measles.</li><li>• have an infection, including fungal and threadworm (Strongyloides)</li></ul>

infections.

- have diabetes or glaucoma or have a family history of diabetes or glaucoma.
- have cataracts.
- have or had tuberculosis.
- have high blood pressure (hypertension).
- have decreased bone mineral density (osteoporosis).
- have stomach ulcers.
- have malaria of the brain (cerebral malaria).
- are pregnant or plan to become pregnant. Budesonide delayed-release capsules may harm your unborn baby. Talk to your healthcare provider about the possible risk to your unborn baby if you take budesonide delayed-release capsules when you are pregnant. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during your treatment with budesonide delayed-release capsules.
- are breastfeeding or plan to breastfeed. It is not known if budesonide passes into your breast milk or if it will affect your baby. Talk to your healthcare provider about the best way to feed your baby if you take budesonide delayed-release capsules.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Budesonide delayed-release capsules and other medicines may affect each other causing side effects.

#### **How should I take budesonide delayed-release capsules ?**

- Take budesonide delayed-release capsules exactly as your healthcare provider tells you.
  - Your healthcare provider will tell you how many budesonide delayed-release capsules to take. Your healthcare provider may change your dose if needed.
  - Take budesonide delayed-release capsules 1 time each day in the morning.
  - Take budesonide delayed-release capsules whole. Do not chew or crush budesonide delayed-release capsules before swallowing.
  - For patients, unable to swallow a whole capsule, budesonide delayed-release capsules can be opened and administered as follows:
1. Place 1 tablespoonful of applesauce into a clean container, such as an empty bowl. The applesauce used should not be hot and should be soft enough to be swallowed without chewing.
  2. Open the capsule. You may need to use more than 1 budesonide delayed-release capsules for the dose prescribed by your healthcare provider.
  3. Carefully empty all of the granules inside the capsule on the applesauce.
  4. Stir the granules with the applesauce.
  5. Swallow the applesauce and granules mixture within 30 minutes after preparing it. Follow the applesauce and granules immediately with a glass (8 ounces) of cool water to help with complete swallowing of the granules.
  6. Do not chew or crush the granules.

7. Do not save the applesauce and granules for later use.

- If you take too much budesonide delayed-release capsules call your healthcare provider right away or go to the nearest hospital emergency room.

### **What should I avoid while taking budesonide delayed-release capsules?**

- Do not drink grapefruit juice during your treatment with budesonide delayed-release capsules. Drinking grapefruit juice can increase the level of budesonide in your blood.

### **What are the possible side effects of budesonide delayed-release capsules?**

**Budesonide delayed-release capsules may cause serious side effects, including:**

- **Effects of having too much corticosteroid medicine in your blood (hypercorticism).** Long-time use of budesonide delayed-release capsules can cause you to have too much corticosteroid medicine in your blood. Tell your healthcare provider if you have any of the following signs and symptoms of hypercorticism:
  - acne
  - thicker or more hair on your body and face
  - bruise easily
  - a fatty pad or hump between your shoulders (buffalo hump)
  - rounding of your face (moon face)
  - pink or purple stretch marks on the skin of your abdomen, thighs, breasts and arms
  - ankle swelling
- **Adrenal suppression.** When budesonide delayed-release capsules are taken for a long period of time (chronic use), adrenal suppression can happen. This is a condition in which the adrenal glands do not make enough steroid hormones. Symptoms of adrenal suppression include: tiredness, weakness, nausea and vomiting and low blood pressure. Tell your healthcare provider if you are under stress or have any symptoms of adrenal suppression during treatment with budesonide delayed-release capsules.
- **Worsening of allergies.** If you take certain other corticosteroid medicines to treat allergies, switching to budesonide delayed-release capsules may cause your allergies to come back. These allergies may include a skin condition called eczema or inflammation inside your nose (rhinitis). Tell your healthcare provider if any of your allergies become worse while taking budesonide delayed-release capsules.
- **Decreased ability of your body to fight infections (immunosuppression) and increased risk of infection.** Corticosteroid medicines, including budesonide delayed-release

capsules, lower the ability of your immune system to fight infections and increase the risk of infections caused by virus, bacteria, fungi, protozoan, or certain parasites. Corticosteroid medicines including budesonide delayed-release capsules can also:

- o make current infections worse
- o increase the risk of infections spreading (disseminated)
- o increase the risk of making infections active again or making infections worse that have not been active (latent)
- o hide (mask) some signs of infection

These infections can be mild, but can be severe and lead to death. Your healthcare provider should check you closely for signs and symptoms of an infection while taking budesonide delayed-release capsules. Tell your healthcare provider right away about any signs or symptoms of a new or worsening infection while taking budesonide delayed-release capsules, including flu-like symptoms such as:

- o fever
- o cough
- o chills
- o pain
- o stomach area (abdominal) pain
- o feeling tired
- o aches
- o nausea and vomiting
- o diarrhea

o Tuberculosis: If you have inactive (latent) tuberculosis, your tuberculosis may become active again while taking budesonide delayed-release capsules your healthcare provider should check you closely for signs and symptoms of tuberculosis while taking budesonide delayed-release capsules.

o Chicken pox and measles: People taking corticosteroid medicines, including E budesonide delayed-release capsules, who have not had chicken pox or measles, should avoid contact with people who have these diseases. Tell your healthcare provider right away if you come in contact with anyone who has chicken pox or measles.

o Hepatitis B Virus (HBV) reactivation: If you are a carrier of HBV, the virus can become an active infection again while taking budesonide delayed-release capsules. Your healthcare provider will test you for HBV before you start taking budesonide delayed-release capsules.

o Amebiasis: Inactive (latent) amebiasis before you start taking budesonide delayed-release capsules. Your healthcare provider should check you for amebiasis before you start taking budesonide delayed-release capsules in people who have spent time in the tropics or have unexplained diarrhea.

- **Kaposi's Sarcoma.** Kaposi's sarcoma has happened in people who receive corticosteroid therapy, most often for treatment of long-lasting (chronic) conditions.

**The most common side effects of budesonide delayed-release capsules in adults include:**

- headache
- vomiting

- stomach area (abdominal) pain
- infection in your air passages (respiratory infection)
- gas
- nausea
- back pain
- tiredness
- indigestion
- pain
- dizziness

**The most common side effects of budesonide delayed-release capsules in children**

8 to 17 years of age, who weigh more than 55 pounds (25 kg), are similar to the most common side effects in adults. Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of budesonide delayed-release capsules. For more information, ask your healthcare provider or pharmacist.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**How should I store budesonide delayed-release capsules?**

- Store budesonide delayed-release capsules at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep budesonide delayed-release capsules in a tightly closed container.
- Budesonide delayed-release capsules come in child-resistant bottles.

**Keep budesonide delayed-release capsules and all medicines out of the reach of children.**

**General information about the safe and effective use of budesonide delayed-release capsules**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use budesonide delayed-release capsules for a condition for which it was not prescribed. Do not give budesonide delayed-release capsules to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about budesonide delayed-release capsules that is written for health professionals.

**Please address medical inquiries to, Northstar Rx LLC at Tel.: 1-800-206-7821.**

**What are the ingredients in budesonide delayed-release capsules?**

**Active ingredient:**budesonide, USP

**Inactive ingredients:**acetyltributyl citrate, ethylcellulose aqueous dispersion, gelatin, iron oxide red, iron oxide yellow, methacrylic acid copolymer dispersion, polysorbate 80, simethicone emulsion, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide and triethyl citrate. The capsule shell is printed with black pharmaceutical ink which contains following ingredients: iron oxide black, potassium hydroxide, propylene

glycol and shellac.

This Patient Information has been approved by the U.S. Food and Drug Administration.

**Repackaged and Distributed By:**

**Remedy Repack, Inc.**

**625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762**

Rev.: 01/25

DRUG: budesonide

GENERIC: budesonide

DOSAGE: CAPSULE, COATED PELLETS

ADMINISTRATION: ORAL

NDC: 70518-4613-0

NDC: 70518-4613-1

COLOR: orange

SHAPE: CAPSULE

SCORE: No score

SIZE: 19 mm

IMPRINT: 720

PACKAGING: 1 in 1 POUCH

OUTER PACKAGING: 50 in 1 BOX

ACTIVE INGREDIENT(S):

- BUDESONIDE 3mg in 1

INACTIVE INGREDIENT(S):

- TRIETHYL CITRATE
- TALC
- SUCROSE
- SODIUM LAURYL SULFATE
- SHELLAC
- PROPYLENE GLYCOL
- POTASSIUM HYDROXIDE
- TITANIUM DIOXIDE
- POLYSORBATE 80
- GELATIN
- FERROSO FERRIC OXIDE
- FERRIC OXIDE YELLOW
- FERRIC OXIDE RED
- ETHYLCELLULOSES
- DIMETHICONE
- METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER
- ACETYLTRIBUTYL CITRATE

# Budesonide DR Capsule

MFG NDC: 16714-0829-01  
MFG: Northstar Rx LLC, Memphis, TN  
38141

**3 mg**

**QTY: 50 Per Box**  
NDC #: **70518-4613-00**  
LOT #:  
Expires:  
Capsule ORANGE/WHITE 720

Usual Dosage: See Insert

Keep this and all medication out of the reach of children  
Store at 20-25°C (68-77°F); excursions permitted to 15-30°C (59-86°F) [See USP]



Repackaged By: RemedyRepack Inc.,  
Indiana, PA 15701, 724.465.8762

**RX ONLY**

# Budesonide DR Capsule

MFG NDC: 16714-0829-01  
MFG: Northstar Rx LLC, Memphis, TN  
38141

**3 mg**

**QTY: 1 Capsule**  
NDC #: **70518-4613-01**  
LOT #:  
Expires:  
Capsule ORANGE/WHITE 720

Usual Dosage: See Insert

Keep this and all medication out of the reach of children  
Store at 20-25°C (68-77°F); excursions permitted to 15-30°C (59-86°F) [See USP]



Repackaged By: RemedyRepack Inc.,  
Indiana, PA 15701, 724.465.8762

**RX ONLY**

## BUDESONIDE

budesonide capsule, coated pellets

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:70518-4613(NDC:16714-829)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BUDESONIDE</b> (UNII: Q3OKS62Q6X) (BUDESONIDE - UNII:Q3OKS62Q6X)	BUDESONIDE	3 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETYLTRIBUTYL CITRATE</b> (UNII: 0ZBX0N59RZ)	

<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>ETHYLCELLULOSES</b> (UNII: 7Z8S9VYZ4B)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)
<b>GELATIN</b> (UNII: 2G86QN327L)
<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)

### Product Characteristics

<b>Color</b>	orange (OPAQUE LIGHT-ORANGE) , white (OPAQUE WHITE)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	720
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70518-4613-0	50 in 1 BOX	04/23/2026	
1	NDC:70518-4613-1	1 in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206134	04/23/2026	

**Labeler** - REMEDYREPACK INC. (829572556)