

**MECLIZINE 25MG- meclizine tablet, chewable**  
**Innovida Pharmaceutique Corporation**

-----

***Active Ingredient (in each chewable tablet)***

Meclizine HCl 25 mg

***Purpose***

Expectorant

***Uses***

- prevents and treats nausea, vomiting or dizziness due to motion sickness

***Warnings***

**Do not use** in children under 12 years of age unless directed by a doctor

**Ask a doctor before use if you have**

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if** you are taking sedatives or tranquilizers

**When using this product**

- do not exceed recommended dosage
- may cause drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away ( **1-800-222-1222**).

***Directions***

Dosage should be taken one hour before travel starts

---

adults and children 12 years of age and over	chew 1 to 2 tablets once daily, or as directed by a doctor
children under 12 years of age	do not give this product to children under 12 years of age unless directed by a doctor

---

Other Information

- Store in a dry place at 15°-30°C (59°-86°F)
- keep lid tightly closed

Inactive Ingredients

Croscarmellose Sodium, Crospovidone, FD&C Red #40 Lake, French Vanilla Flavor, Lactose, Magnesium Stearate, Raspberry Flavor, Silica, Sodium Saccharin, Stearic Acid

Questions or Comments? 1-888-462-4166

Product label

NDC: 71800-052-03

INNVIDA

PHARMACEUTIQUE CORPORATION

Meclizine

25 mg

ANTIEMETIC

1000 Chewable Tablet

Drug Facts

Active ingredient  
(in each chewable tablet)

Meclizine HCl 25 mg.....Expectorant

Purpose

Expectorant

Uses

• prevents and treats nausea, vomiting or dizziness due to motion sickness

Warnings

Do not use in children under 12 years of age unless directed by a doctor

Ask a doctor before use if you have

• glaucoma • a breathing problem such as emphysema or chronic bronchitis • trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

• do not exceed recommended dosage • may cause drowsiness • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

• Dosage should be taken one hour before travel starts

adults and children 12 years of age and over	chew 1 to 2 tablets once daily, or as directed by a doctor
children under 12 years of age	do not give this product to children under 12 years of age unless directed by a doctor

Other Information

• Store in a dry place at 15°-30°C (59°-86°F) • keep lid tightly closed

Inactive ingredients

Croscarmellose Sodium, Crospovidone, FD&C Red #40 Lake, French Vanilla Flavor, Lactose, Magnesium Stearate, Raspberry Flavor, Silica, Sodium Saccharin, Stearic Acid

Questions or Comments? 1-888-462-4166

TEMPER EVIDENT. DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Mfd. for: Innovida Pharmaceutical Corporation, Phoenix, AZ 85040

MECLIZINE 25MG

meclizine tablet, chewable

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71800-052
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)		MECLIZINE HYDROCHLORIDE	25 mg
Inactive Ingredients			
Ingredient Name			Strength

<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>VANILLA</b> (UNII: Q74T35078H)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>RASPBERRY</b> (UNII: 4N14V5R27W)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	

### Product Characteristics

<b>Color</b>	white (Off White)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71800-052-03	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/10/2025	

**Labeler** - Innovida Pharmaceutique Corporation (080892908)

Revised: 2/2025

Innovida Pharmaceutique Corporation