

REGULAR STRENGTH ANTACID ANTIGAS- aluminum hydroxide, magnesium hydroxide, dimethicone liquid
VALU MERCHANDISERS COMPANY

ALS (BC) ANTACID MINT

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 200mg (equivalent to dried gel, USP)
Magnesium hydroxide 200 mg
Simethicone 20mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls two times a day or as directed by a doctor
- do not exceed 16 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks

- children under 12 years: ask a doctor

Other information

- **each 5 mL teaspoonful contains:** magnesium 85 mg, sodium 3 mg
- store at room temperature tightly closed
- do not freeze



Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package label


**REGULAR STRENGTH
ANTACID
ANTI-GAS**

ALUMINA, MAGNESIA,
AND SIMETHICONE
ORAL SUSPENSION

FAST-ACTING

FOR RELIEF OF
ACID INDIGESTION · HEARTBURN
· SOUR STOMACH · PRESSURE & BLOATING

**MINT CREME
FLAVOR**



12 FL OZ (355 mL)

276-11212-2 REV0323

Drug Facts TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

Active ingredients (in each 5 mL teaspoonful)	Purpose
Aluminum hydroxide (equivalent to dried gel, USP) 200 mg	Antacid
Magnesium hydroxide 200 mg	Antacid
Simethicone 20 mg	Antigas

Uses relieves • acid indigestion • heartburn • sour stomach
• the symptoms referred to as gas

Warnings
Ask a doctor before use if you have
• kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last for more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Directions • shake well before use
• adults and children 12 years of age and older: take 2 to 4 teaspoonfuls two times a day or as directed by a doctor
• do not exceed 16 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks
• children under 12 years of age: ask a doctor

Other information
• each 5 mL teaspoonful contains: magnesium 85 mg, sodium 3 mg
• store at room temperature tightly closed • do not freeze

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

PROUDLY DISTRIBUTED BY:
ASSOCIATED WHOLESALE
GROCERS, INC.
KANSAS CITY, KANSAS 66106

ALWAYS SAVE 66742 0723 GER

FOR PRODUCT INFORMATION
PLEASE VISIT:
ALWAYSSAVEBRAND.COM/66742

976-11212-2 REV0323



0 70038 66742 1

 www.alwaysavebrand.com

REGULAR STRENGTH ANTACID ANTIGAS

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-639
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	200 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	200 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	20 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-639-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	08/01/2017	

Labeler - VALU MERCHANDISERS COMPANY (868703513)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(63941-639) , pack(63941-639)

