

ASPIRIN- aspirin tablet, film coated
Better Living Brands, LLC

Signature Select 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- asthma (wheezing)
- hives
- facial swelling
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal Display Panel**COMPARE TO**

Genuine Bayer® Aspirin Active Ingredient†

NDC 21130-957-17

Signature

SELECT™

ASPIRIN

325 mg

Pain Reliever/
Fever Reducer

(NSAID)

Actual
Size

300 TABLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG,
distributors of Genuine Bayer® Aspirin. 50844 REV0122E15717

DISTRIBUTED BY: BETTER LIVING BRANDS LLC

P.O. BOX 99, PLEASANTON, CA 94566-0009

‡1-888-723-3929

**LOVE IT
OR IT'S ON US‡**

COMPARE TO
Genuine Bayer® Aspirin Active Ingredient!
NDC 21130-957-17

Signature SELECT™

ASPIRIN 325 mg

Pain reliever/
Fever reducer
(NSAID)

Actual Size

300 TABLETS

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin, 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves: ■ minor pain of arthritis
■ headache ■ muscle pain ■ menstrual pain
■ toothache ■ pain and fever of colds

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ asthma (wheezing) ■ hives ■ facial swelling ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed ■ have 3 or more alcoholic drinks every day while using this product

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

*This product is not manufactured or distributed by Bayer AG, distributors of Genuine Bayer® Aspirin, 508-44 REV0122E15717
DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0099
#1-888-723-3929

LOVE IT OR IT'S ON US!

S1714
RD 24137
3 21130 17602 4 5
PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn ■ you have asthma
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away, you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ ringing in the ears or a loss of hearing occurs
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ do not take more than directed
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
■ children under 12 years: do not use unless directed by a doctor

Other information
■ store at 25° C (77° F); excursions permitted between 15°-30° C (59°-86° F)
■ use by expiration date on package

Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 1-800-426-9391

STOP PEELING

Signature Select 44-157

ASPIRIN aspirin tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-957
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
Inactive Ingredients			
	Ingredient Name		Strength
	STARCH, CORN (UNII: O8232NY3SJ)		
	HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
	POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-957-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/20/2023	
2	NDC:21130-957-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
3	NDC:21130-957-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	08/31/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/10/1996	

Labeler - Better Living Brands, LLC (009137209)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(21130-957)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(21130-957)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(21130-957)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(21130-957)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(21130-957)