

LANSOPRAZOLE- lansoprazole capsule, delayed release
Rite Aid Corporation

DRUG FACTS

Active ingredient (in each capsule)

Lansoprazole USP, 15 mg

Purpose

Acid Reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole.

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drug

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20°-25°C (68°-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients

black iron oxide, colloidal silicon dioxide, corn starch, FD&C Blue #1, FD&C red #3, FD&C red #40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer type C, polyethylene glycol, polysorbate 80, potassium hydroxide, propylene glycol, shellac, strong ammonia solution, sucrose, sugar spheres (corn starch and sucrose), talc, titanium dioxide, yellow iron oxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

†Compare to the active ingredient in Prevacid® 24HR*

LANSOPRAZOLE

Delayed-Release Capsules USP, 15 mg

24HR

Acid Reducer

TREATS FREQUENT HEARTBURN

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

CAPSULES

†This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®, or by Perrigo Company plc, distributor of the Prevacid® 24HR product.

TAMPER EVIDENT: DO NOT USE IF TAMPER-EVIDENT INNER SEAL OR BLACK BAND AROUND THE CENTER OF EACH CAPSULES IS BROKEN OR MISSING.

KEEP OUTER CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

www.riteaid.com

Package Label

Drug Facts (continued)

- Keep product out of high heat and humidity
- Protect product from moisture
- Close cap tightly after use

Inactive ingredients
 Back iron oxide, colloidal silicon dioxide, corn starch, FD&C Blue No. 1, FD&C Red No. 3, FD&C Red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer type C, polyethylene glycol, polyacrylate 50, potassium hydroxide, propylene glycol, stearic, stearic ammonium solution, sucrose, sugar spheres (corn starch and sucrose), titanium dioxide, yellow iron oxide

Questions or comments?
 Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Teva Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid® or by Perrigo Company, distributor of the Prevacid® 24HR product.

WARNING: KEEP OUT OF REACH OF CHILDREN. NEVER USE ALONE. KEEP THIS AND ALL OTHER MEDICATIONS IN THEIR ORIGINAL CONTAINERS. KEEP THESE CONTAINERS TIGHTLY CLOSED.

NDC 11822-3374-5

Compare to the active ingredient in Prevacid® 24HR

LANSOPRAZOLE

Delayed-Release Capsules USP, 15 mg

24HR
 Acid Reducer

TREATS FREQUENT HEARTBURN
 MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

ONE 14-DAY COURSE OF TREATMENT

ACTUAL SIZE
14
 CAPSULES (1 BOTTLE INSIDE)

ONE 14-DAY COURSE OF TREATMENT

14
 CAPSULES (1 BOTTLE INSIDE)

Drug Facts

Active ingredient (in each capsule)
 Lansoprazole USP, 15 mg

Purpose
 Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Drug Facts (continued)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

Directions

- Swallow 1 capsule with a glass of water before eating in the morning
- Do not take more than 1 capsule a day
- Do not use for more than 14 days unless directed by your doctor
- Do not use for more than 14 days or more often than every 4 months unless directed by a doctor
- Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- Read the directions, warnings and package insert before use
- Keep the carton and package insert. They contain important information.
- Store at 20° to 25°C (68° to 77°F)

DISTRIBUTED BY:
 RITE AID, 30 HUNTER LANE,
 CAMD HILL, PA 17011
 www.riteaid.com

1 BOTTLE INSIDE

SATISFACTION GUARANTEE
 If you're not satisfied, we'll happily refund your money.

PLD: C565B
 F008157
 Lot No.:
 Exp. Date:

0 11822 33113 5

RITE AID Acid Reducer

LANSOPRAZOLE

lansoprazole capsule, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-3374
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LANSOPRAZOLE (UNII: 0K5C5T2QPG) (LANSOPRAZOLE - UNII:0K5C5T2QPG)	LANSOPRAZOLE	15 mg

Inactive Ingredients

Ingredient Name	Strength
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYDROXYPROPYL CELLULOSE, LOW SUBSTITUTED (UNII: 2165RE0K14)	
MAGNESIUM CARBONATE (UNII: 0E53J927NA)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SHELLAC (UNII: 46N107B710)	
AMMONIA (UNII: 5138Q19F1X)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

Product Characteristics

Color	pink, green	Score	no score
Shape	CAPSULE	Size	18mm
Flavor		Imprint Code	MYL;LD15
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3374-5	1 in 1 CARTON	11/29/2019	11/29/2026
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-3374-7	3 in 1 BOX	11/29/2019	11/29/2026
2		14 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA203187	11/29/2019	11/29/2026

Labeler - Rite Aid Corporation (014578892)

Revised: 8/2025

Rite Aid Corporation