

**ASPIRIN- aspirin tablet, film coated**  
**Topco Associates, LLC**

-----  
**TopCare 44-157**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves:

- pain and fever of colds
- toothache
- menstrual pain
- muscle pain
- headache
- minor pain of arthritis

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have asthma
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients**

corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?**

**1-888-423-0139**

**Principal display panel**

**TopCare®**

health

NDC 36800-157-14

COMPARE TO GENUINE BAYER® ASPIRIN  
ACTIVE INGREDIENT†

**REGULAR STRENGTH**

**Aspirin**

**325 mg**

PAIN RELIEVER • FEVER REDUCER

**(NSAID)**

- Safe Pain Relief
- Sodium &  
Caffeine Free

**500** COATED TABLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP  
IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG,  
owner of the registered trademark Genuine Bayer® Aspirin.

50844 REV0122C15714


DISTRIBUTED BY TOPCO ASSOCIATES LLC

ELK GROVE VILLAGE, IL 60007

©TOPCO LNKA0521 QUESTIONS? 1-888-423-0139

topcare@topco.com www.topcarebrand.com

**Visit here for more information: <http://topbrnds.com/48ZZSP>**



NDC 36800-157-14

COMPARE TO GENUINE BAYER® ASPIRIN ACTIVE INGREDIENT!

**REGULAR STRENGTH**


# Aspirin

**325 mg**

PAIN RELIEVER • FEVER REDUCER (NSAID)

- Safe Pain Relief
- Sodium & Caffeine Free

**500** COATED TABLETS



0 36800 09337 9

DISTRIBUTED BY TOPCO ASSOCIATES LLC  
 ELK GROVE VILLAGE, IL 60007  
 ©TOPCO LNK60521 QUESTIONS? 1-888-423-0139  
 topcare@topco.com www.topcareand.com  
 visit here for more information: http://topcareand.com/4238

No print/No varnish  
Lot & Exp date

**Drug Facts**  
**Active ingredient (in each tablet)**  
 Aspirin 325 mg (NSAID) ..... Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Purpose**  
 Pain reliever/fever reducer

**Uses** temporarily relieves: ■ pain and fever of colds ■ toothache ■ menstrual pain  
 ■ muscle pain ■ headache ■ minor pain of arthritis

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ shock ■ facial swelling ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ take a blood thinning (anticoagulant) or steroid drug ■ have had stomach ulcers or bleeding problems ■ take more or for a longer time than directed ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ are age 60 or older

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer  
**Ask a doctor before use** if ■ you have ever had an allergic reaction to this product or any of its ingredients ■ you have asthma ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have high blood pressure, heart disease, liver disease, or kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

**Other information** ■ use by expiration date on package ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients** corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?** 1-888-423-0139

**TopCare 44-157**

**ASPIRIN**

aspirin tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-157
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

**Packaging**

<b>Marketing Start</b>	<b>Marketing End</b>
------------------------	----------------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-157-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
2	NDC:36800-157-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
3	NDC:36800-157-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
4	NDC:36800-157-87	1 in 1 CARTON	02/10/1996	06/13/2026
4		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/10/1996	

**Labeler** - Topco Associates, LLC (006935977)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(36800-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(36800-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(36800-157)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(36800-157)