CLINDAMYCIN- clindamycin capsule Cronus Pharma LLC

Clindamycin Hydrochloride Capsules, USP

For Use In Dogs Only

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

Clindamycin Hydrochloride Capsules, USP contain clindamycin hydrochloride which is the hydrated salt of clindamycin. Clindamycin is a semisynthetic antibiotic produced by a 7(S)-chlorosubstitution of the 7(R)-hydroxyl group of a naturally produced antibiotic produced by *Streptomyces lincolnensis var. lincolnensis.*

Clindamycin Hydrochloride Capsules, USP (For Use in Dogs Only):

25 mg Capsule, each yellow and colorless capsule contains clindamycin hydrochloride equivalent to 25 mg of clindamycin.

75 mg Capsule, each green and colorless capsule contains clindamycin hydrochloride equivalent to 75 mg of clindamycin.

150 mg Capsule, each blue and colorless capsule contains clindamycin hydrochloride equivalent to 150 mg of clindamycin.

300 mg Capsule, each turquoise and colorless capsule contains clindamycin hydrochloride equivalent to 300 mg of clindamycin.

CLINICAL PHARMACOLOGY

Absorption

Clindamycin hydrochloride is rapidly absorbed from the canine gastrointestinal tract.


Dog Serum Levels

Serum levels at or above 0.5 μ g/mL can be maintained by oral dosing at a rate of 2.5 mg/lb of clindamycin hydrochloride every 12 hours. This same study revealed that average peak serum concentrations of clindamycin occur 1 hour and 15 minutes after oral dosing. The elimination half -life for clindamycin in dog serum was approximately 5 hours. There was no bioactivity accumulation after a regimen of multiple oral doses in healthy dogs.

Clindamycin Serum Concentrations

2.5 mg/lb (5.5 mg/kg) After B.I.D. Oral

Dose of Clindamycin Hydrochloride Capsules to Dogs



Metabolism and Excretion

Extensive studies of the metabolism and excretion of clindamycin hydrochloride administered orally in animals and humans have shown that unchanged drug and bioactive and bioinactive metabolites are excreted in urine and feces. Almost all of the bioactivity detected in serum after clindamycin hydrochloride product administration is due to the parent molecule (clindamycin). Urine bioactivity, however, reflects a mixture of clindamycin and active metabolites, especially N-demethyl clindamycin and clindamycin sulfoxide.

Site and Mode of Action

Clindamycin is an inhibitor of protein synthesis in the bacterial cell. The site of binding appears to be in the 50S sub-unit of the ribosome. Binding occurs to the soluble RNA fraction of certain ribosomes, thereby inhibiting the binding of amino acids to those ribosomes. Clindamycin differs from cell wall inhibitors in that it causes irreversible modification of the protein-synthesizing subcellular elements at the ribosomal level.

Microbiology

Clindamycin is a lincosaminide antimicrobial agent with activity against a wide variety of aerobic and anaerobic bacterial pathogens. Clindamycin is a bacteriostatic compound that inhibits bacterial protein synthesis by binding to the 50S ribosomal sub-unit. The minimum inhibitory concentrations (MICs) of Gram-positive and obligate anaerobic pathogens isolated from dogs in the United States are presented in Table 1. Bacteria were isolated in 1998-1999. All MICs were performed in accordance with the Clinical and Laboratory Standards Institute (CLSI).

Table 1. Clindamycin MIC Values (µg/mL) from Diagnostic Laboratory Survey Data Evaluating Canine Pathogens in the U.S. during 1998-991

Organism Soft Tissue/Wound2	Number of Isolates	MIC50	MIC85	MIC90	Range
Staphylococcus aureus	17	0.5	0.5	≥ 4.0	0.25- ≥ 4.0
Staphylococcus intermedius	28	0.25	0.5	≥ 4.0	0.125- ≥ 4.0
<i>Staphylococcus</i> spp	18	0.5	0.5	≥ 4.0	0.25- ≥ 4.0
Beta-hemolytic streptococci	46	0.5	0.5	≥ 4.0	0.25- ≥ 4.0
Streptococcus spp	11	0.5	≥ 4.0	≥ 4.0	0.25-≥ 4.0
Osteomyelitis/Bone3					
Staphylococcus aureus	20	0.5	0.5	0.5	0.54
Staphylococcus intermedius	15	0.5	≥ 4.0	≥ 4.0	0.25-≥ 4.0
<i>Staphylococcus</i> spp	18	0.5	≥ 4.0	≥ 4.0	0.25-≥ 4.0
Beta-hemolytic streptococci	21	0.5	2.0	2.0	0.25-≥ 4.0
Streptococcus spp	21	≥ 4.0	≥ 4.0	≥ 4.0	0.25-≥ 4.0
Dermal/Skin5					
Staphylococcus aureus	25	0.5	≥ 4.0	≥ 4.0	0.25-≥ 4.0
Staphylococcus intermedius	48	0.5	≥ 4.0	≥ 4.0	0.125- ≥ 4.0
<i>Staphylococcus</i> spp	32	0.5	≥ 4.0	≥ 4.0	0.25-≥ 4.0
Beta-hemolytic streptococci	17	0.5	0.5	0.5	0.25-0.5

1 The correlation between the in vitro susceptibility data and clinical response has not been determined.

2 Soft Tissue/Wound: includes samples labeled wound, abscess, aspirate, exudates, draining tract, lesion, and mass.

- 3 Osteomyelitis/Bone: includes samples labeled bone, fracture, joint, tendon
- 4 No range, all isolates yielded the same value
- 5 Dermal/Skin: includes labeled skin, skin swab, biopsy, incision, lip

INDICATIONS

Clindamycin Hydrochloride Capsules, USP (for use in dogs only) are indicated for the treatment of infections caused by susceptible strains of the designated microorganisms in the specific conditions listed below:

Dogs: Skin infections (wounds and abscesses) due to coagulase positive staphylococci (Staphylococcus aureus or Staphylococcus intermedius). **Deep wounds and abscesses** due to Bacteroides fragilis, Prevotella melaninogenicus, Fusobacterium necrophorum and Clostridium perfringens.

Dental infections due to *Staphylococcus aureus*, *Bacteroides fragilis*, *Prevotella melaninogenicus*, *Fusobacterium necrophorum and Clostridium perfringens*.**Osteomyelitis** due to *Staphylococcus aureus*, *Bacteroides fragilis*, *Prevotella melaninogenicus*, *Fusobacterium necrophorum and Clostridium perfringens*.

CONTRAINDICATIONS

Clindamycin Hydrochloride Capsules, USP are contraindicated in animals with a history of

hypersensitivity to preparations containing clindamycin or lincomycin.

Because of potential adverse gastrointestinal effects, do not administer to rabbits, hamsters, guinea pigs, horses, chinchillas or ruminating animals.

WARNINGS

Keep out of reach of children. Not for human use.

PRECAUTIONS

During prolonged therapy of one month or greater, periodic liver and kidney function tests and blood counts should be performed.

The use of Clindamycin Hydrochloride Capsules, USP occasionally results in overgrowth of non-susceptible organisms such as clostridia and yeasts. Therefore, the administration of Clindamycin Hydrochloride Capsules should be avoided in those species sensitive to the gastrointestinal effects of clindamycin (see **CONTRAINDICATIONS**). Should superinfections occur, appropriate measures should be taken as indicated by the clinical situation.

Patients with very severe renal disease and/or very severe hepatic disease accompanied by severe metabolic aberrations should be dosed with caution, and serum clindamycin levels monitored during high-dose therapy.

Clindamycin hydrochloride has been shown to have neuromuscular blocking properties that may enhance the action of other neuromuscular blocking agents. Therefore, Clindamycin Hydrochloride Capsules, USP should be used with caution in animals receiving such agents.

Safety in gestating bitches and queens or breeding male dogs has not been established.

ADVERSE REACTIONS

Side effects occasionally observed in either clinical trials or during clinical use were vomiting and diarrhea.

To report a suspected adverse reaction call 1-844-227-6687 (1-844-2-CRONUS)

DOSAGE AND ADMINISTRATION

Dogs: Infected Wounds, Abscesses, and Dental Infections

Oral: 2.5-15.0 mg/lb body weight every 12 hours.

Duration: Treatment with Clindamycin Hydrochloride Capsules, USP may be continued up to a maximum of 28 days if clinical judgment indicates. Treatment of acute infections should not be continued for more than three or four days if no response to therapy is seen.

Dosage Schedule

Capsules

Clindamycin Hydrochloride Capsules, USP 25 mg, administer 1-6 capsules every 12 hours for each 10 pounds of body weight.

Clindamycin Hydrochloride Capsules, USP 75 mg, administer 1 -6 capsules every 12 hours for each 30 pounds of body weight.

Clindamycin Hydrochloride Capsules, USP 150 mg, administer 1-6 capsules every 12 hours for each 60 pounds of body weight.

Clindamycin Hydrochloride Capsules, USP 300 mg, administer 1-6 capsules every 12 hours for each 120 pounds of body weight.

Dogs: Osteomyelitis

Oral: 5.0-15.0 mg/lb body weight every 12 hours.

Duration: Treatment with Clindamycin Hydrochloride Capsules is recommended for a minimum of 28 days. Treatment should not be continued for longer than 28 days if no response to therapy is seen.

Dosage Schedule:

Capsules

Clindamycin Hydrochloride Capsules, USP 25 mg, administer 2-6 capsules every 12 hours for each 10 pounds of body weight.

Clindamycin Hydrochloride Capsules, USP 75 mg, administer 2 -6 capsules every 12 hours for each 30 pounds of body weight.

Clindamycin Hydrochloride Capsules, USP 150 mg, administer 2-6 capsules every 12 hours for each 60 pounds of body weight.

Clindamycin Hydrochloride Capsules, USP 300 mg, administer 2-6 capsules every 12 hours for each 120 pounds of body weight

ANIMAL SAFETY SUMMARY

Rat and Dog Data: One year oral toxicity studies in rats and dogs at doses of 30, 100 and 300 mg/kg/day (13.6, 45.5 and 136.4 mg/lb/day) have shown clindamycin hydrochloride capsules to be well tolerated. Differences did not occur in the parameters evaluated to assess toxicity when comparing groups of treated animals with contemporary controls. Rats administered clindamycin hydrochloride at 600 mg/kg/day (272.7 mg/lb/day) for six months tolerated the drug well; however, dogs orally dosed at 600 mg/kg/day (272.7 mg/lb/day) vomited, had anorexia, and subsequently lost weight. At necropsy these dogs had erosive gastritis and focal areas of necrosis of the mucosa of the gallbladder.

Safety in gestating bitches or breeding males has not been established.

STORAGE

Store at controlled room temperature 68° to 77°F (20° to 25°C).

How Supplied

Clindamycin Hydrochloride Capsules are available as: 25 mg - bottles of 200 75 mg - bottles of 200 150 mg - bottles of 100 and 500 300 mg - bottles of 100

To report suspected adverse drug events, for technical assistance or to obtain a copy of the safety data sheet (SDS), contact Cronus Pharma LLC at 1-844-227-6687 or 1-844-2-CRONUS.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at https://www.fda.gov/animal-veterinary/safety-health.

Approved by FDA under ANADA # 200-298

Manufactured for: **Cronus Pharma LLC**, East Brunswick, NJ 08816. Contact No: 1-844-227-6687 (1-844-2-CRONUS)

Made in India

Distributed by:

Cronus Pharma LLC 2 Tower Center Boulevard, Suite 1101A, East Brunswick, NJ 08816 Contact No: 1-844-227-6687 (1-844-2-CRONUS) e-Fax No: 732-647-1272 Email: contact@cronuspharma.com Issued: 02/2023

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 69043-013-20

Clindamycin Hydrochloride Capsules, USP

25 mg

Equivalent to 25 mg clindamycin

Approved by FDA under ANADA # 200-298

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

200 Capsules

For Use in Dogs Only



PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 69043-014-20

Clindamycin Hydrochloride Capsules, USP

75 mg

Equivalent to 75 mg clindamycin

Approved by FDA under ANADA # 200-298

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

200 Capsules

For Use in Dogs Only



PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 69043-015-01

Clindamycin Hydrochloride Capsules, USP

150 mg

Equivalent to 150 mg clindamycin

Approved by FDA under ANADA # 200-298

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

100 Capsules

For Use in Dogs Only



PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 69043-015-05

Clindamycin Hydrochloride Capsules, USP

150 mg

Equivalent to 150 mg clindamycin

Approved by FDA under ANADA # 200-298

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

500 Capsules

For Use in Dogs Only

NDC 69043-015-05	Approved for use in dogs. Recommended dosage: For therapy of wounds, abscesses and dental infections: 1-6 capsules for each 60 pounds body weight every 12 hours.	Manufactured for: Cronus Pharma LLC, East Brunswick, NJ 08816. Contact No: 1-844-227-6687 (1-844-2-CRONUS)
Clindamycin Hydrochloride Capsules, USP 150 mg	For therapy of osteomyelitis, 2-6 capsules for each 60 pounds of body weight every 12 hours. See package insert for complete product	Made in India Code No.: TN/Drugs/TN00002327
500 Capsules	information.	LC01505-02 Issued: 08/2019
Equivalent to 150 mg clindamycin Approved by FDA under ANADA # 200-298	Warning : Keep out of reach of children. Not for human use.	1550E0. 00/2019
CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.	Keep container tightly closed. Store at controlled room temperature 68° to 77°F (20° to 25°C).	
For Use in Dogs Only	. ,	
Cronus	Each capsule contains: Clindamycin hydrochloride equivalent to clindamycin, 150 mg.	

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 69043-016-01

Clindamycin Hydrochloride Capsules, USP

300 mg

Equivalent to 300 mg clindamycin

Approved by FDA under ANADA # 200-298

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

100 Capsules

For Use in Dogs Only

NDC 69043-016-01 Clindamycin Hydrochloride Capsules, USP 300 mg 100 Capsules Equivalent to 300 mg clindamycin Approved by FDA under ANADA # 200-298 CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian. For Use in Dogs Only	 Approved for use in dogs. Recommended dosage: For therapy of wounds, abscesses and dental infections: 1-6 capsules for each 120 pounds body weight every 12 hours. For therapy of osteomyelitis, 2-6 capsules for each 120 pounds of body weight every 12 hours. For therapy of osteomyelitis, 2-6 capsules for each 120 pounds of body weight every 12 hours. See package insert for complete product information. Warning: Keep out of reach of children. Not for human use. Keep container tightly closed. Store at controlled room temperature 68° to 77°F (20° to 25°C). Each capsule contains: Clindamycin hydrochloride equivalent to clindamycin, 300 mg. 	Code No.: TN/Drugs/TN00002327 LC01601-02 Issued: 08/2019 Manufactured for: Cronus Pharma LLC, East Brunswick, NJ 08816. Contact No: 1-844-227-6687 (1-844-2-CRONUS) Made in India	N 3 6 9 0 4 3 0 1 6 0 1 1
--	--	---	---------------------------

CLINDAMYCIN

clindamycin capsule

Product Infor	mation						
Product Type		PRESCRIPTION ANIMA	AL DRUG	ltem Code	(Source)	NDC:6	9043-013
Route of Admini	istration	ORAL					
Active Ingredi	ient/Active	Moiety					
	Ing	redient Name			Basis Streng		Strength
CLINDAMYCIN HYI UNII:3U02EL437C)	DROCHLORIDE	(UNII: T200Q1YN1W)	(CLINDAMYCII	N -	CLINDAMYCIN	J	25 mg
Product Chara	acteristics						
Color	YELLOW (YELLO	W-WHITE)	Score		no sco	re	
Shape	CAPSULE		Size		16mm		
Flavor			Imprin	nt Code	CRONU	S;CLIN2	5
Contains							
Packaging							
# Item Code	e Packa	ge Description	Marketin	ig Start Dat	e Marke	eting E	nd Date
1 NDC:69043-013-2	20 200 in 1 B	OTTLE					
Marketing	Informat	ion					

Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date
ANADA	ANADA200298	10/04/2019	

CLINDAMY								
Product Info	ormat	tion						
Product Type			PRESCRIPTION ANIM	AL DRUG	ltem Code (Source) NDC:	69043-014
Route of Admi	nistra	ition	ORAL					
Active Ingree	dient	Active	Moiety					
		Ing	redient Name				sis of ength	Strengt
CLINDAMYCIN H UNII:3U02EL437C)	YDROC	CHLORIDE	(UNII: T200Q1YN1W)	(CLINDAMYCIN	-	CLINDAM	IYCIN	75 mg
Product Cha	racte	ristics						
Color		EEN (GREEN	N-WHITE)	Score		no	score	
Shape		SULE	·····-,	Size		16r		
Flavor				Imprint	Code	CRO	DNUS;CLIN7	5
Contains								
Packaging								
# Item Coc	-		ge Description	Marketin	g Start Dat	e Ma	arketing I	End Date
1 NDC:69043-014	1-20	200 in 1 B	OTTLE					
Marketing	Inf	ormat	ion					
Marketing Category		Applicat	tion Number or M Citation	onograph	Marketin Dat			eting End Date
ANADA		NADA20029	0.0		10/04/2019			

clindamycin capsule			
Product Information			
Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:69043-015
Route of Administration	ORAL		

•	ant/Active	Moiety					
	Ing	redient Name			Basis Stren		Strength
CLINDAMYCIN HYD UNII:3U02EL437C)	ROCHLORIDE	(UNII: T200Q1YN1W)	(CLINDAMYCIN	l - (CLINDAMYC	•	150 mg
Product Chara	cteristics						
Color	BLUE (BLUE-V	WHITE)	Score		no score		
Shape	CAPSULE		Size		19mm		
Flavor			Imprint Co	de	CRONUS;	CLIN150	
Contains							
Packaging							
# Item Code	Packa	ge Description	Marketin	g Start Date	Mark	ceting E	nd Date
1 NDC:69043-015-0		BOTTLE					
2 NDC:69043-015-0	5 500 in 1 E	BOTTLE					
Marketing I	nformat	ion					
Marketing Category		tion Number or M Citation	onograph	Marketing Date			ting End ate
ANADA	ANADA2002	98		10/04/2019			
	le						
clindamycin capsı	le	PRESCRIPTION ANIM	AL DRUG	ltem Code (S	ource)	NDC:6	9043-016
clindamycin capsu Product Inforr	ule nation	PRESCRIPTION ANIMA ORAL	AL DRUG	ltem Code (S	iource)	NDC:6	9043-016
clindamycin capsu Product Inforr Product Type	ule nation stration	ORAL	AL DRUG	ltem Code (S	ource)	NDC:6	9043-016
clindamycin capsu Product Inforr Product Type Route of Adminis	ule nation stration ent/Active	ORAL	AL DRUG	ltem Code (S	iource) Basis Stren	s of	
Product Type Route of Adminis Active Ingredie	ule nation stration ent/Active Ing	ORAL Moiety			Basis	s of igth	
Clindamycin capsu Product Inforr Product Type Route of Adminis Active Ingredie CLINDAMYCIN HYD UNII: 3U02EL437C)	ule nation stration ent/Active Ing ROCHLORIDE	ORAL Moiety redient Name			Basis Stren	s of igth	Strength
Clindamycin capsu Product Inforr Product Type Route of Adminis Active Ingredia CLINDAMYCIN HYD UNII: 3U02EL437C) Product Chara	ule nation stration ent/Active Ing ROCHLORIDE	ORAL Moiety redient Name	(CLINDAMYCIN	I - (Basis Stren	s of igth IN	Strength
Clindamycin capsu Product Inforr Product Type Route of Adminis Active Ingredic CLINDAMYCIN HYD UNII:3U02EL437C) Product Chara Color TU	ule nation stration ent/Active Ing ROCHLORIDE cteristics RQUOISE (TUF	ORAL Moiety redient Name	(CLINDAMYCIN	l- ore	Basis Stren CLINDAMYC no s	s of ogth IN core	Strength
Clindamycin capsu Product Inforr Product Type Route of Adminis Active Ingredie CLINDAMYCIN HYD UNII:3U02EL437C) Product Chara Color TU Shape CA	ule nation stration ent/Active Ing ROCHLORIDE	ORAL Moiety redient Name	(CLINDAMYCIN Sc Siz	l- ore ze	Basis Stren CLINDAMYC	s of igth IN core	Strength 300 mg
Clindamycin capsu Product Inforr Product Type Route of Adminis Active Ingredic CLINDAMYCIN HYD UNII:3U02EL437C) Product Chara Color TU	ule nation stration ent/Active Ing ROCHLORIDE cteristics RQUOISE (TUF	ORAL Moiety redient Name	(CLINDAMYCIN Sc Siz	l- ore	Basis Stren CLINDAMYC	s of ogth IN core	Strength 300 mg

Packaging					
# Item Code	Package Description	Marketing	Start Date	Marke	eting End Date
1 NDC:69043-016-01	100 in 1 BOTTLE				
Marketing II	nformation				
Marketing II Marketing Category	nformation Application Number or Mo Citation	onograph	Marketing S Date	tart	Marketing End Date

Labeler - Cronus Pharma LLC (079421067)

Revised: 11/2023

Cronus Pharma LLC