

**DAYTIME AND NIGHTTIME COLD/FLU RELIEF SOFTGELS - COMBO PACK-
acetaminophen, dextromethorphan hbr, phenylephrine hcl, doxylamine
succinate
ARMY AND AIR FORCE EXCHANGE SERVICE**

**exchange select DayTime and Nighttime Cold/Flu Relief Softgels - Combo
Pack**

DayTime COLD/FLU RELIEF

Drug Facts

**Active ingredients
(in each Softgel)**

Acetaminophen325 mg

Dextromethorphan HBr10 mg

Phenylephrine HCl5 mg

Purpose

AcetaminophenPain reliever/Fever reducer

Dextromethorphan HBrCough suppressant

Phenylephrine HCl Nasal Decongestant

Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever

Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash,

nausea or vomiting, consult a doctor promptly.

Do not use:

■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease
- thyroid disease ■ diabetes
- high blood pressure
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for

adults as well as children, even if you do not notice any signs or symptoms.

Directions

- take only as directed (see Warnings)
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 4 hours
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

■ even if taking DayTime during day and Nighttime during night, total limit per day is 4 doses per 24-hour period.

Other information

- Store at room temperature 20-25° C (68-77° F)
- avoid excessive heat

Inactive ingredients FD&C Red # 40, FD&C Yellow # 6, Gelatin, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide

Questions? Call Weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Nighttime COLD/FLU RELIEF

Drug Facts

**Active ingredients
(in each Softgel)**

- Acetaminophen325 mg
- Dextromethorphan HBr15 mg
- Doxylamine succinate6.25 mg

Purpose

- AcetaminophenPain reliever/fever reducer
- Dextromethorphan HBrCough suppressant
- Doxylamine succinate Antihistamine

Uses temporarily relieves common cold/flu symptoms: ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches and pain ■ fever ■ runny nose & sneezing

Warnings Liver warning: This product contains acetaminophen. Severe liver

damage may occur if you take ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include ■ skin reddening ■ blisters ■ rash. If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use: ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ to make a child sleep.

Ask a doctor before use if you have

■ liver disease ■ glaucoma

■ cough that occurs with too much phlegm (mucus)

■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

■ trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

■ taking sedatives or tranquilizers

■ taking the blood thinning drug warfarin

When using this product

■ do not use more than directed

■ excitability may occur, especially in children

■ marked drowsiness may occur

- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control center right away. Quick medical attention is critical for adults as well as children, even if you do not notice any signs or symptoms.

Directions

- take only as directed (see Warnings)
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hours
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

- even if taking DayTime during day and Nighttime during night, total limit per day is 4 doses per 24-hour period.

Other information

- Store at room temperature 20-25° C (68-77° F)
- avoid excessive heat

Inactive ingredients

D&C Yellow # 10, FD&C Blue # 1, Gelatin, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide.

Questions? Call Weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Principal Display Package Label: exchange select DayTime and Nighttime Cold/Flu Relief Softgels - Combo Pack

DayTime Cold/Flu Relief Nighttime Cold/Flu Relief

32 Softgels / 16 Softgels

**Compare to the active ingredients in ingredients in

Vicks® DayQuil® Cold and Flu LiquiCaps® and Flu Liquicaps®

**This product is not manufactured or manufactured or distributed by Procter & Gamble, owner of Gamble, owner of the registered trademark Vicks® DayQuil® trademark Vicks® NyQuil® Cold & Flu LiquiCaps®.

**Compare to the active

Vicks® NyQuil® Cold

**This product is not distributed by Procter & the registered Cold & Flu LiquiCaps®.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

SATISFACTION GUARANTEED OR YOUR MONEY BACK.

Manufactured For Your Military Exchanges
By AptaPharma Inc.,
1533 Union Avenue, Pennsauken, NJ 08110
1-877-798-5944 aptapharma.com

Drug Facts (continued) DayTime	Drug Facts (continued) Nighttime
Directions ■ take only as directed (see Warnings) ■ do not exceed 4 doses per 24 hrs ■ adults & children 12 yrs & over 2 softgels with water every 4 hours ■ children 4 to under 12 yrs ask a doctor ■ children under 4 yrs do not use ■ even if taking Daytime during day and Nighttime during night, total limit per day is 4 doses per 24-hour period	Directions ■ take only as directed (see Warnings) ■ do not exceed 4 doses per 24 hrs ■ adults & children 12 yrs & over 2 softgels with water every 6 hours ■ children 4 to under 12 yrs ask a doctor ■ children under 4 yrs do not use ■ even if taking Daytime during day and Nighttime during night, total limit per day is 4 doses per 24-hour period
Other information ■ store at room temperature 20-25°C (68-77°F) ■ avoid excessive heat	Other information ■ store at room temperature 20-25°C (68-77°F) ■ avoid excessive heat
Inactive ingredients FD&C Red # 40, FD&C Yellow # 6, Sorbitol, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide	Inactive ingredients D&C Yellow # 10, FD&C Blue # 1, Sorbitol, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide
Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944	Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

**This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® DayQuil® Cold & Flu LiquiCaps®.

SATISFACTION GUARANTEED OR YOUR MONEY BACK.
Manufactured For Your Military Exchanges
By AptaPharma Inc.,
1533 Union Ave., Pennsauken, NJ 08110
1-877-798-5944 aptapharma.com

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



Drug Facts DayTime	Drug Facts Nighttime
<p>Active ingredients (in each softgel) Purpose</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Phenylephrine HCl 5 mg Nasal decongestant</p> <p>Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product</p> <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping</p>	<p>Active ingredients (in each softgel) Purpose</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Doxylamine Succinate 6.25 mg Antihistamine</p> <p>Uses temporarily relieves common cold/flu symptoms: ■ sore sore ■ sore throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever ■ runny nose & sneezing</p> <p>Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product</p> <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping</p>

Drug Facts (continued) DayTime	Drug Facts (continued) Nighttime
<p>Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma or emphysema</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.</p> <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ symptoms get worse or last more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children, even if you do not notice any signs or symptoms.</p>	<p>Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema ■ trouble urinating due to enlarged prostate gland</p> <p>Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin</p> <p>When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives & tranquilizers may increase drowsiness</p> <p>Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 5 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children, even if you do not notice any signs or symptoms.</p>

Drug Facts (continued) DayTime	Drug Facts (continued) Nighttime
<p>Directions ■ take only as directed (see Warnings) ■ do not exceed 4 doses per 24 hrs</p> <p>adults & children 12 yrs & over 2 softgels with water every 4 hours children 4 to under 12 yrs ask a doctor children under 4 yrs do not use</p> <p>■ even if taking DayTime during day and Nighttime during night, total limit per day is 4 doses per 24-hour period</p> <p>Other information ■ store at room temperature 20-25°C (68-77°F) ■ avoid excessive heat</p> <p>Inactive ingredients FD&C Red # 40, FD&C Yellow # 6, Gelatin, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide</p> <p>Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-786-5844</p>	<p>Directions ■ take only as directed (see Warnings) ■ do not exceed 4 doses per 24 hrs</p> <p>adults & children 12 yrs & over 2 softgels with water every 6 hours children 4 to under 12 yrs ask a doctor children under 4 yrs do not use</p> <p>■ even if taking DayTime during day and Nighttime during night, total limit per day is 4 doses per 24-hour period</p> <p>Other information ■ store at room temperature 20-25°C (68-77°F) ■ avoid excessive heat</p> <p>Inactive ingredients D&C Yellow # 10, FD&C Blue # 1, Gelatin, Glycerin, Polyethylene glycol, Povidone, Propylene glycol, Purified water, Sorbitol sorbitan solution, Titanium dioxide</p> <p>Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-786-5844</p>

SATISFACTION GUARANTEED OR YOUR MONEY BACK.
Manufactured For Your Military Exchanges
By: AptaPharma Inc.,
1533 Union Ave., Pennsauken, NJ 08110
1-877-786-5844 aptapharma.com

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLENDED SMELL IS NOTICED. BREAK IN OR SEE ANY SIGNS OF UNUSUALNESS.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

exchange select

DayTime

COLD/FLU RELIEF

Acetaminophen
Dextromethorphan HBr/ Phenylephrine HCl

• Headache, Fever, Sore Throat, Minor Aches & Pains
• Nasal Congestion, Sinus Pressure
• Cough

Actual Size

32 Softgels

exchange select

Nighttime

COLD/FLU RELIEF

Acetaminophen
Dextromethorphan HBr/ Doxylamine Succinate

• Aches, Fever & Sore Throat
• Sneezing, Runny Nose
• Cough

Actual Size

16 Softgels

exchange select

Nighttime

COLD/FLU RELIEF

Acetaminophen
Dextromethorphan HBr/ Phenylephrine HCl

Actual Size

Quality Value

BK-152
P000000
5 142299 034133 6

res

DAYTIME AND NIGHTTIME COLD/FLU RELIEF SOFTGELS - COMBO PACK

acetaminophen, dextromethorphan hbr, phenylephrine hcl, doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55301-605
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-605-05	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	01/01/2025	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	4 BLISTER PACK	32
Part 2	2 BLISTER PACK	16

Part 1 of 2

DAYTIME COLD/FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Item Code (Source)	NDC:55301-604
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O921CV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	604
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2025	

Part 2 of 2

NIGHTTIME COLD/FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:55301-603
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	603
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2025	

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Registrant - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(55301-605, 55301-604, 55301-603)

Revised: 1/2025

ARMY AND AIR FORCE EXCHANGE SERVICE