

CUTAR- coal tar lotion
Summers Laboratories Inc

SUMMERS LABS (as PLD) - CUTAR (11086-105)

ACTIVE INGREDIENT

7.5% L.C.D. (EQUIVALENT TO 1.5% COAL TAR)

PURPOSE

PSORIASIS, SEBORREIC DERMATITIS TREATMENT

USE

- CONTROLS ITCHING, FLAKING AND REDNESS DUE TO PSORIASIS AND SEBORRHEIC DERMATITIS

WARNINGS

FOR EXTERNAL USE ONLY

ASK A DOCTOR BEFORE USE IF YOU HAVE

- CONDITION THAT COVERS A LARGE AREA OF THE BODY
- CONDITION IN OR AROUND THE RECTUM, GENITAL AREA OR GROIN

WHEN USING THIS PRODUCT

- AVOID CONTACT WITH EYES. IF CONTACT OCCURS, RINSE EYES THOROUGHLY WITH WATER
- USE CAUTION IN EXPOSING SKIN TO SUNLIGHT AFTER APPLYING THIS PRODUCT. IT MAY INCREASE YOUR TENDENCY TO SUNBURN FOR UP TO 24 HOURS AFTER APPLICATION

DO NOT USE

- FOR PROLONGED PERIODS WITHOUT CONSULTING A DOCTOR
- WITH OTHER FORMS OF PSORIASIS THERAPY UNLESS DIRECTED TO DO SO BY A DOCTOR

STOP USE AND ASK A DOCTOR IF

- CONDITION WORSENS OR DOES NOT IMPROVE AFTER REGULAR USE AS DIRECTED
- EXCESSIVE IRRITATION OCCURS

KEEP OUT OF REACH OF CHILDREN

- IF SWALLOWED, GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY
- TO PREVENT STAINING DO NOT USE IN PLASTIC OR FIBERGLASS TUBS, SINKS OR BASINS. CUTAR MAY STAIN LIGHT COLORED FABRICS

DIRECTIONS

- APPLY TO AFFECTED AREAS ONE TO FOUR TIMES DAILY OR AS DIRECTED BY A DOCTOR
- AS A MEDICATED LOTION: GENTLY RUB A SMALL AMOUNT ONTO AFFECTED AREAS.

BLOT OFF EXCESS WITH A TISSUE TO HELP PREVENT STAINING OF CLOTHING OR LINENS

- IN THE BATH: POUR A LIBERAL QUANTITY (2 TO 4 OZ.) INTO THE TUB, MAKING SURE CUTAR IS WELL MIXED. SOAK FOR 10 TO 20 MINUTES, OR AS DIRECTED BY YOUR DOCTOR. AFTER TREATMENT, PAT SKIN DRY.
- GUARD AGAINST SLIPPING.
- AS A HAND OR FOOT SOAK: ADD 1 TO 2 OZ. TO A SINK OR BASIN. MIX WELL AND SOAK FOR 5 TO 10 MINUTES OR AS DIRECTED BY YOUR DOCTOR.

INACTIVE INGREDIENTS

WATER, MINERAL OIL, ISOPROPYL MYRISTATE, POLYSORBATE 80, SORBITAN SESQUIOLEATE, LANOLIN ALCOHOLS EXTRACT, METHYL AND PROPYL PARABENS, FRAGRANCE, XANTHAN GUM, CARBOMER

<p>Drug Facts</p> <table border="1"> <tr> <td>Active Ingredient 7.5% L.C.D. (equiv. to 1.5% coal tar)</td> <td>Purpose Psoriasis, seborrheic dermatitis treatment</td> </tr> </table>	Active Ingredient 7.5% L.C.D. (equiv. to 1.5% coal tar)	Purpose Psoriasis, seborrheic dermatitis treatment	<h1>cūtar[®]</h1> <h2>emulsion</h2>
Active Ingredient 7.5% L.C.D. (equiv. to 1.5% coal tar)	Purpose Psoriasis, seborrheic dermatitis treatment		
<p>Use Controls itching, flaking and redness due to psoriasis and seborrheic dermatitis.</p>			
<p>Warnings For external use only</p>	<p>✓ Recommended by dermatologists</p>		
<p>Ask a doctor before use if you have a</p> <ul style="list-style-type: none"> ■ condition that covers a large area of the body ■ condition in or around the rectum or genital area or groin 	<p>✓ Controls itching, redness and scaling associated with psoriasis and seborrheic dermatitis</p>		
<p>When using this product ■ avoid contact with eyes. If contact occurs rinse eyes thoroughly with water. ■ use caution exposing skin to sunlight after applying this product. It may increase your tendency to sunburn for up to 24 hours after application.</p>	<p>✓ One product - two uses</p> <ol style="list-style-type: none"> 1) apply as a medicated lotion 2) mix with water for soothing therapeutic soaks 		
<p>Do not use ■ for prolonged periods without consulting a doctor. ■ with other psoriasis therapies unless directed by a doctor.</p>	<p>SATISFACTION GUARANTEED</p>		
<p>Stop use and ask a doctor if ■ condition worsens or does not improve after regular use. ■ excessive irritation occurs.</p>			
<p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>SUMMERS LABORATORIES INC</p>		
<p>Directions To prevent staining, do not use in plastic or fiberglass tubs, sinks or basins. May stain light colored fabrics. ■ Medicated lotion: apply to affected areas one to four times daily or as directed by a doctor. Gently rub a small amount onto affected areas. Blot off excess with a tissue. ■ Bath: pour a liberal quantity (2-4 oz.) into the tub, mixing well. Soak for 10-20 minutes, or as directed by your doctor. After treatment pat skin dry. ■ Hand/foot soak: add 1-2 oz. to a sink or basin. Mix well and soak for 5-10 minutes or as directed by a doctor.</p>			
<p>Inactive ingredients water, mineral oil, isopropyl myristate, polysorbate 80, sorbitan sesquioleate, lanolin alcohol, methyl and propyl parabens, fragrance, xanthan gum, carbomer</p>	<p>Distributed by: Summers Laboratories, Inc. Collegeville, PA 19426 800-533-SKIN (7546) www.summlab.com</p>		
<p>NDC 11086-105-01</p>	<p>SHAKE WELL BEFORE USING</p>		
<p>WARNING: This product contains a chemical known to the state of California to cause cancer.</p>	<p>6 oz. (177mL)</p>		
<p>L-0005-01-04</p>			

CUTAR
coal tar lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11086-105
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
COAL TAR (UNII: R533ESO2EC) (COAL TAR - UNII:R533ESO2EC)	COAL TAR	1.5 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
MINERAL OIL (UNII: T5L8T28FGP)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SORBITAN SESQUIOLEATE (UNII: 0W8RRI5W5A)	
LANOLIN ALCOHOLS (UNII: 884C3FA9HE)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
XANTHAN GUM (UNII: TTV12P4NEE)	
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11086-105-01	180 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/29/2013	
2	NDC:11086-105-02	3785 mL in 1 JUG; Type 0: Not a Combination Product	10/29/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M032	10/29/2013	

Labeler - Summers Laboratories Inc (002382612)

Revised: 10/2023

Summers Laboratories Inc