EXCHANGE SELECT ESOMEPRAZOLE MAGNESIUM- esomeprazole capsule, delayed release

Army & Air Force Exchange Service

Perrigo Company Esomeprazole Magnesium Drug Facts

Active ingredient (in each capsule)

Esomeprazole 20 mg

(Each delayed-release capsule corresponds to 22 mg esomeprazole magnesium dihydrate)

Purpose

Acid reducer

Uses

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

• taking a prescription drug. Acid reducers may interact with certain prescription

drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients

FD&C blue no. 1, FD&C red no. 3, ferric oxide, gelatin, glyceryl monostearate, hypromellose, magnesium stearate, meglumine, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, shellac, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Exchange Select[™] Compare to Nexium[®] 24 HR active ingredient See New Warning Information Esomeprazole Magnesium Delayed-Release Capsules, 20 mg/Acid Reducer 24 HOUR TREATS FREQUENT HEARTBURN CAPSULES Actual size 42 CAPSULES Three 14-day courses of treatment May take 1 to 4 days for full effect guality value




ССІИІСАLLY РКОУЕМ ТО ССІМІСАLLY РКОУЕМ ТО

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

HOUR

Drug Facts Active ingredient (in each capsule) Esomeprazole 20 mgAcid reducer (Each delayed-release capsule corresponds to 22 mg esomeprazole magnesium dihydrate)	Drug Facts (continued) If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	ļ	
 Uses treats frequent heartburn (occurs <u>2 or more</u> days a week) not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect 	Directions adults 18 years of age and older this product is to be used once a day (every 24 hours), every day for 14 days may take 1 to 4 days for full effect		
Warnings Allergy alert: Do not use if you are allergic to esomeprazole Do not use if you have: Trouble or pain swallowing food, vomiting with blood, or bloody or black stools heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have	 14-Day Course of Treatment swallow 1 capsule with a glass of water before eating in the morning take every day for 14 days do not take more than 1 capsule a day swallow whole. Do not crush or chew capsules. do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) you may repeat a 14-day course every 4 months do not take for more than 14 days or more often than every 4 months unless directed by a doctor children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. 		
 had heartburn over 3 months. This may be a sign of a more serious condition. frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain 	Other information read the directions and warnings before use keep the carton. It contains important information. store at 20-25°C (68-77°F)		
Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days	Inactive ingredients FD&C blue no. 1, FD&C red no. 3, ferric oxide, gelatin, glyceryl monostearate, hypromellose, magnesium stearate, meglumine, methacrylic acid and ethyl acrylate copolymer dispersion, polyethylene glycol, polysorbate 80, shellac, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide, triethyl citrate		
 you need to take more than 1 course of treatment every 4 months you get diarrhea you develop a rash or joint pain 	Questions or comments? 1-800-719-9260		



EXCHANGE SELECT ESOMEPRAZOLE MAGNESIUM

Product Infor	mation						
				6			01.000
Product Type		HUMAN OTC DRUG	Item	Code (So	urce)	NDC:553	01-898
Route of Admini	stration	ORAL					
Active Ingredi	ent/Active	Moiety					
Ingredient Name Basis of Str						trength	Strengt
ESOMEPRAZOLE (U	JNII: N3PA6559	FT) (ESOMEPRAZOLE - UNII	:N3PA6	559FT)	ESOMEPRAZ O	DLE	20 mg
Inactive Ingre	dients						
		Ingredient Name				S	trength
FD&C BLUE NO. 1	(UNII: H3R47K3	TBD)					
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)							
FERRIC OXIDE RED							
GELATIN, UNSPEC							
GLYCERYL MONOS							
		(UNII: 3NXW29V3WO)					
MAGNESIUM STEA MEGLUMINE (UNII:		J97M0I30)					
		CIFIED (UNII: 3WJQ0SDW1	Δ)				
POLYSORBATE 80			, ,				
SHELLAC (UNII: 46N	-						
SODIUM LAURYL S	ULFATE (UNII:	368GB5141J)					
TALC (UNII: 7SEV7J4	IR1U)						
TITANIUM DIOXIDE	UNII: 15FIX9V	2JP)					
TRIETHYL CITRATE	(UNII: 8Z96Q)	(D6UM)					
Product Chara	cteristics						
Color	BLUE (opa	aque)	Sco	re		no sco	ore
Shape	CAPSULE	(oblong)	Siz	e		14mm	
Flavor			Imp	rint Code		L898	
Contains							
Packaging				Markat	ing Start	Marka	ting End
# Item Code	Pac	kage Description			ing Start ate		ting End ate
1 NDC:55301-898- 01	1 in 1 CARTON	I		12/04/2017			
1	14 in 1 BOTTL Product	E; Type 0: Not a Combinat	ion				
2 NDC:55301-898- 03	3 in 1 CARTON			12/04/2017			
2	14 in 1 BOTTL Product	E; Type 0: Not a Combinat	ion				

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
ANDA	ANDA207193	12/04/2017					

Labeler - Army & Air Force Exchange Service (001695568)

Revised: 11/2022

Army & Air Force Exchange Service