

**HUMCO CALACLEAR- pramoxine hydrochloride and zinc acetate lotion**  
**Humco Holding Group, Inc.**

-----  
**Humco Calaclear**

**Humco Calamine Lotion**

Drug Facts

**Active Ingredient**

Pramoxine HCl 1%

**Purpose**

External Analgesic

**Active Ingredient**

Zinc Acetate 0.1%

**Purpose**

Skin Protectant

**Use**

Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.

**Warnings**

**For external use only.** Use only as directed.

**Avoid contact** with eyes and mucous membranes.

**Ask a doctor before using**

on children under 2 years of age.

**When using this product. Discontinue use if**

condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days. and consult a doctor.

**Keep out of reach of children.**

In case of accidental ingestion, seek professional assistance or contact a Poison Control

Center immediately.

## Directions

- Adults and children 2 yrs. of age and older. Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.
- Children under 2 years of age: Consult a doctor before use.

## Other Information

- store at room temperature 15 ° - 30 °C (50 ° - 85 °F)

## Inactive Ingredient

SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.

## Label

Drug Facts		Drug Facts (CONTINUED)
<b>Active Ingredients</b> Pramoxine HCl 1%.....External analgesic Zinc Acetate 0.1%.....Skin protectant	<b>Purpose</b> External analgesic Skin protectant	<b>Directions</b> ■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort. ■ Children under 2 years of age: Consult a doctor before use.
<b>Uses</b> Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.		<b>Other information</b> Store at room temperature 15-30°C (50-86°F)
<b>Warnings</b> ■ For external use only. Use only as directed. ■ Avoid contact with eyes and mucous membranes. ■ Ask a doctor before using on children under 2 years of age.		<b>Inactive ingredients</b> SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.
<b>When using this product</b> discontinue use if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.		HUMCO, Texarkana, TX 75501 Questions or Comments? ☎ 1-800-662-3435 Go to: <a href="http://www.simplehomemedies.com">www.simplehomemedies.com</a> to find helpful hints and learn more about Calaclear™ Lotion.
<b>Keep out of reach of children.</b> In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.		<b>MADE IN USA</b>
<b>Continued</b> →		

*\*Compare to Caladryl® Clear™ Lotion*

NDC 0395-0400-86

**CALACLEAR®**  
**LOTION**  
LOCIÓN CALACLEAR®  
The Healthy Skin Care Solution

- External analgesic
- Skin protectant
- Drying agent & itch relief

6 FL OZ (177 mL)  
6 ONZAS FL (177 mL)

HUMCO

03950 40096 9

A45240  
02 2024

Levante etiqueta para información



**Drug Facts**

**Active ingredients Purpose**  
 Pramoxine HCl 1% and Zinc Acetate 0.1%  
 .....External analgesic/  
 Skin protectant

**Uses** Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.

**Warnings**  
 ■ **For external use only.** Use only as directed.  
 ■ **Avoid contact with eyes and mucous membranes.**  
 ■ **Ask a doctor before using** on children under 2 years of age.

**When using this product**  
**Discontinue use if** condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.

**Keep out of reach of children.** In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

*Continued* ▶

This product is not manufactured or distributed by Pfizer, owner of the registered trademark "Caladryl® Clear™ Lotion"

\*Compare to  
Caladryl® Clear™ Lotion

NDC 0395-0400-96

# CALACLEAR™ LOTION

*The Healthy  
Skin Care Solution*

- External analgesic
- Skin protectant
- Drying agent & itch relief

**6 FL OZ  
(177 mL)**

**HUMCO™**

**Drug Facts (CONTINUED)**

**Directions**  
 ■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.  
 ■ Children under 2 years of age: Consult a doctor before use.

**Other information**  
 Store at room temperature 15-30°C (50-86°F)

**Inactive ingredients**  
 SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.

R121211.RLG

HUMCO, Texarkana, TX 75501  
 Questions or Comments? ☎1-800-662-3435  
 \* Go to: [www.simplehomeremedies.com](http://www.simplehomeremedies.com)  
 to find helpful hints and learn more about Calaclear™ Lotion.

## HUMCO CALACLEAR

pramoxine hydrochloride and zinc acetate lotion

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0395-0400
<b>Route of Administration</b>	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>PRAMOXINE HYDROCHLORIDE</b> (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 mL
<b>ZINC ACETATE</b> (UNII: FM5526K07A) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	1 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>CAMPHOR (NATURAL)</b> (UNII: N20HL7Q941)	
<b>DIAZOLIDINYL UREA</b> (UNII: H5RIZ3MPW4)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	

<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KOOR)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0395-0400-96	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/25/1998	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M016	03/25/1998	

**Labeler** - Humco Holding Group, Inc. (825672884)

**Registrant** - Pharma Nobis, LLC (118564114)

### Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(0395-0400) , analysis(0395-0400) , pack(0395-0400) , label(0395-0400)

Revised: 12/2023

Humco Holding Group, Inc.