

# SUCROMATE EQUINE- deslorelin acetate and deslorelin injection

## Dechra Veterinary Products

-----

### SucroMate® Equine (deslorelin acetate)

#### Sterile Suspension

**Injectable sustained release gonadotropin releasing hormone (GnRH) analog**

**For use in horses only**

**For intramuscular injection**

**Refrigerate after each use**

#### CAUTION:


Federal law restricts this drug to use by or on the order of a licensed veterinarian.

#### DESCRIPTION:

SucroMate Equine (deslorelin acetate) is a sterile, synthetic gonadotropin-releasing hormone (GnRH) analog suspension. SucroMate Equine is a sustained release formulation that forms an in situ gel upon intramuscular injection. Deslorelin acetate is [(6-D-tryptophan-9-(N-ethyl-L-prolinamide)-10-deglycinamide)] GnRH. The molecular weight is 1282.6.

#### SucroMate Equine

The deslorelin acetate structural formula is:



[(6-D-tryptophan-9-(N-ethyl-L-prolinamide)-10-deglycinamide)] GnRH

Each mL contains 1.8 mg deslorelin acetate (1.7 mg deslorelin) for injection as a sterile suspension in sucrose acetate isobutyrate/propylene carbonate (70:30 wt:wt). Ten doses per vial.

## **INDICATION:**

SucroMate Equine is indicated for inducing ovulation within 48 hours of treatment in cyclic estrous mares with an ovarian follicle between 30 and 40 mm in diameter.

## **DOSAGE and ADMINISTRATION:**

**Shake well before use.** Administer 1 mL (one dose) per estrus cycle, 48 hours prior to desired ovulation. Verify that the mare is in estrus and has at least one ovarian follicle between 30 and 40 mm in diameter.

SucroMate Equine is a suspension of deslorelin and settling will occur over time. Warm the product for 2 minutes by rolling the vial between the palms (to reach room temperature), or allow the vial to sit for 30 minutes at room temperature prior to administration. Shake the vial vigorously for 1 minute before use. Administer a 1 mL intramuscular injection into the thick musculature of the neck.

Prior to administration, it should be determined by rectal palpation and/or ultrasonography that the cyclic estrous mare has an ovarian follicle between 30 and 40 mm in diameter. Effectiveness is contingent upon accurate diagnosis of estrus and detection of a developing follicle between 30 and 40 mm in diameter. Only 1 mL should be administered per mare during a given estrus.

## **CONTRAINDICATIONS:**

SucroMate Equine is contraindicated in horses known to be hypersensitive to deslorelin acetate.

## **WARNING:**

For use in horses (estrous mares) only. Do not use in horses intended for human consumption.

For intramuscular (IM) use only. Do not administer intravascularly. Not for use in humans. Keep this and all drugs out of reach of children.

## **HUMAN WARNING - NOT FOR HUMAN USE**

Pregnant women and women of childbearing age should exercise caution when handling this product. Accidental administration may lead to a disruption of the menstrual cycle. Direct contact with the skin should be avoided. If exposure occurs, contact areas should be washed immediately with alcohol followed by soap and water, as this product is insoluble in water. In case of accidental human injection, consult a physician immediately.

To obtain a copy of the Safety Data Sheet (SDS), contact Dechra at (866) 933-2472.

## **PRECAUTIONS:**

The use of GnRH analogs in cycling mares has been associated with prolonged interovulatory intervals. SucroMate Equine has not been evaluated in mares less than 3 years of age.

### **ADVERSE REACTIONS:**

Injection site swelling was observed following the administration of SucroMate Equine during the effectiveness and safety studies; all injection site swellings resolved within 5 days, and 7 - 14 days, respectively.

### **CONTACT INFORMATION:**

To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Dechra at (866) 933-2472.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS, or online at <http://www.fda.gov/reportanimalae>

### **CLINICAL PHARMACOLOGY:**

Deslorelin induces ovulation by increasing the levels of endogenous luteinizing hormone. SucroMate Equine has been shown to induce ovulation within 48 hours of treatment in 77% of mares with a developing follicle between 30 and 40 mm in diameter. This product is intended to optimize breeding management through induction of ovulation within 48 hours.

### **EFFECTIVENESS:**

Two hundred eighty three (283) mares, ranging in age from 2 to 20 years (mean 10.6 years) of age randomly received either SucroMate Equine or placebo after the presence of an ovarian follicle between 30 and 40 mm in diameter was confirmed. All 283 mares were evaluated for field safety under actual conditions of use. A total of 191 mares (97 placebo and 94 mares in the SucroMate Equine group) were evaluated for effectiveness over a total of 273 cycles (131 placebo and 142 SucroMate Equine cycles). Breeds included Standardbreds, Thoroughbreds, Quarter Horses and Paints. Mares were evaluated for up to 3 cycles to determine if SucroMate Equine induced ovulation in cycling mares, and to determine the effects on the interovulatory interval, pregnancy rate and general health characteristics of the resultant foals.

The proportion of mares ovulating within 48 hours of treatment was significantly higher in the SucroMate Equine group as compared to the placebo group (see Table 1).

**Table 1. Percentage of Mares Ovulating Within 48 Hours of Treatment (Yes = ovulated / No = did not ovulate)**

	<b>Placebo (1.0 mL)</b>	<b>SucroMate Equine (1.0 mL)</b>	<b>P value</b>
<b>1<sup>st</sup> Cycle</b>			
Number of mares			

Number of mares treated	97	94	
Yes	23 (24%)	68 (72%)	p = 0.000*
No	73 (75%)	25 (27%)	
Not determined	1 (1%)	1 (1%)	
<b>2<sup>nd</sup> Cycle</b>			
Number of mares treated	25	35	
Yes	9 (36%)	29 (83%)	p = 0.000*
No	16 (64%)	6 (17%)	
<b>3<sup>rd</sup> Cycle</b>			
Number of mares treated	9	13	
Yes	3 (33%)	12 (92%)	p = 0.006*
No	6 (67%)	0 (0%)	
Not determined	0 (0%)	1 (8%)	
<b>Percent of mares ovulating within 48 hours after treatment in any cycle</b>	<b>27%</b>	<b>77%</b>	

\* statistically significant

Following treatment in each of three estrus cycles, the mean hours to ovulation was reduced in the SucroMate Equine treated group compared to the placebo group (See Table 2).

**Table 2. Mean Hours to Ovulation\***

	<b>Placebo (1.0 mL)</b>	<b>SucroMate Equine (1.0 mL)</b>	<b>P value</b>
<b>1<sup>st</sup> Cycle</b>	91.8 (n=96)	65.1 (n=90)	p = 0.019†
<b>2<sup>nd</sup> Cycle</b>	83.0 (n=24)	56.2 (n=35)	p = 0.099
<b>3<sup>rd</sup> Cycle</b>	85.3 (n=9)	41.0 (n=12)	p = 0.153

\* data could not be obtained on all mares

† statistically significant

The interovulatory interval was calculated between cycles for all mares that did not become pregnant in the previous cycle. There was no significant difference in the length of interovulatory intervals between the SucroMate Equine (deslorelin acetate) treated group and the placebo-treated mares as represented in Table 3.

**Table 3. Mean Interovulatory Interval (days)\***

		<b>SucroMate</b>	
--	--	------------------	--

	<b>Placebo (1.0 mL)</b>	<b>SucroMate Equine (1.0 mL)</b>	<b>P value</b>
<b>Cycles 1-2</b>	26.1 (n=27)	22.3 (n=32)	p = 0.182
<b>Cycles 2-3</b>	21.3 (n=9)	18.5 (n=11)	p = 0.336
<b>Cycles 3-4</b>	13.0 (n=1)	18.0 (n=1)	

\* data could not be obtained on all mares

Twins were detected in 8 placebo mares and 2 SucroMate Equine mares. The nursing behavior, mobility and general health of the live foals were similar for foals from mares in the SucroMate Equine and placebo groups. See Table 4 for a summary of the foaling and foal assessment data.

**Table 4. Foaling and Foal Assessment Data\***

<b>Assessment</b>	<b>Treatment Group</b>	
	<b>Placebo</b>	<b>SucroMate Equine</b>
Mares confirmed pregnant in 1999 breeding season out of total # of mares per group % (n/N)	83% (118/143)	81% (114/140)
Mares confirmed to have a live foal out of # of pregnant mares % (n/N)	94% (111/118)	84% (96/114)
Mares confirmed to have a stillbirth out of # of pregnant mares % (n/N)	0% (0/118)	3.0% (3/114)
72 Hour-General Health Foal Assessment % (n/N)		
Good	80% (89/111)	79% (76/96)
Satisfactory	16% (18/111)	20% (19/96)
Not Satisfactory	2% (2/111)	2% (2/96)
4 Week-General Health Foal Assessment % (n/N)		
Good	34% (38/111)	36% (35/96)
Not Assessed	66% (73/111)	67% (64/96)

\* data could not be obtained on all mares

When administered to cycling, estrous mares with a pre-ovulatory follicle between 30 mm and 40 mm in diameter, a single, 1 mL injection of SucroMate Equine (1.8 mg of deslorelin acetate) resulted in ovulation within 48 hours of treatment in 77% of treated mares. The length of the interovulatory interval between cycles was not altered.

The pregnancy rate of the treated mares was similar to the placebo mares.

**Animal Safety:** In a target animal safety study, 32 reproductively sound, mixed breed mares, ranging in age from 3-15 years were randomly assigned to one of four dose groups; each dose group contained 8 mares. The dose groups were administered 0X (placebo), 1X (1 mL), 3X (3 mL) or 5X (5 mL) the recommended 1.8 mg dose of

SucroMate Equine for three consecutive cycles.

Treatment with SucroMate Equine at all doses reduced the mean hours to ovulation during all three treatment cycles compared to the placebo group. Of the sixty-five treatment experiences for the SucroMate Equine treated mares across the three cycles 55 (84.6%) were followed by ovulation within 48 hours compared to five of twenty-three (21.7%) for placebo mares. There was no statistically significant effect on the length of estrus at any dose level.

Some mares in all SucroMate Equine treatment groups exhibited mild to moderate swelling at the injection site six hours post-injection during all three treatment cycles. All injection site reactions in the 1X treated mares resolved by Day 3 post-treatment. Two mares, one 3X mare and one 5X mare, had swelling and hardness noted at injection sites at Day 7 post-treatment. The incidence of injection site reactions for the placebo, 1X, 3X and 5X groups was 37%, 75%, 50%, and 87%, respectively.

The LH concentrations in the SucroMate Equine treated groups in all three cycles exhibited an expected surge following treatment and prior to ovulation, followed by lower concentrations during the midluteal phase, with a gradual return to pretreatment levels.

The mean interovulatory interval in the SucroMate Equine groups ranged from 17.8 - 23 days compared to 21 - 23.5 days in the placebo group.

All mares that ovulated after the 3rd treatment with SucroMate Equine or placebo were bred.

These mares were bred until they became pregnant (not to exceed three cycles). In the placebo group, seven mares ovulated after the 3rd treatment and were bred. Six of the seven mares became pregnant (85.7%). In the 1X group, six mares ovulated after the 3rd treatment and were bred. All six became pregnant (100%). In the 3X group, eight mares ovulated after the 3rd treatment and were bred. Six of the eight mares became pregnant (75%). In the 5X group, five mares ovulated after the 3rd treatment and all became pregnant (100%).

There were no significant differences across the treatment groups in terms of time to pregnancy or the number and proportion of mares that became pregnant after three consecutive SucroMate Equine injections.

In a separate dose tolerance study mares received a single IM injection of placebo (8 mares) or SucroMate Equine at 10X the recommended 1.8 mg dose (8 mares). One SucroMate Equine 10X mare exhibited moderate tremors and hives at 6 hours post-treatment.

#### **STORAGE:**

Store at or below 8°C.

#### **HOW SUPPLIED:**

One (1) 10 mL vial (10 doses) per vial.

Approved by FDA under NADA # 141-319

Manufactured for:  
Dechra Veterinary Products  
7015 College Boulevard, Suite 525  
Overland Park, KS 66211 USA

SucroMate is a registered trademark of Dechra Limited

Rev. August 2022  
SP-1000595

**Dechra**

**PRINCIPAL DISPLAY PANEL - 10 mL Vial Carton**

SucroMate®  
Equine

(deslorelin acetate)

Sterile Suspension

For use in horses only

For Intramuscular Injection

CAUTION: Federal law restricts  
this drug to use by or on the  
order of a licensed veterinarian.

A sustained release gonadotropin  
releasing hormone (GnRH) analog  
(deslorelin acetate) for ovulation  
induction in estrous mares

Approved by FDA under  
NADA # 141-319

Net Contents:  
10 mL

Dechra

**SucroMate® Equine**  
 (deslorelin acetate)  
 Sterile Suspension  
 For use in horses only  
 For intramuscular injection  
 A sustained release gonadotropin  
 releasing hormone (GnRH) analog  
 (deslorelin acetate) for ovulation  
 induction in estrous mares  
 Net Contents: 10 mL 10 doses

**INDICATION:**  
 SucroMate Equine is indicated for inducing ovulation within 48 hours of treatment in cyclic estrous mares with an ovarian follicle between 30 and 40 mm in diameter.

**DOSAGE and ADMINISTRATION:**  
**Shake well before use.**  
 Administer 1 mL (one dose) per estrus cycle, 48 hours prior to desired ovulation. Verify that the mare is in estrus and has at least one ovarian follicle between 30 and 40 mm in diameter.

SucroMate Equine is a suspension of deslorelin and settling will occur over time. Warm the product for 2 minutes by rolling the vial between the palms (to reach room temperature), or allow the vial to sit for 30 minutes at room temperature prior to administration. Shake the vial vigorously for 1 minute before use. Administer a 1 mL intramuscular injection into the thick musculature of the neck.



**WARNING:**  
 For use in horses (estrous mares) only. Do not use in horses intended for human consumption. For IM use only. Do not administer intravascularly. **Not for use in humans. Keep this and all drugs out of reach of children.**

**HUMAN WARNINGS - NOT FOR HUMAN USE:**  
**Pregnant women and women of childbearing age should exercise caution when handling this product.** Accidental administration may lead to a disruption of the menstrual cycle. Direct contact with the skin should be avoided. If exposure occurs, contact areas should be washed immediately with alcohol followed by soap and water, as this product is insoluble in water. In case of accidental human injection, consult a physician immediately. **See package insert for complete product information.**

**SucroMate®  
 Equine**  
 (deslorelin acetate)  
 Sterile Suspension

For use in horses only  
 For Intramuscular Injection

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

A sustained release gonadotropin releasing hormone (GnRH) analog (deslorelin acetate) for ovulation induction in estrous mares

Approved by FDA under NADA # 141-319



Net Contents:  
 10 mL



**STORAGE:**  
 Store at or below 8°C.  
**NET CONTENTS:**  
 10 mL vial; each mL contains 1.8 mg deslorelin acetate (1.7 mg deslorelin) as a sterile suspension in sucrose acetate isobutyrate/propylene carbonate (70:30 wt:wt)  
**MANUFACTURED FOR:**  
 Dechra Veterinary Products  
 Overland Park, KS 66211 USA  
 Rev. March 2022



17033-906-10 1  
 N 3

LOT:  
 EXP:

SP-1000594

**SUCROMATE EQUINE**

deslorelin acetate and deslorelin injection

**Product Information**



<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17033-906	
<b>Route of Administration</b>	INTRAMUSCULAR			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	DESLORELIN ACETATE (UNII: 679007NR5C) (DESLORELIN - UNII:TKG3I66TVE)	DESLORELIN ACETATE	1.8 mg in 1 mL	
	DESLORELIN (UNII: TKG3I66TVE) (DESLORELIN - UNII:TKG3I66TVE)	DESLORELIN	1.7 mg in 1 mL	
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:17033-906-10	1 in 1 CARTON		
1		10 mL in 1 VIAL, MULTI-DOSE		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
NADA	NADA141319	11/05/2010		

**Labeler** - Dechra Veterinary Products (362142734)

Revised: 12/2023

Dechra Veterinary Products