

**ASPIRIN 50 CT- aspirin tablet**  
**JHK Inc dba American Safety & First Aid**

-----  
**Aspirin 50 CT**

***Drug Facts***

**Active ingredient (In Each Tablets)**

Aspirin 325 mg

**Purpose**

pain reliever/fever reducer

**Use**

temporarily relieves fever and minor aches and pains associated with:

- headache
- colds
- muscular aches
- toothache
- minor arthritis pain
- menstrual cramps

**Warnings**

**Reye's Syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Allergy Warning**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other prescription or nonprescription drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- If you are allergic to aspirin
- with any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn

**Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**When using this product** do not exceed recommended dose.

**Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain or fever persists or gets worse
- new symptoms occur
- redness or swelling is present

**If pregnant or breast-feeding,** ask a health professional before use. IT IS ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS DEFINITELY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

Adults and children 12 years of age and older: Take 2 tablets with water every 4 hours as needed, not to exceed 12 tablets in 24 hours, or as directed by doctor  
Children under 12 years of age: Ask a doctor.

## **Other Information**

- do not use if packet is opened or torn
- Store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)

## **Inactive Ingredients**

microcrystalline cellulose, povidone, pregelatinized starch

## **Questions?**

1-866-651-3660 Monday-Friday 8:00am-5:00pm EST

## **PRINCIPAL DISPLAY PANEL - 50 Tablet Packet Box**

Product #1112

Aspirin

Pain Reliever/Fever Reducer (NSAID)

- Temporary relief from pain and fever
- Soothes muscle aches and minor arthritis pain

These quality comfort tablets are formulated and packaged for use in the workplace and meet ANSI Z308.1-2015 standards. They are sealed in tamper-evident packets for safety and convenience.

50 Tablets • 2 Tablets Per Packet

Manufactured for American Safety & First Aid  
Osceola, IN 46561

Product #1112



# Aspirin

Pain Reliever/Fever Reducer (NSAID)

- Temporary relief from pain and fever
- Soothes muscle aches and minor arthritis pain

These quality comfort tablets are formulated and packaged for use in the workplace and meet ANSI Z308.1-2015 standards. They are sealed in tamper-evident packets for safety and convenience.

**50** Tablets • 2 Tablets Per Packet

Manufactured for American Safety & First Aid  
Osceola, IN 46561

#### Other Information

■ do not use if packet is opened or torn ■ Store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)

**Inactive Ingredients** microcrystalline cellulose, povidone, pregelatinized starch

**Questions?** 1-866-851-3660 Mon.- Fri. 8:00am-5:00pm EST

V1\_10/11/19



#### Drug Facts (continued)

**When using this product** do not exceed recommended dose.

#### Stop use and ask a doctor

If you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ pain or fever persists or gets worse ■ new symptoms occur ■ redness or swelling is present

#### If pregnant or breast-feeding,

ask a health professional before use. IT IS ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS DEFINITELY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** Adults and children 12 years of age and older: Take 2 tablets with water every 4 hours as needed, not to exceed 12 tablets in 24 hours, or as directed by doctor. Children under 12 years of age: Ask a doctor.

Product #1112

50 Tablets • 2 Tablets Per Packet

### Drug Facts

**Active Ingredient** (in Each Tablet).....*Purpose*  
Aspirin 325 mg.....pain reliever/fever reducer

**Use** temporarily relieves fever and minor aches and pains associated with: ■ headache ■ colds ■ muscular aches ■ toothache ■ minor arthritis pain ■ menstrual cramps

**Warnings**  
**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Allergy Warning:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other prescription or nonprescription drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you are allergic to aspirin ■ with any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have a history of stomach problems, such as heartburn

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for: ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis



Pain Reliever/Fever Reducer (NSAID)

PULL TO OPEN

Do not use if individual packet is open or torn.

# ASPIRIN 50 CT

aspirin tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73598-1112
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	ASPIRIN;44;157
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:73598-1112-1	25 in 1 BOX	02/02/2000	
1		2 in 1 DOSE PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph drug	M013	02/02/2000	

**Labeler** - JHK Inc dba American Safety & First Aid (867236309)