

**MAXIMUM STRENGTH ANTACID AND GAS RELIEF- aluminum hydroxide,
magnesium hydroxide, dimethicone suspension
RITE AID PHARMACY**

RITEAID max antacid

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)

Magnesium hydroxide 400 mg

Simethicone 40mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Do not take more than 12 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use

- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each 5 mL teaspoonful contains:** magnesium 165 mg, sodium 5 mg
- protect from freezing
- store at room temperature
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package Label

Compare to the active ingredients in
Maximum Strength Mylanta®*

NDC 11822-3727-6

MAXIMUM STRENGTH ANTACID & GAS RELIEF

ALUMINA, MAGNESIA,
AND SIMETHICONE ORAL
SUSPENSION USP

FAST ACTING

Soothing relief of:
heartburn • acid indigestion • sour
stomach • pressure & bloating



218-06122-6
REV GC-1222

ORIGINAL
FLAVOR
ALCOHOL 0.5%
12 FL OZ (355 mL)

Drug Facts

TAMPER-EVIDENT: Do not use if breakaway band on bottle cap is missing or broken.

Active ingredients

(in each 5 mL teaspoonful)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

Purposes

Uses relieves • heartburn • sour stomach • acid indigestion
• the symptoms referred to as gas

Warnings

Do not take more than 12 teaspoonsful in 24 hours or use the maximum dosage for more than 2 weeks

Ask a doctor before use if you have

• kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Directions

• shake well before use
• adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
• children under 12 years: ask a doctor

Other information

• each 5 mL teaspoonful contains:
magnesium 165 mg, sodium 5 mg • store at room temperature
• protect from freezing • keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*RITE AID is not affiliated with the owner of the registered trademark MYLANTA®. 918-06122-6 REV1222

DISTRIBUTED BY: RITE AID,
200 NEWBERRY COMMONS
ETTERS, PA 17319

www.riteaid.com

MADE IN USA

SATISFACTION GUARANTEE:

If you're not satisfied,
we'll happily refund your money.



0 11822 37276 3

MAXIMUM STRENGTH ANTACID AND GAS RELIEF

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-3727
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	400 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3727-6	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	04/01/2014	

Labeler - RITE AID PHARMACY (014578892)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(11822-3727) , pack(11822-3727)

Revised: 11/2023

RITE AID PHARMACY