

**GOOD SENSE EFFERVESCENT COLD RELIEF- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent
Tower Laboratories Ltd.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Good Sense Effervescent Cold Relief

Active ingredients

(in each effervescent tablet)

Aspirin 325 mg **(NSAID)***

Chlorpheniramine maleate 2 mg

Phenylephrine bitartrate 7.8 mg

*Nonsteroidal anti-inflammatory drug

Purpose

Aspirin 325 mg **(NSAID)***.....Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg.....Antihistamine

Phenylephrine bitartrate 7.8 mg.....Nasal decongestant

*Nonsteroidal anti-inflammatory drug

Uses

temporarily relieves these symptoms of the common cold:

■ fever ■ sneezing ■ nasal congestion ■ sore throat ■ headache ■ minor aches and pains ■ runny nose ■ sinus congestion and pressure

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood

thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic. **You have** ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ an allergic reaction occurs. Seek medical help right away. ■ pain or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present ■ ringing in the ears or a loss of hearing occurs ■ nervousness, dizziness or sleeplessness occurs

When using this product

■ do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours
- do not take more than 8 tablets in 24 hours
- *children under 12 years: ask a doctor*

Other information

each tablet contains: sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Serious side effects associated with the use of this product can be reported to the following address:

Tower Laboratories LTD.

P.O. Box 306

8 Industrial Park Rd.
Centerbrook, CT 06409

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have diabetes
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a medicine for asthma or chronic bronchitis
- you have a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs or symptoms:
- stomach bleeding
- fever
- vomiting
- black stools
- stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- nervousness, dizziness or sleeplessness occurs
- ringing in the ears or a loss of hearing occurs

Drug Facts

Active Ingredients (in each effervescent tablet)

- Aspirin 325 mg (NSAID)
- Chlorpheniramine maleate 2 mg
- Phenylephrine bitartrate 7.5 mg
- Nasal decongestant

Purpose

Temporarily relieves these symptoms of the common cold:

- fever
- sneezing
- nasal congestion
- sore throat
- headache
- minor aches and pains
- runny nose

Warnings

Flu symptoms: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting may occur. Consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains a non-steroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning prescription (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Take more or for a longer time than directed

GOODSENSE[®]

Effervescent Cold Relief

Nasal Decongestant, Antihistamine
Pain Reliever/Fever Reducer

Distributed by:
Geiss, Destin & Dunn, Inc.
Peachtree City, GA 30269
www.valuelabels.com
1-866-696-0957


GOODSENSE[®]

NDC 50201-8736-0

Fast Relief Plus

Effervescent Cold Relief

Aspirin (NSAID)/Pain Reliever/Fever Reducer
Chlorpheniramine Maleate/Antihistamine
Phenylephrine Bitartrate/Nasal Decongestant



- FAST RELIEF OF:**
- Nasal Congestion
 - Runny Nose
 - Sneezing
 - Headache
 - Sinus Pain & Pressure



20 Effervescent Tablets

Drug Facts (continued)

- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- each tablet contains:
- sodium 464 mg
 - phenylketonurics: contains phenylalanine 9 mg per tablet
 - store at room temperature (59°-86°F)

Inactive ingredients

- acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Serious side effects associated with the use of this product can be reported to the following address:

Tower Laboratories LTD.
P.O. Box 306 8 Industrial Park Rd.
Centerbrook, CT 06409

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED



PN: 87360 GS C1
REV: 08/19

*This product is not manufactured, distributed or licensed by the maker of Alka-Seltzer Plus® Cold.

Drug Facts (continued)


When using this product

- do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.
- do not use more than directed
- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours



GOOD SENSE EFFERVESCENT COLD RELIEF

aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50201-8736
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ASPARTAME (UNII: Z0H242BBR1)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50201-8736-0	10 in 1 CARTON	08/15/2019	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC monograph final	part341	02/19/2013	12/31/2024
---------------------	---------	------------	------------

Labeler - Tower Laboratories Ltd. (001587203)

Registrant - Tower Laboratories Ltd. (001587203)

Establishment

Name	Address	ID/FEI	Business Operations
Tower Laboratories Ltd.		869024500	manufacture(50201-8736) , pack(50201-8736) , label(50201-8736)

Revised: 12/2022

Tower Laboratories Ltd.