

SUDAFED PE SINUS PRESSURE PLUS PAIN- acetaminophen and phenylephrine hydrochloride tablet, film coated
Kenvue Brands LLC

Sudafed PE Sinus Pressure Plus Pain

Drug Facts

| <i>Active ingredients (in each tablet)</i> | <i>Purpose</i> |
|---|-----------------------------|
| Acetaminophen 325 mg | Pain reliever/fever reducer |
| Phenylephrine HCl 5 mg | Nasal decongestant |

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - nasal congestion
- promotes sinus drainage
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**

| | |
|---------------------------------------|---|
| adults and children 12 years and over | <ul style="list-style-type: none">▪ take 2 tablets every 4 hours▪ do not take more than 10 tablets in 24 hours |
| children under 12 years | ask a doctor |

Other information

- store between 20-25°C (68-77°F)

- **do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, FD&C yellow no. 6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polysorbate 80, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide

Questions or comments?

call **1-888-217-2117** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

PREVIOUSLY SUDAFED PE [®] PRESSURE + PAIN
NDC 50580-435-01

SUDAFED PE[®]

SINUS
PRESSURE + PAIN

Acetaminophen, Phenylephrine HCl
Pain Reliever/Fever Reducer, Nasal Decongestant

actual size

MAXIMUM STRENGTH

- SINUS PRESSURE + CONGESTION
- SINUS HEADACHE

24 TABLETS

NON-DROWSY

SUDAFED^{PE}
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19084 USA ©J&JCI 2018

DrugFacts (continued)
Questions or comments?
call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

Open from other side



Do not use
■ If you are not sure whether a drug contains acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychobiotics, or Parkinson's disease), or if you have ever had an allergic reaction to this product or any of its ingredients, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients, ask a doctor or pharmacist before taking this product.

Warnings
Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg of acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.
■ With other drugs containing acetaminophen
■ 3 or more a day to help with pain while using this product
■ Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness
■ blisters
■ rash
■ If a skin reaction occurs, stop use and seek medical help right away.

Uses
Temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
■ sinus congestion and pressure
■ headache
■ sore throat and pain
■ nasal congestion
■ promotes sinus drainage
■ temporarily reduces fever

Active ingredients (in each tablet)
Phenylephrine HCl 5 mg.
Pain reliever/fever reducer
Nasal decongestant

Drug Facts
Keep this box for important information before using.
Important: Read a product information before using.

Other information
■ Store between 20-25°C (68-77°F)
■ do not use if blister units are torn or broken
Inactive ingredients carnau wax, FD&C yellow no. 6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polysorbate 80, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide

Directions
■ do not use more than directed (see overdose warning)
adults and children 12 years and over
■ take 2 tablets every 4 hours
■ do not take more than 10 tablets in 24 hours
children under 12 years
ask a doctor

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition:
■ redness or swelling
■ new symptoms occur
■ fever gets worse or lasts more than 3 days
■ pain or nasal congestion gets worse or lasts more than 7 days

Stop use and ask a doctor if
■ nervousness, dizziness, or sleepiness occur
■ When using this product do not exceed recommended dose
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

DrugFacts (continued)
Ask a doctor before use if you have
■ liver disease
■ heart disease
■ high blood pressure
■ thyroid disease
■ diabetes
■ trouble urinating due to an enlarged prostate gland

Does Not Contain
Pseudoephedrine


The makers of the SUDAFED[®] family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

PREVIOUSLY SUDAFED PE[®] PRESSURE + PAIN NDC 50580-435-01

SUDAFED^{PE}

**SINUS
PRESSURE + PAIN**

Acetaminophen, Phenylephrine HCl
Pain Reliever/Fever Reducer, Nasal Decongestant



24 TABLETS

MAXIMUM STRENGTH
• SINUS PRESSURE + CONGESTION
• SINUS HEADACHE

NON-DROWSY

30042 477/
120 483

30042 477/
120 483

SUDAFED PE SINUS PRESSURE PLUS PAIN

acetaminophen and phenylephrine hydrochloride tablet, film coated

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:50580-435 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-----------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| ALUMINUM OXIDE (UNII: LMI26O6933) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | |
| POWDERED CELLULOSE (UNII: SMD1X3XO9M) | |
| SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D) | |

Product Characteristics

| | | | |
|----------|----------------|--------------|-----------|
| Color | orange (PEACH) | Score | no score |
| Shape | OVAL | Size | 18mm |
| Flavor | | Imprint Code | SUPE;WL89 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:50580-435-01 | 2 in 1 CARTON | 06/17/2019 | |
| 1 | | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing | Application Number or Monograph | Marketing Start | Marketing End |
|-----------|---------------------------------|-----------------|---------------|
|-----------|---------------------------------|-----------------|---------------|

| Category | Citation | Date | Date |
|--------------------|----------|------------|------|
| OTC Monograph Drug | M012 | 06/17/2019 | |

Labeler - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC