

LINCOBAC 300- lincomycin injection
MWI/VetOne

LincoBac™ 300

(lincomycin injection, USP)

Injection

Swine Antibiotic

For use in swine weighing 300 pounds or more.

For Intramuscular Use in Swine Only

LincoBac™ 300 contains lincomycin hydrochloride, an antibiotic produced by *Streptomyces lincolnensis* var. *lincolnensis* which is chemically distinct from all other clinically available antibiotics and is isolated as a white crystalline solid.

ANADA 200-368, Approved by FDA

ADVERSE REACTIONS: The intramuscular administration to swine may cause a transient diarrhea or loose stools. Although this effect has rarely been reported, one must be alert to the possibility that it may occur. Should this occur, it is important that the necessary steps be taken to prevent the effects of dehydration.

DOSAGE AND ADMINISTRATION: For arthritis or mycoplasma pneumonia – 5 mg per pound of body weight intramuscularly once daily for three to seven days as needed. One mL per 60 lb body weight will provide 5 mg/lb.

For optimal results, initiate treatment as soon as possible.

As with any multi-dose vial, practice aseptic techniques in withdrawing each dose. Adequately clean and disinfect the vial closure prior to entry with a sterile needle and syringe. No vial closure should be entered more than 20 times.

□ **HOW SUPPLIED:** 100 mL multiple dose vial.

INDICATIONS FOR SWINE:

LincoBac™ 300 is indicated for the treatment of infectious forms of arthritis caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such as staphylococci, streptococci, *Erysipelothrix* and *Mycoplasma* spp. It is also indicated for the treatment of mycoplasma pneumonia.

□ **STORAGE:** Store at 20°-25°C (68°-77°F). Protect from freezing.

CONTRAINDICATIONS: As with all drugs, the use of **LincoBac™ 300** is contraindicated in animals previously found to be hypersensitive to the drug.

Residue Warning: Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

Human Warnings: Not for use in humans. Keep out of reach of children.

CAUTION: If no improvement is noted within 48 hours, consult a veterinarian.

For Intramuscular Use in Swine Over 300 lbs. NDC 13985-576-10

100 mL

VET ONE
Lincobac[®] 300
(lincomycin injection, USP)
Injection

Swine Antibiotic

Residue Warning: Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

HUMAN WARNINGS: Not for use in humans. Keep out of reach of children.

STORAGE: Store at 20°-25°C (68°-77°F). Protect from freezing.

4281012 100 mL (3.3 FL OZ)

INGREDIENTS: Contains per mL: Lincomycin hydrochloride equivalent to lincomycin 300 mg; also benzyl alcohol 9.45 mg added as a preservative.

INDICATIONS: Indicated for the treatment of arthritis caused by susceptible organisms and for mycoplasma pneumoniae.

Manufactured by: **Stranco-MTC Animal Health Inc.**, Carlsbad, CA, Canada NDC 294
www.lincobac.com

Distributed by: **MHI**, Boise, ID 83705
(888) 854-4381
www.VetOne.net

139857 5UN037 Rev. 04/14

3 13985 01239 6

FULL

USE THE OBSERVE/LABEL DIRECTIONS

NDC 13985-576-10

VET ONE
Lincobac[®] 300
(lincomycin injection, USP)
Injection

Swine Antibiotic

For use in swine weighing 300 pounds or more.

For Intramuscular Use in Swine Only

ANADA 200-368, Approved by FDA

Lincobac[®] 300 contains lincomycin hydrochloride, an antibiotic produced by *Streptomyces lincolnensis* var. *lincolnensis* which is chemically distinct from all other clinically available antibiotics and is isolated as a white crystalline solid.

INDICATIONS FOR SWINE: **Lincobac[®] 300** is indicated for the treatment of infectious forms of arthritis caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such

as staphylococci, streptococci, *Erysipelothrix* and *Mycoplasma* spp. It is also indicated for the treatment of mycoplasma pneumoniae.

CONTRAINDICATIONS: As with all drugs, the use of **Lincobac[®] 300** is contra-

indicated in animals previously found to be hypersensitive to the drug.

Residue Warning: Swine intended for human consumption should not be slaughtered within 48 hours of latest treatment.

Human Warnings: Not for use in humans. Keep out of reach of children.

CAUTION: If no improvement is noted within 48 hours, consult a veterinarian.

ADVERSE REACTIONS: The intramuscular administration to swine may cause a transient diarrhea or loose stools. Although this effect has rarely been reported, one must be alert to the possibility that it may occur. Should this occur, it is important that the necessary steps be taken to prevent the effects of dehydration.

DOSAGE AND ADMINISTRATION: For arthritis or mycoplasma pneumoniae - 5 mg per pound of body weight intramuscularly once daily for three to seven days as needed. One mL per 60 lb body weight will provide 5 mg/lb. For optimal results, initiate treatment as soon as possible.

As with any multi-dose vial, practice aseptic techniques in withdrawing each dose. Adequately clean and disinfect the vial closure prior to entry with a sterile needle and syringe. No vial closure should be entered more than 20 times.

INGREDIENTS: Each mL contains: Lincomycin hydrochloride equivalent to lincomycin 300 mg; also benzyl alcohol 9.45 mg added as a preservative.

HOW SUPPLIED: 100 mL multiple dose vial.

STORAGE: Store at 20°-25°C (68°-77°F). Protect from freezing.

TACTIC  OBSERVE/LABEL DIRECTIONS

VET ONE

Distributed by: **MHI**, Boise, ID 83705
(888) 854-4381
www.VetOne.net

INGREDIENTS: Contains per mL: Lincomycin hydrochloride equivalent to lincomycin 300 mg; also benzyl alcohol 9.45 mg added as a preservative.

INDICATIONS: Indicated for the treatment of arthritis caused by susceptible organisms and for mycoplasma pneumoniae.

Manufactured by: **Stranco-MTC Animal Health Inc.**, Carlsbad, CA, Canada NDC 294
www.lincobac.com

Distributed by: **MHI**, Boise, ID 83705
(888) 854-4381
www.VetOne.net

139857 5UN037 Rev. 04/14

3 13985 01239 6

LINCOBAC 300

lincomycin injection

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:13985-576
Route of Administration	INTRAMUSCULAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LINCOMYCIN HYDROCHLORIDE (UNII: M6T05Z2B68) (LINCOMYCIN - UNII:BOD072YW0F)	LINCOMYCIN	300 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	9.45 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-576-10	100 mL in 1 VIAL, GLASS		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200368	08/02/2006	

Labeler - MWI/VetOne (019926120)

Registrant - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda-MTC Animal Health		256232216	manufacture

Revised: 12/2018

MWI/VetOne