

CALACLEAR- pramoxine hydrochloride and zinc acetate lotion
Humco Holding Group, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Private Label Calaclear

Humco Calamine Lotion

Drug Facts

Active Ingredient

Pramoxine HCl 1%

Purpose

External Analgesic

Active Ingredient

Zinc Acetate 0.1%

Purpose

Skin Protectant

Use

Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.

Warnings

For external use only. Use only as directed.

Avoid contact with eyes and mucous membranes.

Ask a doctor before using

on children under 2 years of age.

When using this product. Discontinue use if

condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days. and consult a doctor.

Keep out of reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- Adults and children 2 yrs. of age and older. Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.
- Children under 2 years of age: Consult a doctor before use.


Other Information

- store at room temperature 15 ° - 30 °C (50 ° - 85 °F)

Inactive Ingredient

SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.

GNP Label

 NDC 24385-439-30

Caldyphen

Lotion Clear

External Analgesic / Skin Protectant

Relieves ITCHING due to:

- Poison Ivy • Insect Bites
- Poison Oak • Minor Skin Irritations

6 FL OZS (177 mL)

Drug Facts	Purpose External analgesic Skin protectant
Active ingredients Pramoxine HCl 1% Zinc Acetate 0.1%	External analgesic Skin protectant
Uses Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac and other minor skin irritations.	
Warnings For External Use Only. Use only as directed. ■ Avoid contact with eye and mucous membranes. ■ Ask a doctor before using on children under 2 years of age.	
Stop use and ask a doctor if ■ condition worsens ■ symptoms last for more than 7 days or clear up and occur again within a few days. Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions Adults and children 2 yrs. of age and older Children under 2 yrs. of age	Shake well before using. Cleanse the skin with soap and water. Let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort. Consult a doctor before use.
Inactive ingredients SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, and Purified Water.	
Other information Store at room temperature 15° - 30° (59°-86°F).	

Distributed by: AmerisourceBergen
1300 Morris Drive, Chesterbrook, PA 19087
Questions or Comments? ☎1-800-662-3435 • www.goodneighborpharmacy.com


0 87701 40045 2
ABC# 643-751
R082913RLG



Leader Label

LEADER[™]

NDC 70000-0397-1

Calamine Clear Lotion

External Analgesic | Skin Protectant

Relieves Itching Due to:
Poison Ivy, Poison Oak,
Insect Bites, Minor Skin
Irritations

COMPARE TO CALADRYL[®] CLEAR[®] LOTION active ingredients*

100% Money Back Guarantee

A45238
02 2024

6 FL OZ (177 mL)

Drug Facts

Active ingredients	Purpose
Pramoxine HCl 1%	External analgesic
Zinc Acetate 0.1%	Skin protectant

Uses

- temporarily relieves pain and itching associated with:
 - rashes due to poison ivy, poison oak or poison sumac
 - insect bites
 - minor skin irritation
 - minor cuts
- dries the oozing and weeping of poison ivy, poison oak and poison sumac

Warnings

- For external use only. Use only as directed.
- Avoid contact with eyes and mucous membranes.
- Ask a doctor before using on children under 2 years of age.

Stop use and ask a doctor if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately (1-800-222-1222).

Directions

- Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.
- Children under 2 years of age: Consult a doctor before use.

Other information

- Store at room temperature 15-30°C (59-86°F)

Inactive ingredients

- SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.

Questions or Comments? 1-800-662-3435

*This product is not manufactured or distributed by Valeant Pharmaceuticals, owner of the registered trademark Caladryl[®] Clear[®] Lotion.

© 2018 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY:
CARDINAL HEALTH
DUBLIN, OHIO 43007
www.mylister.com
1-800-300-6333

CIN 5-455795 REV. 12/18

100% Money Back Guarantee

Return to place of purchase.

0 96295 13509 1

Quality Choice Label

Drug Facts

Active ingredients	Purpose
Pramoxine HCl 1%	External analgesic
Zinc Acetate 0.1%	Skin protectant

Indications Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.

Warnings

- For external use only. Use only as directed.
- Avoid contact with eyes and mucous membranes.
- Ask a doctor before using on children under 2 years of age.

When using this product discontinue use if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately. *Continued* >

*This product is not manufactured or distributed by Johnson & Johnson Consumer Products Company, owner of the registered trademark Caladryl[®] Clear[™] Lotion

NDC 63868-097-06

QC QUALITY CHOICE

*Compare to Ingredient in Caladryl[®] Clear[™] Lotion

Anti-Itch Clear Lotion

Skin Protectant

External Analgesic,
Drying Agent
Helps Relieve Pain & Itch Caused by
Poison Ivy, Poison Oak
& Poison Sumac

6 fl OZ (177 mL)

Drug Facts (CONTINUED)

Directions

- Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion in the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.
- Children under 2 years of age: Consult a doctor before use.

Other information

- Store at room temperature 15-30°C (59-86°F)

Inactive ingredients

- SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.

R062814RLG

06 355 15 94565 4

SATISFACTION 100% GUARANTEED

Distributed by C.D.M.A., Inc.
43157 W. Nine Mile
Novi, MI 48376-0995
www.qualitychoice.com
Questions: 248-449-9300

MADE IN USA

Sunmark Label

sunmark™ COMPARE TO CALADRYL® CLEAR™ ACTIVE INGREDIENT* NDC 49348-610-36

caldyphen lotion

Clear

External analgesic
Skin protectant
Drying agent & itch relief

0 10939 14633 5

Distributed By McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee

6 FL OZ (177 mL)

Drug Facts	
Active Ingredients	Purpose
Pramoxine HCl 1% and Zinc Acetate 0.1%.....	External Analgesic/Skin Protectant
Uses Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac and other minor skin irritations.	
Warnings For External Use Only. Use only as directed. ■ Avoid contact with eye and mucous membranes. ■ Ask a doctor before using on children under 2 years of age.	
When using this product Discontinue use if condition worsens, does not improve or if symptoms persist for more than 7 days or clear up and occur again within a few days, and consult a doctor.	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a poison Control Center immediately.	
Directions	
Adults and children 2 yrs. of age and older	Shake well before using. Cleanse the skin with soap and water. Let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.
Children under 2 yrs. of age	Consult a doctor before use.
Inactive ingredients SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, and Purified Water.	
Other information Store at room temperature 15° - 30° (59°-86°F).	

*Caladryl® Clear™ is the registered trade mark of Warner-Lambert Co.

Publix Label

Publix

clear

ANTI-ITCH LOTION

Skin Protectant

Compare to Active Ingredients in
Caladryl® Clear™ Lotion

external analgesic • skin protectant
drying agent & itch relief

0 41415 06775 2

6 FL OZ (177mL)

Drug Facts	
Active Ingredients	Purpose
Pramoxine HCl 1% and Zinc Acetate 0.1%.....	External Analgesic/Skin Protectant
Uses Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.	
Warnings: For External Use Only. Use only as directed. ■ Avoid contact with eye and mucous membranes. ■ Ask a doctor before using on children under 2 years of age.	
When using this product Discontinue use if conditions worsens, does not improve or if symptoms persist for more than 7 days or clear up and occur again within a few days, and consult a doctor.	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions	
Adults and children 2 yrs. of age and older	Shake well before using. Cleanse the skin with soap and water. Let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.
Children under 2 yrs. of age	Consult a doctor before use.
Inactive Ingredients: SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, and Purified Water.	
Other Information: Store at room temperature 15° - 30° (59° - 86°F).	

*Caladryl® Clear™ is the registered trade mark of Warner-Lambert Co.
DISTRIBUTED BY PUBLIX SUPER MARKETS, L.L.C. 3300 PUBLIX CORPORATE PARKWAY, LAKELAND, FL 33811
800-242-1227 • publix.com • PUBLIX GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK.

Premiere Value Label

* Compare to the active ingredients in Caladryl® Clear™



CALACLEAR™

LOTION

For the Temporary Relief of Pain and Itching Associated with Minor Skin Irritations.



NET 6 FL. OZ. (177ML)

Drug Facts	
Active Ingredients Pramoxine HCl 1%.....External analgesic Zinc Acetate 0.1%.....Skin protectant	Purpose External analgesic.....Skin protectant
Uses Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac and other minor skin irritations.	
Warnings For External Use Only. Use only as directed. <ul style="list-style-type: none"> ■ Avoid contact with eye and mucous membranes. ■ Ask a doctor before using on children under 2 years of age. 	
Stop use and ask a doctor if ■ condition worsens ■ symptoms last for more than 7 days or clear up and occur again within a few days.	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions Adults and children 2 yrs. of age and older Shake well before using. Cleanse the skin with soap and water. Let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.	
Children under 2 yrs. of age Consult a doctor before use.	
Inactive ingredients SD Alcohol 368 2.5%, Camphor, Diazolidrilyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polyisobutyl 80, Propylene Glycol, Propylparaben, and Purified Water.	
Other information Store at room temperature 15° - 30° (59°-86°F).	
Questions or Comments? ☎ 1-800-662-3435	

* This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Caladryl® Clear®.

FROM HPLC



Distributed By:
 Pharmacy Value Alliance, LLC
 407 East Lancaster Avenue,
 Wayne, PA 19087

MADE IN USA

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund

DDM Label



Calaclear®

Lotion

Topical Analgesic • Skin Protectant
Drying Agent • Itch Relief

* Compare to the Active Ingredients of Caladryl® Clear® Lotion

6 fl oz (177 mL)

Drug Facts	
Active Ingredients Pramoxine HCl 1%.....External analgesic Zinc Acetate 0.1%.....Skin protectant	Purpose External analgesic.....Skin protectant
Uses dries the oozing and weeping of poison ivy, poison oak and poison sumac and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.	
Warnings <ul style="list-style-type: none"> ■ For external use only. Use only as directed. Avoid contact with eyes and mucous membranes. Ask a doctor before using on children under 2 years of age. 	
When using this product discontinue use if condition worsens or does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions Adults and children 2 yrs. of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.	
Children under 2 yrs. of age: Consult a doctor before use.	
Other information Store at room temperature 15° - 30° C (59°-86° F)	
Inactive ingredients SD Alcohol 368 2.5%, Camphor, Diazolidrilyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polyisobutyl 80, Propylene Glycol, Propylparaben, Purified Water.	

* This product is not manufactured or distributed by Johnson & Johnson Consumer Products Company, owner of the registered trademark Caladryl® Clear® Lotion.

Satisfactions guaranteed - If dissatisfied, return unused portion and package to store where purchased for a full refund. If unable to return to store, send reason for dissatisfaction, name, address and copy package to: Discount Drug Mart, 211 Commerce Dr., Medina, Ohio 44125



DISTRIBUTED BY:
 DRUG MART-FOOD FAIR
 211 Commerce Dr.
 MEDINA, OHIO 44125
 Questions or Comments? 1-800-933-6278
 www.DISCOUNT-DRUGMART.COM

CVS Label



Compare to the active ingredients in Caladryl Clear®*

Itch Relief

CLEAR SKIN PROTECTANT

Topical analgesic
Skin protectant
Drying action plus
itch relief

6 FL OZ (177 mL)



Drug Facts	Purpose External analgesic Skin protectant
Active Ingredients Pramoxine HCl 1% Zinc Acetate 0.1%	
Uses Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac. ■ other minor skin irritations.	
Warnings ■ For external use only	
When using this product ■ do not get into eyes	
Stop use and ask a doctor if ■ condition worsens ■ symptoms last more than 7 days or clear up and occur again within a few days	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions ■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, no more than 3 to 4 times daily. ■ Children under 2 years age: Consult a doctor before use.	
Other information Store at room temperature 15°-30°C (59°-86°F)	
Inactive ingredients SD Alcohol 388 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.	

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Caladryl Clear®.

#796623



0 50428 29088 0

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive,
Woonsocket, RI 02895
© 2015 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-17624



Compare to the active ingredients in Caladryl Clear®*

Itch Relief

CLEAR SKIN PROTECTANT

Topical analgesic
Skin protectant
Drying action plus
itch relief

6 FL OZ (177 mL)



Drug Facts	Purpose External analgesic Skin protectant
Active Ingredients Famoxone HCl 1% Zinc Acetate 0.1%	
Uses Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac and other minor skin irritations.	
Warnings ■ For External Use Only. Use only as directed. ■ Avoid contact with eyes and mucous membranes. ■ Ask a doctor before using on children under 2 years of age.	
Stop use and ask a doctor if ■ condition worsens ■ symptoms last for more than 7 days or clear up and occur again within a few days.	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a poison Control Center immediately.	
Directions Adults and children 2 yrs. of age and older Shake well before using. Cleanse the skin with soap and water. Let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort. Children under 2 yrs. of age: Consult a doctor before use.	
Inactive ingredients SD Alcohol 388 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, and Purified Water.	
Other information Store at room temperature 15°-30° (59°-86°F)	

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Caladryl Clear®.

#796623



0 50428 29088 0

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive,
Woonsocket, RI 02895
© 2015 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-17624



Harris Teeter Label

Drug Facts	
Active ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc Acetate 0.1%.....	Skin protectant
Uses	
Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.	
Warnings	
■ For External use only. Use only as directed.	
■ Avoid contact with eyes and mucous membranes.	
■ Ask a doctor before using on children under 2 years of age.	
When using this product	
Discontinue use if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
<i>Continued</i> ▶	

This product is not manufactured or distributed by Pfizer, owner of the registered trademark "Caladryl® Clear™ Lotion"

Compare to active ingredients in Caladryl® Clear™ Lotion*



Calaclear® Lotion

**Skin Protectant • Drying Agent
External Analgesic**

Topical relief of minor irritations due to:
Poison Ivy, Poison Oak, Poison Sumac
and other minor skin irritations.

6 FL OZ (177 mL)

Drug Facts (CONTINUED)	
Directions	
■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.	
■ Children under 2 years of age: Consult a doctor before use.	
Other information	
Store at room temperature 15-30°C (50-86°F)	
Inactive ingredients	
SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.	
R020311.RL.G	
0 72036 98725 9	
PROUDLY DISTRIBUTED BY: HARRIS TEETER® MATTHEWS, NC 28105	
your <i>plus</i> guarantee.	
1-800-432-6111 or harristeeter.com	

Best Choice Label

Compare to the Active Ingredients of Caladryl® Clear® Lotion

Best Choice®

Anti-Itch Clear

**Topical Analgesic • Skin Protectant
Drying Agent • Itch Relief Lotion**

6 fl oz (177 mL)

Drug Facts	Purpose
Active Ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc Acetate 0.1%.....	Skin protectant
Indication	Indication
Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.	Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.
Warnings	Warnings
■ For external use only. Use only as directed.	■ For external use only. Use only as directed.
■ Avoid contact with eyes and mucous membranes.	■ Avoid contact with eyes and mucous membranes.
■ Ask a doctor before using on children under 2 years of age.	■ Ask a doctor before using on children under 2 years of age.
When using this product	When using this product
Discontinue use if condition worsens or does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.	Discontinue use if condition worsens or does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.
Keep out of reach of children.	Keep out of reach of children.
In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.
Directions	Directions
■ Adults and children 2 yrs. of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.	■ Adults and children 2 yrs. of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.
■ Children under 2 yrs. of age: Consult a doctor before use.	■ Children under 2 yrs. of age: Consult a doctor before use.
Other information	Other information
Store at room temperature.	Store at room temperature.
Inactive ingredients	Inactive ingredients
SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water	SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water

R04103RLG

BEST CHOICE

Best Choice
100% Guaranteed

www.bestchoicebrand.com

PROUDLY DISTRIBUTED BY:
VALU MERCHANTSERS. CO.
5000 KANSAS AVE
KANSAAS CITY, MO 66106

0 70038 20052 9

HEB Label



Drug Facts

Active ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc acetate 0.1%.....	Skin protectant

Uses Dries the oozing and weeping, and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.

Warnings: ■ **For External use only.** Use only as directed
 ■ **Avoid contact with eyes and mucous membranes**
 ■ **Ask a doctor before using on children under 2 yrs. of age**

When using this product ■ **Discontinue use** if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days and consult a doctor.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions
 ■ **Adults and children 2 yrs. and older:** Shake well before using. Wash the skin with soap and water and let dry. Apply to the affected area using cotton or soft cloth not more than 3 to 4 times daily as needed for comfort.
 ■ **Children under 2 yrs. of age:** Consult a doctor.
 ■ **Other information** ■ Store at room temperature 15° - 30°C (50° - 86°F)

Inactive ingredients ■ SD alcohol 38B 2.5%, camphor, diazolidinyl urea, fragrances, glycerin, hydroxypropyl methylcellulose, methylparaben, polysorbate 80, propylene glycol, propylparaben and purified water.

MADE WITH PRIDE AND CARE FOR H-E-B, SAN ANTONIO, TX 78204
 Questions or Comments? Call 1-800-662-3435 1104
 We believe the high quality of this H-E-B product makes it an outstanding value. We hope you'll agree. If not, we'll cheerfully refund your money. 0

Equate Label

Drug Facts

Active ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc Acetate 0.1%.....	Skin protectant

Uses Dries the oozing and weeping and temporarily relieves pain and itching of ■ poison ivy ■ poison oak ■ poison sumac ■ other minor skin irritations

Warnings
 ■ **For external use only**

When using this product ■ do not get into eyes

Stop use and ask a doctor if
 ■ condition worsens ■ symptoms last more than 7 days or clear up and occur again within a few days

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center (1-800-222-1222) right away.

Continued ▶

NDC 49035-402-96

equate™

Calamine Clear LOTION

Drug Facts (continued)

Directions
 ■ **Adults and children 2 years of age and older:** Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, no more than 3 to 4 times daily.
 ■ **Children under 2 years of age:** Consult a doctor before use.

Other information
 Store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients
 Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water, SD Alcohol 38B 2.5%.

- Skin protectant
- Poison Ivy, Oak and Sumac drying lotion
- Relieves itching from minor skin irritations

6 FL OZ (177mL)

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
 Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

R040721

*This product is not manufactured or distributed by Valeant Pharmaceuticals, owner of the registered trademark Caladryl® Clear® Lotion.

Drug Facts

Active ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc Acetate 0.1%.....	Skin protectant

Uses Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations

Warnings

- For external use only. Use only as directed.
- Avoid contact with eyes and mucous membranes
- Ask a doctor before using on children under 2 years of age.

When using this product Discontinue use if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Continued ▶

*This product is not manufactured or distributed by Valeant Pharmaceuticals, owner of the registered trademark Caladryl® Clear® Lotion.

R080017.RLG

NDC 49035-402-95

equate™

Calamine Clear LOTION



- Skin Protectant
- Poison Ivy, Oak and Sumac drying lotion
- Relieves itching from minor skin irritations



6 FL OZ (177mL)

Drug Facts (continued)

Directions

■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, no more than 3 to 4 times daily.

■ Children under 2 years age: Consult a doctor before use.

Other information

Store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients

SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.

DISTRIBUTED BY: Wal-Mart Stores, Inc., Bentonville, AR 72716

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.



WM Label

Drug Facts

Active Ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc Acetate 0.1%.....	Skin protectant

Uses Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.

Warnings

- For external use only. Use only as directed.
- Avoid contact with eyes and mucous membranes.
- Ask a doctor before using on children under 2 years of age.

When using this product Discontinue use if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days, and consult a doctor.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Continued ▶

*This product is not manufactured or distributed by Johnson & Johnson Consumer Products Company, owner of the registered trademark Caladryl® Clear™ Lotion

NDC 0095 400-95

*Compare to Caladryl® Clear™ Lotion

CALACLEAR® LOTION

LOCIÓN CALACLEAR®

The Healthy Skin Care Solution

- External analgesic
- Skin protectant
- Drying agent & itch relief

6 FL OZ (177 mL)

6 ONZAS FL (177 mL)

HUMCO

Drug Facts (CONTINUED)

Directions

■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.

■ Children under 2 years of age: Consult a doctor before use.

Other information

Store at room temperature 15-30°C (50-86°F)

Inactive ingredients

SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water.

HUMCO, Texarkana, TX 75501
Questions or Comments? ☎1-800-662-3435

Go to: www.simplehomeremedies.com to find helpful hints and learn more about Caladryl™ Lotion.

MADE IN USA



R082316RLGF



A45240
02 2024

Levante etiqueta para información

RA Label

Compare to the active ingredients of Caladryl® Clear® Lotion*

CALACLEAR LOTION

PRAMOXINE HCl 1%
ZINC ACETATE 0.1%

TOPICAL ANALGESIC SKIN PROTECTANT

relieves itching due to
poison ivy, poison oak, insect bites
& minor skin irritations

RITE AID PHARMACY

6 FL OZ (177 mL)

Drug Facts	
Active ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc Acetate 0.1%.....	Skin protectant
Uses Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.	
Warnings ■ For external use only. Use only as directed	
■ Avoid contact with eyes and mucous membranes	
■ Ask a doctor before using on children under 2 years of age	
Stop use and ask a doctor if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.	
Directions	
■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.	
■ Children under 2 years of age: Consult a doctor before use	
Other information ■ Store at room temperature 15-30°C (59-86°F)	
Inactive ingredients SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water	
Questions or Comments? 1-800-662-3435	

*This product is not manufactured or distributed by Valeant Pharmaceuticals, distributor of Caladryl® Clear Lotion.

DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE,
CAMP HILL, PA 17011
www.riteaid.com



0 11822 46461 1

Top Care Label

NDC 06809-287-56

TopCare health™

COMPARE TO CALADRYL® CLEAR® LOTION ACTIVE INGREDIENT*

Calaclear Lotion

TOPICAL ANALGESIC SKIN PROTECTANT

Relieves Itching Due to:

- Poison Ivy
- Poison Oak
- Insect Bites
- Minor Skin Irritations

OUR PHARMACISTS RECOMMEND

6 FL OZ (177 mL)

Drug Facts	
Active ingredients	Purpose
Pramoxine HCl 1%.....	External analgesic
Zinc Acetate 0.1%.....	Skin protectant
Uses Dries the oozing and weeping and temporarily relieves pain and itching of poison ivy, poison oak, and poison sumac or other minor skin irritations.	
Warnings ■ For external use only. Use only as directed	
■ Avoid contact with eyes and mucous membranes	
■ Ask a doctor before using on children under 2 years of age	
Stop use and ask a doctor if condition worsens, does not improve or if symptoms persist for more than 7 days, or clear up and occur again within a few days.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center 1-800-222-1222 immediately.	
Directions	
■ Adults and children 2 years of age and older: Shake well before using. Cleanse the skin with soap and water and let dry before each use. Apply lotion to the affected area using cotton or soft cloth, not more than 3 to 4 times daily as needed for comfort.	
■ Children under 2 years of age: Consult a doctor before use	
Other information ■ Store at room temperature 15-30°C (59-86°F)	
Inactive ingredients SD Alcohol 38B 2.5%, Camphor, Diazolidinyl Urea, Fragrances, Glycerin, Hydroxypropyl Methylcellulose, Methylparaben, Polysorbate 80, Propylene Glycol, Propylparaben, Purified Water	

*This product is not manufactured or distributed by Valeant Pharmaceuticals, owner of the registered trademark Caladryl® Clear® Lotion.

DISTRIBUTED BY: TOPCO ASSOCIATES, LLC
ELI CRONE VILLAGE, L 5007 ©TOPCO HEALTHCARE
QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com
Visit here for more information:
<http://topbrands.com/48uibe>

ROSC019



0 36800 44340 2

CALACLEAR

pramoxine hydrochloride and zinc acetate lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0395-9108
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 mL

ZINC ACETATE (UNII: FM5526K07A) (ZINC CATION - UNII:13S1S8SF37)

ZINC CATION

1 mg
in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
CAMPHOR (NATURAL) (UNII: N20HL7Q941)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0395-9108-96	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/25/1998	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part347	03/25/1998	

Labeler - Humco Holding Group, Inc. (825672884)

Registrant - Humco Holding Group, Inc. (825672884)

Establishment

Name	Address	ID/FEI	Business Operations
Humco Holding Group, Inc.		825672884	manufacture(0395-9108) , analysis(0395-9108) , pack(0395-9108) , label(0395-9108)

Revised: 12/2021

Humco Holding Group, Inc.