

EQUALINE ASPIRIN- aspirin tablet, film coated
United Natural Foods, Inc. dba UNFI

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

SuperValu Inc. Aspirin Tablets Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - redness or swelling is present
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Inactive ingredients

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

Questions?

1-855-423-2630

Package/Label Principal Display Panel

compare to Bayer® Aspirin active ingredient

for adults

aspirin tablets

pain reliever/fever reducer (NSAID)

TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER BEFORE USING THIS PRODUCT FOR YOUR HEART.

100 tablets, 325 mg

actual size

compare to Bayer® Aspirin active ingredient**
NDC 41163-405-78

EQUALINE®
for adults
aspirin tablets
pain reliever/fever reducer (NSAID)

TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER BEFORE USING THIS PRODUCT FOR YOUR HEART.

100 tablets, 325 mg actual size

DO NOT USE IF FINGERED SEAL (NEVER OPEN) IS BROKEN OR MISSING.

MADE IN CHINA
MAY CONTAIN MILK AND SOY PROTEIN
DISTRIBUTED BY UNR
OF CALIFORNIA, PHOENIX, AZ 02018 USA
855-423-2630

Drug Facts (continued)
Inactive ingredients: corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

Directions ■ Drink a full glass of water with each dose. ■ Adults and children 12 years and over: 1 or 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours. ■ Children under 12 years: consult a doctor.

Warnings
■ Bleeding: Stop using this product and call your doctor if you notice any of the following signs of bleeding: stomach pain, black stools, or blood in your urine or stool. ■ Pregnancy: If you are pregnant or breastfeeding, ask a health professional before use. ■ Allergic reactions: If you have had a severe allergic reaction to aspirin, you should not use this product. ■ Liver and kidney disease: If you have liver or kidney disease, you should not use this product. ■ Stomach ulcers: If you have stomach ulcers, you should not use this product. ■ Heart disease: If you have heart disease, you should not use this product. ■ High blood pressure: If you have high blood pressure, you should not use this product. ■ Low blood pressure: If you have low blood pressure, you should not use this product. ■ Headaches: If you have headaches, you should not use this product. ■ Dizziness: If you have dizziness, you should not use this product. ■ Stomach pain: If you have stomach pain, you should not use this product. ■ Nausea: If you have nausea, you should not use this product. ■ Vomiting: If you have vomiting, you should not use this product. ■ Diarrhea: If you have diarrhea, you should not use this product. ■ Constipation: If you have constipation, you should not use this product.

Drug Facts
100 tablets, 325 mg actual size

Drug Facts (continued)

Warnings

- Bleeding: Stop using this product and call your doctor if you notice any of the following signs of bleeding: stomach pain, black stools, or blood in your urine or stool.
- Pregnancy: If you are pregnant or breastfeeding, ask a health professional before use.
- Allergic reactions: If you have had a severe allergic reaction to aspirin, you should not use this product.
- Liver and kidney disease: If you have liver or kidney disease, you should not use this product.
- Stomach ulcers: If you have stomach ulcers, you should not use this product.
- Heart disease: If you have heart disease, you should not use this product.
- High blood pressure: If you have high blood pressure, you should not use this product.
- Low blood pressure: If you have low blood pressure, you should not use this product.
- Headaches: If you have headaches, you should not use this product.
- Dizziness: If you have dizziness, you should not use this product.
- Stomach pain: If you have stomach pain, you should not use this product.
- Nausea: If you have nausea, you should not use this product.
- Vomiting: If you have vomiting, you should not use this product.
- Diarrhea: If you have diarrhea, you should not use this product.
- Constipation: If you have constipation, you should not use this product.

Directions

- Drink a full glass of water with each dose.
- Adults and children 12 years and over: 1 or 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours.
- Children under 12 years: consult a doctor.

Warnings

- Bleeding: Stop using this product and call your doctor if you notice any of the following signs of bleeding: stomach pain, black stools, or blood in your urine or stool.
- Pregnancy: If you are pregnant or breastfeeding, ask a health professional before use.
- Allergic reactions: If you have had a severe allergic reaction to aspirin, you should not use this product.
- Liver and kidney disease: If you have liver or kidney disease, you should not use this product.
- Stomach ulcers: If you have stomach ulcers, you should not use this product.
- Heart disease: If you have heart disease, you should not use this product.
- High blood pressure: If you have high blood pressure, you should not use this product.
- Low blood pressure: If you have low blood pressure, you should not use this product.
- Headaches: If you have headaches, you should not use this product.
- Dizziness: If you have dizziness, you should not use this product.
- Stomach pain: If you have stomach pain, you should not use this product.
- Nausea: If you have nausea, you should not use this product.
- Vomiting: If you have vomiting, you should not use this product.
- Diarrhea: If you have diarrhea, you should not use this product.
- Constipation: If you have constipation, you should not use this product.

EQUALINE ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-405
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TALC (UNII: 7SEV7J4R1U)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L37J
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-405-78	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/15/2021	04/30/2024
2	NDC:41163-405-87	300 in 1 BOTTLE; Type 0: Not a Combination Product	07/15/2021	08/31/2023

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/15/2021	04/30/2024

Labeler - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 8/2023

United Natural Foods, Inc. dba UNFI