

MOTION SICKNESS LESS DROWSY FORMULA- meclizine hcl tablet
Strategic Sourcing Services LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to an enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, the first dose should be taken 1 hour before starting activity
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor
- do not exceed 2 tablets in 24 hours

Other information

- store at room temperature 20°-25°C (68°-77°F)

Inactive ingredients

anhydrous lactose, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminium lake, magnesium stearate, microcrystalline cellulose, sodium starch glycolate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO DRAMAMINE® LESS DROWSY FORMULA ACTIVE INGREDIENT*

Helps to prevent nausea, vomiting, or dizziness associated with motion sickness

Motion Sickness

Less Drowsy Formula

For children & adults

MECLIZINE HYDROCHLORIDE

ANTIEMETIC

ALL DAY RELIEF

TABLETS 25 mg EACH

*This product is not manufactured or distributed by Prestige Brands, Inc., owner of the registered trademark Dramamine® Less Drowsy Formula.

Another Quality Product Distributed by McKesson


One Post Street, San Francisco, CA 94104

Please visit us at www.sunmarkbrand.com

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Product Label



sunmark®

motion sickness

Less Drowsy Formula



For children & adults
MECLIZINE HYDROCHLORIDE
ANTIEMETIC

ALL DAY RELIEF
8 TABLETS 25 mg EACH

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

*This product is not manufactured or distributed by Prestige Brands, Inc., owner of the registered trademark Dramamine® Less Drowsy Formula.

MCKESSON

Another Quality Product Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
Please visit us at www.sunmarkbrand.com

PLD-C
FC001824



Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.
Warnings Do not use for children under 12 years of age unless directed by a doctor.
Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to an enlargement of the prostate gland
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ to prevent motion sickness, the first dose should be taken 1 hour before starting activity ■ adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor ■ do not exceed 2 tablets in 24 hours
Other information ■ store at room temperature 20°-25°C (68°-77°F)
Inactive ingredients anhydrous lactose, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, magnesium stearate, microcrystalline cellulose, sodium starch glycolate
Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST



Sunmark Motion Sickness Less Drowsy Tablet

MOTION SICKNESS LESS DROWSY FORMULA

meclizine hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-363
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

Product Characteristics

Color	WHITE, YELLOW	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	TCL086
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-363-67	1 in 1 CARTON	07/09/2010	12/31/2020
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 9/2019

Strategic Sourcing Services LLC