

**HEMAX- ascorbic acid, dl-alpha tocopheryl acetate, l-5-methyltetrahydrofolate calcium, cyanocobalamin, biotin, carbonyl iron, cupric sulfate tablet**

**Pronova Corporation**

-----  
**HEMAX™**

<b>Supplement Facts</b>		
<b>Serving Size:</b> 1 Tablet / Servings Per Container: 90		
	<b>Amount Per Serving</b>	<b>% Daily Value</b>
Vitamin C (as Ascorbic Acid)	500 mg	556%
Vitamin E (as dl-Alpha Tocopheryl Acetate)	13.5 mg	90%
Folate (as L-5-methyltetrahydrofolate Calcium)	1,020 mcg DFE (600 mcg L-5-MTH)	255%
Vitamin B12 (as Cyanocobalamin)	60 mcg	2,500%
Biotin	150 mcg	500%
Iron (as Carbonyl Iron)	150 mg	833%
Copper (as Cupric Sulfate)	3 mg	333%
*Daily Value not established		

**OTHER INGREDIENTS:** Microcrystalline Cellulose, Citric Acid, Stearic Acid, Dicalcium Phosphate, Croscarmellose Sodium, Polyvinylpyrrolidone, Hypromellose, Magnesium Stearate, Silicon Dioxide, Starch, Glucose, Calcium Carbonate, Crospovidone, Isomalt, Sodium Lauryl Sulfate, Maltodextrin, Triglycerides, Sucrose, Soy Polysaccharide, Glycerin, Mineral Oil, Polyethylene Glycol, Artificial French Vanilla Flavor, Annatto Extract (color), Propylene Glycol, Tricalcium Phosphate, Potassium Hydroxide and Talc.

**Contains** soy, a bioengineered food ingredient.

**HEMAX™** is a multivitamin/multimineral dietary supplement indicated for use in improving the nutritional status of patients with iron deficiency.†

†This statement has not been evaluated by the FDA. This product is not intended to diagnose, treat, cure, or prevent any disease.

**WARNING** Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6 years of age. **KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.

**DO NOT USE HEMAX™** for individuals with a known hypersensitivity to any of the ingredients. Hemochromatosis and hemosiderosis are contraindications to iron therapy. This product is not for use in children under 12 years of age.

**WARNING:**

Consuming this product can expose you to chemicals including, lead and mercury, which are known to the State of California to cause cancer and birth defects or other reproductive harm. For more information go to [www.P65Warnings.ca.gov/food](http://www.P65Warnings.ca.gov/food).

**CAUTION:**

This product contains 1.7mg DFE (1mg) of folic acid, the maximum safe Reference Daily Intake (RDI) amount for this nutrient. Folic Acid alone should not be used for individuals with pernicious anemia and other megaloblastic anemias where Vitamin B12 is deficient. Folic acid in amounts above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Allergic sensitization has been reported following both oral and parenteral administration of folic acid. Pregnant women, nursing mothers or women prior to conception who take **HEMAX™** should take this supplement only after their medical condition and other drugs, herbs and/or supplements consumption are considered. Use for elderly individuals should be cautious due to the greater frequency of decreased hepatic, renal, or cardiac function.

**ADVERSE REACTIONS:**

Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

**DIRECTIONS FOR USE:**

One tablet daily with or without food or as recommended by your healthcare provider.

**OTHER INFORMATION:**

**HEMAX™** is supplied in child-resistant bottles of 90, 30 and 4 tablets. Store at room temperature, USP.

**To report a serious adverse event or to obtain product information, contact (866) 703-3508.**

**Manufactured in the USA for:**

**Pronova Corporation  
Miami, Florida 33155**

**DIETARY SUPPLEMENT**

**With 100% Bioactive Folate**

**EZY-DOSE TABLETS**

**DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

# Packaging

90 TABLETS PACKAGE

DIETARY SUPPLEMENT

With 100% Bioactive Folate

# HEMAX™

90 EZY-DOSE TABLETS

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

## Supplement Facts

Serving Size: 1 Tablet / Servings Per Container: 90

	Amount Per Serving	% Daily Value
Vitamin C (as Ascorbic Acid)	500 mg	556%
Vitamin E (as dl-Alpha Tocopheryl Acetate)	13.5 mg	90%
Folate (as L-5-methyltetrahydrofolate Calcium) (600 mcg L-5-MTH)	1,020 mcg DFE	255%
Vitamin B12 (as Cyanocobalamin)	60 mcg	2500%
Biotin	150 mcg	500%
Iron (as Carbonyl Iron)	150 mg	833%
Copper (as Cupric Sulfate)	3 mg	333%

\*Daily Value not established.

**OTHER INGREDIENTS:** Microcrystalline Cellulose, Citric Acid, Stearic Acid, Dicalcium Phosphate, Croscarmellose Sodium, Polyvinylpyrrolidone, Hypromellose, Magnesium Stearate, Silicon Dioxide, Starch, Glucose, Calcium Carbonate, Crospovidone, Isomalt, Sodium Lauryl Sulfate, Maltodextrin, Triglycerides, Sucrose, Soy Polysaccharide, Glycerin, Mineral Oil, Polyethylene Glycol, Artificial French Vanilla Flavor, Annatto Extract (color), Propylene Glycol, Tricalcium Phosphate, Potassium Hydroxide and Talc. **Contains** soy, a bioengineered food ingredient.

3 67555 13190 1

PEEL HERE FOR MORE INFORMATION

Lot # & Exp. Date:

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

## Supplement Facts

Serving Size: 1 Tablet / Servings Per Container: 90

	Amount Per Serving	% Daily Value
Vitamin C (as Ascorbic Acid)	500 mg	556%
Vitamin E (as dl-Alpha Tocopheryl Acetate)	13.5 mg	90%
Folate (as L-5-methyltetrahydrofolate Calcium) (600 mcg L-5-MTH)	1,020 mcg DFE	255%
Vitamin B12 (as Cyanocobalamin)	60 mcg	2500%
Biotin	150 mcg	500%
Iron (as Carbonyl Iron)	150 mg	833%
Copper (as Cupric Sulfate)	3 mg	333%

\*Daily Value not established.

**OTHER INGREDIENTS:** Microcrystalline Cellulose, Citric Acid, Stearic Acid, Dicalcium Phosphate, Croscarmellose Sodium, Polyvinylpyrrolidone, Hypromellose, Magnesium Stearate, Silicon Dioxide, Starch, Glucose, Calcium Carbonate, Crospovidone, Isomalt, Sodium Lauryl Sulfate, Maltodextrin, Triglycerides, Sucrose, Soy Polysaccharide, Glycerin, Mineral Oil, Polyethylene Glycol, Artificial French Vanilla Flavor, Annatto Extract (color), Propylene Glycol, Tricalcium Phosphate, Potassium Hydroxide and Talc. **Contains** soy, a bioengineered food ingredient.

Lot # & Exp. Date:

3 67555 13190 1

PEEL HERE FOR MORE INFORMATION

**HEMAX™** is a multivitamin/multimineral dietary supplement indicated for use in improving the nutritional status of patients with iron deficiency. †

†This statement has not been evaluated by the FDA. This product is not intended to diagnose, treat, cure, or prevent any disease.

**WARNING:** Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6 years of age. **KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.

**DO NOT USE HEMAX™** for individuals with a known hypersensitivity to any of the ingredients. Hemochromatosis and hemosiderosis are contraindications to iron therapy. This product is not for use in children under 12 years of age.

**WARNING:** Consuming this product can expose you to chemicals including lead and mercury, which are known to the State of California to cause cancer and birth defects or other reproductive harm. For more information go to [www.P65Warnings.ca.gov/food](http://www.P65Warnings.ca.gov/food).

**CAUTION:** This product contains 1.7mg DFE (1mg) of folic acid, the maximum safe Reference Daily Intake (RDI) amount for this nutrient. Folic Acid alone should not be used for individuals with pernicious anemia and other megaloblastic anemias where Vitamin B12 is deficient. Folic acid in amounts above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Allergic sensitization has been reported following both oral and parenteral administration of folic acid. Pregnant women, nursing mothers or women prior to conception who take **HEMAX™** should take this supplement only after their medical condition and other drugs, herbs and/or supplements consumption are considered. Use for elderly individuals should be cautious due to the greater frequency of decreased hepatic, renal, or cardiac function.

**ADVERSE REACTIONS:** Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

**DIRECTIONS FOR USE:** One tablet daily with or without food or as recommended by your healthcare provider.

**OTHER INFORMATION:** **HEMAX™** is supplied in child-resistant bottles of 90, 30 and 4 tablets. Store at room temperature, USP. **To report a serious adverse event or to obtain product information, contact (866) 703-3508.**

**Manufactured in the USA for:**

**PRONOVA**

Pronova Corporation  
Miami, Florida 33155

30 TABLETS PACKAGE

DIETARY SUPPLEMENT

With 100% Bioactive Folate

**HEMAX™**

30 EZY-DOSE TABLETS

PRONOVA

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

**Supplement Facts**  
Serving Size: 1 Tablet

	Amount Per Serving	% Daily Value
Vitamin C (as Ascorbic Acid)	500 mg	556%
Vitamin E (as d-Alpha Tocopheryl Acetate)	13.5 mg	90%
Folate (as L-5-Methyltetrahydrofolate Calcium) (600 mcg L-5-MTHF)	1,020 mcg DFE	255%
Vitamin B12 (as Cyanocobalamin)	60 mcg	2,500%
Biotin	150 mcg	500%
Iron (as Carbonyl Iron)	150 mg	833%
Copper (as Cupric Sulfate)	3 mg	333%

\*Daily Value not established.

**OTHER INGREDIENTS:** Microcrystalline Cellulose, Citric Acid, Stearic Acid, Dicalcium Phosphate, Croscarmellose Sodium, Polyvinylpyrrolidone, Hypromellose, Magnesium Stearate, Silicon Dioxide, Starch, Glucose, Calcium Carbonate, Crospovidone, Isomalt, Sodium Lauryl Sulfate, Maltodextrin, Triglycerides, Sucrose, Soy Polysaccharide, Glycerin, Mineral Oil, Polyethylene Glycol, Artificial French Vanilla Flavor, Annatto Extract (color), Propylene Glycol, Tricalcium Phosphate, Potassium Hydroxide and Talc. **Contains soy, a bioengineered food ingredient.**

Lot #: \_\_\_\_\_  
Exp. Date: \_\_\_\_\_

3 67555 13130 7

PEEL HERE FOR MORE INFORMATION

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

**Supplement Facts**

Serving Size: 1 Tablet

	Amount Per Serving	% Daily Value
Vitamin C (as Ascorbic Acid)	500 mg	556%
Vitamin E (as d-Alpha Tocopheryl Acetate)	13.5 mg	90%
Folate (as L-5-Methyltetrahydrofolate Calcium) (600 mcg L-5-MTHF)	1,020 mcg DFE	255%
Vitamin B12 (as Cyanocobalamin)	60 mcg	2,500%
Biotin	150 mcg	500%
Iron (as Carbonyl Iron)	150 mg	833%
Copper (as Cupric Sulfate)	3 mg	333%

\*Daily Value not established.

**OTHER INGREDIENTS:** Microcrystalline Cellulose, Citric Acid, Stearic Acid, Dicalcium Phosphate, Croscarmellose Sodium, Polyvinylpyrrolidone, Hypromellose, Magnesium Stearate, Silicon Dioxide, Starch, Glucose, Calcium Carbonate, Crospovidone, Isomalt, Sodium Lauryl Sulfate, Maltodextrin, Triglycerides, Sucrose, Soy Polysaccharide, Glycerin, Mineral Oil, Polyethylene Glycol, Artificial French Vanilla Flavor, Annatto Extract (color), Propylene Glycol, Tricalcium Phosphate, Potassium Hydroxide and Talc. **Contains soy, a bioengineered food ingredient.**

Lot# &amp; Exp. Date:



PEEL HERE FOR MORE INFORMATION

HEMAX™ is a multivitamin/multimineral dietary supplement indicated for use in improving the nutritional status of patients with iron deficiency.<sup>†</sup>

<sup>†</sup>This statement has not been evaluated by the FDA. This product is not intended to diagnose, treat, cure, or prevent any disease.

**WARNING:** Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6 years of age. **KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.

**DO NOT USE HEMAX™** for individuals with a known hypersensitivity to any of the ingredients. Hemochromatosis and hemosiderosis are contraindications to iron therapy. This product is not for use in children under 12 years of age.

**CAUTION:** This product contains 1.7mg DFE (1mg) of folic acid, the maximum safe Reference Daily Intake (RDI) amount for this nutrient. Folic Acid alone should not be used for individuals with pernicious anemia and other megaloblastic anemias where Vitamin B12 is deficient. Folic acid in amounts above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Allergic sensitization has been reported following both oral and parenteral administration of folic acid. Pregnant women, nursing mothers or women prior to conception who take HEMAX™ should take this supplement only after their medical condition and other drugs, herbs and/or supplements consumption are considered. Use for elderly individuals should be cautious due to the greater frequency of decreased hepatic, renal, or cardiac function.

**ADVERSE REACTIONS:** Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

**DIRECTIONS FOR USE:** One tablet daily with or without food or as recommended by your healthcare provider.

**OTHER INFORMATION:** HEMAX™ is supplied in child-resistant bottles of 90, 30 and 4 tablets. Store at room temperature, USP.

To report a serious adverse event or to obtain product information, contact (866) 703-3508.

Manufactured in the USA for:



## HEMAX

ascorbic acid, dl-alpha tocopheryl acetate, l-5-methyltetrahydrofolate calcium, cyanocobalamin, biotin, carbonyl iron, cupric sulfate tablet

### Product Information

<b>Product Type</b>	DIETARY SUPPLEMENT	<b>Item Code (Source)</b>	NHRIC:67555-131
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASCORBIC ACID</b> (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	500 mg
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8) (.ALPHA.-TOCOPHEROL, DL- - UNII:7QWA1RIO01)	.ALPHA.-TOCOPHEROL, DL-	13.5 mg
<b>LEVOMEFOLATE CALCIUM</b> (UNII: A9R10K3F2F) (LEVOMEFOLIC ACID - UNII:8S95DH25XC)	LEVOMEFOLATE CALCIUM	600 ug
<b>CYANOCOBALAMIN</b> (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	60 ug
<b>BIOTIN</b> (UNII: 6S06U10H04) (BIOTIN - UNII:6S06U10H04)	BIOTIN	150 ug
<b>IRON</b> (UNII: E1UOL152H7) (IRON - UNII:E1UOL152H7)	IRON	150 mg
<b>CUPRIC SULFATE</b> (UNII: LRX7AJ16DT) (CUPRIC CATION - UNII:8CBV67279L)	CUPRIC CATION	3 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>ANHYDROUS DIBASIC CALCIUM PHOSPHATE</b> (UNII: L11K75P92J)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, RICE</b> (UNII: 4DGK8B7I3S)	
<b>ANHYDROUS DEXTROSE</b> (UNII: 5SL0G7R0OK)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>ISOMALT</b> (UNII: S870P55O2W)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	

<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>SOYBEAN COTYLEDON CELL WALL POLYSACCHARIDES</b> (UNII: 4UL6DF56YQ)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)
<b>ANNATTO</b> (UNII: 6PQP1V1B6O)
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>TRICALCIUM PHOSPHATE</b> (UNII: K4C08XP666)
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)
<b>TALC</b> (UNII: 7SEV7J4R1U)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NHRIC:67555-131-30	30 in 1 BOTTLE		
2	NHRIC:67555-131-90	90 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
dietary supplement		06/01/2024	

### Supplement Facts

**Serving Size :**

**Serving per Container :**

	Amount Per Serving	% Daily Value
color		
shape		
size (solid drugs)	21 mm	
scoring	3	
imprint		
flavor		

**Labeler** - Pronova Corporation (111421496)