

FAMILY WELLNESS EFFERVESCENT ANTACID AND PAIN RELIEF- aspirin, citric acid, sodium bicarbonate tablet, effervescent
Family Wellness Services, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Family Wellness Effervescent Antacid & Pain Relief

Purpose

Aspirin 325 mg (NSAID)*.....Pain reliever

Citric acid 1000 mg Antacid

Sodium bicarbonate (heat-treated) 1916 mg Antacid

*Nonsteroidal anti-inflammatory drug

Active ingredients

Aspirin 325 mg (NSAID)*

Citric acid 1000 mg

Sodium bicarbonate (heat-treated) 1916 mg

*Nonsteroidal anti-inflammatory drug

Uses for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- pain alone (headaches or body aches and pains)

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin ■ if overly full from food or drink. Ask a doctor if severe stomach pain

occurs after taking this product.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 10 days

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose

get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
- adults and children 12 years and over (up to 60 years of age): 2 tablets every 4 hours - do not exceed 8 tablets in 24 hours

- 60 years and over: 2 tablets every 4 hours - do not exceed 4 tablets in 24 hours

Other information

- **each tablet contains:** sodium 524 mg
- store at room temperature (59°-86°F)

Inactive ingredients

- none

Serious side effects associated with the use of this product can be reported to the following address:

Distributed by: Midwood Brands, LLC

10611 Monroe Rd, Matthews, NC 28105

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



Drug Facts (continued)

or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ if you are allergic to aspirin ■ if overly full from food or drink ■ Ask a doctor if severe stomach pain occurs after taking this product

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 10 days

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or loss of hearing occurs

It is especially important not to use aspirin during the last 3 months of pregnancy or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy or breast-feeding, ask a health professional before use.

Drug Facts

Active ingredients

Aspirin 325 mg (NSAID)
Citric acid 1000 mg
Sodium bicarbonate 1916 mg

Pain Reliever

Purpose

Uses for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- pain alone (headaches or body aches and pains)

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- facial swelling
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription

COMPARE TO ALKA-SELTZER®

FAMILY wellness™

EFFERVESCENT

Antacid & Pain Relief

For Relief of:

- Acid Indigestion
- Heartburn with Headache
- Body Aches

24 TABLETS



ORIGINAL FLAVOR



COMPARE TO ALKA-SELTZER®

FAMILY wellness™

EFFERVESCENT

Antacid & Pain Relief

Aspirin (NSAID)/Pain Reliever
Citric Acid/Antacid
Sodium Bicarbonate/Antacid

For Relief of:

- Acid Indigestion
- Heartburn with Headache
- Body Aches

24 TABLETS



ORIGINAL FLAVOR



Drug Facts (continued)

pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
- adults and children 12 years and over (up to 60 yrs of age): 2 tablets every 4 hours - do not exceed 8 tablets in 24 hrs
- 60 years and over: 2 tablets every 4 hours - do not exceed 4 tablets in 24 hrs

Other information

- each tablet contains: sodium 524 mg
- store at room temperature (59°-86°F)

Inactive ingredients

- none



Serious side effects associated with the use of this product can be reported to the following address.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
10611 MONROE RD., MATTHEWS, NC 28105

NOT 100% SATISFIED?

Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

*This product is not manufactured or distributed by Bayer Corporation, owner of the registered trademark Alka-Seltzer®.


Manufactured in the USA with domestic and imported ingredients.

SKU 0902755



DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

PN: 42124 FA C1
R: 11/17



FAMILY WELLNESS EFFERVESCENT ANTACID AND PAIN RELIEF

aspirin, citric acid, sodium bicarbonate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-421
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	1916 mg

Product Characteristics				
Color		white	Score	no score
Shape		ROUND	Size	25mm
Flavor			Imprint Code	L421
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-421-36	36 in 1 CARTON	06/13/2017	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:55319-421-24	24 in 1 CARTON	12/16/2017	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final		part343	06/13/2017	

Labeler - Family Wellness Services, Inc. (024472631)

Revised: 5/2019

Family Wellness Services, Inc.