

**MAXIMUM STRENGTH ANTACID AND ANTIGAS- aluminum hydroxide,  
magnesium hydroxide, dimethicone liquid  
RITE AID PHARMACY**

-----  
**riteaid antacid max cherry**

**Active ingredients (in each 5 mL teaspoonful)**

Aluminum hydroxide 400mg (equivalent to dried gel, USP)  
Magnesium hydroxide 400 mg  
Simethicone 40mg

**Purposes**

Antacid

Antigas

**Uses**

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

**Warnings**

**Do not take** more than 12 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug.

Antacids may interact with certain prescription drugs.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Directions**

- shake well before each use
- adults and children 12 years of age and older: take 2 to 4 teaspoonfuls two times a

day or as directed by a doctor

- children under 12 years: ask a doctor

### **Other information**

- **each 5 mL teaspoonful contains:** magnesium 165 mg, sodium 5 mg
- do not freeze
- store at room temperature tightly closed

### **Inactive ingredients**

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

### **Questions or comments?**

**1-800-540-3765**

**package Label**

NDC 11822-5742-0

# MAXIMUM STRENGTH ANTACID & ANTIGAS

ALUMINUM HYDROXIDE 400 mg  
MAGNESIUM HYDROXIDE 400 mg  
SIMETHICONE 40 mg

Fast relief of:  
heartburn • pressure  
acid indigestion  
bloating

218-11322-1 REV 1222



CHERRY  
FLAVOR

12 FL OZ (355 mL)



## Drug Facts

**TAMPER-EVIDENT:** Do not use if Imprinted neckband is missing or broken.

### Active ingredients

(in each 5 mL teaspoonful)

Active ingredients	Purposes
Aluminum hydroxide 400 mg (equivalent to dried gel USP)	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

**Uses** relieves • heartburn • sour stomach  
• acid indigestion • the symptoms referred to as gas

### Warnings

**Do not take** more than 12 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks.

**Ask a doctor before use if you have**

• kidney disease • a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are** presently taking a prescription drug. Antacids may interact with certain prescriptions drugs.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

### Directions

• shake well before each use  
• adults and children 12 years and older: take 2 to 4 teaspoonfuls two times a day or as directed by a doctor  
• children under 12 years: ask a doctor

### Other information

• each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg  
• store at room temperature • protect from freezing  
• keep tightly closed

**Inactive ingredients** benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

**Questions or comments?** 1-800-540-3765

DISTRIBUTED BY:  
RITE AID  
200 NEWBERRY COMMONS,  
ETTERS, PA 17319  
www.riteaid.com



918-11322-3

REV GC1222



0 11822 57420 4

**SATISFACTION GUARANTEE:**  
If you're not satisfied, we'll  
happily refund your money.

## MAXIMUM STRENGTH ANTACID AND ANTIGAS

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-5742
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	400 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 5 mL
<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>BUTYLPARABEN</b> (UNII: 3QPI1U3FV8)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-5742-0	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2012	08/31/2026

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	12/01/2012	08/31/2026

**Labeler** - RITE AID PHARMACY (014578892)

**Registrant** - GCP Laboratories (965480861)

## Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(11822-5742) , pack(11822-5742)

