

**CHEST CONGESTION RELIEF DM- dextromethorphan hydrobromide,
guaifenesin liquid
RUGBY LABORATORIES**

Rugby- Chest Cong DM 1313

Active ingredient

Dextromethorphan HBr, 10 mg

Guaifenesin 100 mg

Purpose

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation
- helps loosen phlegm (mucus)
- helps thin bronchial secretions to make coughs more productive

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Do not use if you have ever had an allergic reaction to any of the ingredients in this product.

Ask a doctor before use if you have

- a cough with too much phlegm (mucus)
- a cough that lasts or is chronic such as occurs with smoking asthma, chronic bronchitis or emphysema.

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away

Directions

- take every 4 hours as needed, or as directed by a doctor
- do not take more than 6 doses in 24 hours
- do not exceed recommended dose

Children under 12 years

do not use

Inactive ingredients: Cherry flavor, citric acid, FD&C Red #40, menthol, methylparaben, propylene glycol, propylparaben, water, sodium citrate, sucralose, sucrose.

1-800-645-2158

Drug Facts
Active ingredient (per 5mL teaspoonful) Purposes
Dextromethorphan HBr, USP 10 mgCough suppressant
Guaifenesin, USP 100 mgExpectorant
Uses • temporarily relieves cough due to minor throat and bronchial irritation • helps loosen phlegm (mucus) • helps thin bronchial secretions to make coughs more productive
Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Do not use if you have ever had an allergic reaction to any of the ingredients in this product.
Ask a doctor before use if you have
• a cough with too much phlegm (mucus) • a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash, or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Chest Congestion Relief DM**
Oral Solution
Guaifenesin Dextromethorphan Expectorant Cough Suppressant
Cherry Flavored
• Controls Cough
• Loosens and Relieves Chest Congestion
• Alcohol Free
Cough Formula for Adults and Children 12 Years and Over
16 FL OZ (473 mL)
PACKAGE NOT CHILD RESISTANT

NDC 0536-1313-85
Compare to the active ingredients in Robitussin® DM*

Drug Facts (continue)
Directions
• take every 4 hours as needed, or as directed by a doctor
• do not take more than 6 doses in 24 hours
• do not exceed recommended dose
Adults and children 12 years and over 2 teaspoonfuls (10 mL)
Children under 12 years do not use
Other information
• each teaspoonful (5 mL) contains: **sodium 2 mg**
• store at room temperature 15°- 30°C (59°-86°F)
• protect from freezing • do not refrigerate • protect from light
• Pharmacist-Preserve and dispense in a tight, light-resistant container with a child resistant cap as defined in the USP
• **TAMPER-EVIDENT:** Do not use if foil seal over bottle opening is torn broken or missing.
Inactive ingredients: cherry flavor, citric acid, FD&C Red #40, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose, sucrose.
Questions or comments? 1-800-645-2158
*This product is not manufactured or distributed by the owner of the registered trademark Robitussin® DM.
THIS IS A BULK CONTAINER NOT INTENDED FOR DISPENSING.
Distributed by:
RUGBY® LABORATORIES
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
www.rugbylaboratories.com
Code# L73
Lot#
Exp. Date:

Rev. 08/20 R-164 Re-order No. 371050

3 05361 31385 2
REV0820

CHEST CONGESTION RELIEF DM

dextromethorphan hydrobromide, guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-1313
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MENTHOL (UNII: L7T10EIP3A)	

METHYLPARABEN (UNII: A2I8C7HI9T)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
SODIUM CITRATE (UNII: 1Q73Q2JULR)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
SUCROSE (UNII: C151H8M554)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1313-85	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2020	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	10/01/2020	

Labeler - RUGBY LABORATORIES (079246066)