

**NEO-SOL 50- neomycin sulfate powder, for solution**  
**Huvepharma, Inc.**

-----  
**Neo-Sol® 50**

**100 Gram Packet Front Panel**

**Neo-Sol® 50**  
**(neomycin sulfate soluble powder)**

**Antibacterial**

**For Oral Use in Cattle, Swine,  
Sheep, Goats and Turkeys**

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Each Packet Contains:  
71.5 gm neomycin sulfate  
equivalent to 50 gm neomycin.

Store between 20°C - 25°C  
(68°F - 77°F) with excursions  
permitted between 15°C - 40°C  
(59°F - 104°F).

Restricted Drug (California) - Use only as directed.

Not for Use in Humans.

Keep Out of Reach of Children

Net wt  
3.5 oz (100 gm)



Approved by FDA under ANADA # 200-130

P08-90074BF (Rev 04-2021)

Huvepharma and Neo-Sol 50 are registered trademarks of Huvepharma EOOD

Huvepharma®

Manufactured for:

Huvepharma, Inc.

525 Westpark Dr., Suite 230

Peachtree City, GA 30269

Add to drinking water - Not for use in liquid supplements.

**Dosage and Administration:**

Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

**Herd/Flock Treatment:**

Each packet will treat 7150 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 7150 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required.

**Fresh medicated water should be prepared each day.**

**Individual Animal Treatment:** To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12 - 24 hours.

**Drinking Water:** Swine - Use the number of packets indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs weighing 25 to 50 lbs..... 2 packets

Pigs weighing 50 to 100 lbs... 3 packets  
Pigs weighing over 100 lbs.... 4 packets

### **Precautions:**

To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If

symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated

as the sole treatment since systemic levels of neomycin are not obtained due to low absorption

from the gastrointestinal tract.

**Important:** Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep, goats, and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the

Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or [www.huvepharma.us](http://www.huvepharma.us). For

additional information about adverse drug experience reporting for animal drugs, contact FDA

at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

### **RESIDUE WARNINGS:**

Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:

Turkeys ..... 0 days

Cattle ..... 1 day

Sheep ..... 2 days

Swine and goats ..... 3 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

### **2000 Gram Packet Front Panel**

**Neo-Sol® 50**

**(neomycin sulfate soluble powder)**

**Antibacterial**

**For Oral Use in Cattle, Swine,  
Sheep, Goats and Turkeys**

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli*

susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Each Packet Contains:

1430 gm neomycin sulfate equivalent to 1000 gm neomycin.

Store between 20°C - 25°C  
(68°F - 77°F) with excursions  
permitted between 15°C - 40°C  
(59°F - 104°F).

Restricted Drug (California) - Use only as directed.

**Not for Use in Humans.**

**Keep Out of Reach of Children.**

**Net wt**

**70.5 oz (2000 gm)**

Take Time



**Observe Label  
Directions**

Approved by FDA under ANADA # 200-130

P08-23369BF (04-2021)

Huvepharma and Neo-Sol 50 are registered trademarks of Huvepharma EOOD

Huvepharma

Manufactured for:

Huvepharma, Inc.  
525 Westpark Dr., Suite 230  
Peachtree City, GA 30269

Add to drinking water - Not for use in liquid supplements.

**Dosage and Administration:**

Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

**Herd/Flock Treatment:**

Each packet will treat 143,000 pounds body weight. Therefore, estimate the total number

of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 143,000 pounds. The product should be added to the amount

of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required.

**Fresh medicated water should be prepared each day.**

**Individual Animal Treatment:** To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12 - 24 hours.

**Drinking Water: Swine** - Use the quantity indicated below in 5120 gallons of water, or in 40 gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs weighing 25 to 50 lbs..... 2 packets

Pigs weighing 50 to 100 lbs... 3 packets

Pigs weighing over 100 lbs.... 4 packets

**Precautions:**

To administer the stated dosage, the concentration of neomycin required in medicated water

must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects

water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult

a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

**Important:** Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep, goats, and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or [www.huvepharma.us](http://www.huvepharma.us). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

### **RESIDUE WARNINGS:**

Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:

Turkeys .....	0 days
Cattle .....	1 day
Sheep .....	2 days
Swine and goats .....	3 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

### **500 Gram Packet Front Panel**

**Neo-Sol® 50**  
**(neomycin sulfate soluble powder)**

**Antibacterial**

**For Oral Use in Cattle, Swine,  
Sheep, Goats and Turkeys**

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.  
For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Each Packet Contains:  
357.5 gm neomycin sulfate equivalent to 250 gm neomycin.

Store between 20°C - 25°C  
(68°F - 77°F) with excursions  
permitted between 15°C - 40°C  
(59°F - 104°F).

Restricted Drug (California) - Use only as directed.

**Not for Use in Humans.**

**Keep Out of Reach of Children.**

**Net wt**  
**17.6 oz (500 g)**



Approved by FDA under ANADA # 200-130

KLA-N00001 (Rev. 04-2021)

Huvepharma and Neo-Sol 50 are registered trademarks of Huvepharma EOOD

Huvepharma®  
Manufactured for:  
Huvepharma, Inc.  
525 Westpark Dr., Suite 230  
Peachtree City, GA 30269

Add to drinking water - Not for use in liquid supplements

**Dosage and Administration:**

Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

**Herd/Flock Treatment:**

Each packet will treat 35,750 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 35,750 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required.

**Fresh medicated water should be prepared each day.**

**Individual Animal Treatment:** To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

**Drinking Water:** Swine - Use the quantity indicated below in 1280 gallons of water, or in 10 gallons of stock solution used in proportioner set to meter one ounce per gallon.

- Pigs weighing 25 to 50 lbs ..... 2 packets
- Pigs weighing 50 to 100 lbs ..... 3 packets
- Pigs weighing over 100 lbs ..... 4 packets

**Precautions:** To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

**Important:** Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep, goats, and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.



To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or [www.huvepharma.us](http://www.huvepharma.us). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

**RESIDUE WARNINGS:**

Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:

- Turkeys ..... 0 days
- Cattle ..... 1 day
- Sheep ..... 2 days
- Swine and goats ..... 3 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

**100 GRAM Packet Label Image**

**Drinking Water: Swine** - Use the number of packets indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs weighing 25 to 50 lbs..... 2 packets  
Pigs weighing 50 to 100 lbs.... 3 packets  
Pigs weighing over 100 lbs.... 4 packets

**Precautions:** To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

**Important:** Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep, goats, and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

**To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4663 or [www.huvepharma.us](http://www.huvepharma.us). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimal>.**

**Residue Warnings:** Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:

Turkeys..... 0 days  
Cattle..... 1 day  
Sheep..... 2 days  
Swine and goats..... 3 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

# Neo-Sol<sup>®</sup> 50

(neomycin sulfate soluble powder)

Antibacterial

For Oral Use in Cattle, Swine,  
Sheep, Goats and Turkeys

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.



Manufactured for:  
Huvepharma, Inc.  
525 Westpark Drive, Suite 230  
Peachtree City, GA 30269

Each Packet Contains:  
71.5 gm neomycin sulfate  
equivalent to 50 gm neomycin.

Store between 20°C - 25°C  
(68°F - 77°F) with excursions  
permitted between 15°C - 40°C  
(59°F - 104°F).

Restricted Drug (California) - Use only  
as directed.

Not for Use in Humans.

Keep Out of Reach of Children.

Net wt  
3.5 oz (100 gm)

Take Time



Observe Label  
Directions

Approved by FDA under ANADA # 200-130

P08-90074BF (Rev. 04-2021)

Huvepharma and Neo-Sol 50 are registered  
trademarks of Huvepharma EOOD

Add to drinking water - Not for use in liquid supplements.

**Dosage and Administration:**  
Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.  
Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

**Herd/Flock Treatment:** Each packet will treat 7150 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 7150 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required.

Fresh medicated water should be prepared each day.

**Individual Animal Treatment:** To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk to reach 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

Neo-Sol<sup>®</sup> 50  
(neomycin sulfate soluble powder)

**Drinking Water: Swine** - Use the quantity indicated below in 5120 gallons of water, or in 40 gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs weighing 25 to 50 lbs.... 2 packets  
Pigs weighing 50 to 100 lbs... 3 packets  
Pigs weighing over 100 lbs.... 4 packets

**Precautions:** To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

**Important:** Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep, goats, and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

**To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-894-4883 or www.huvepharma.us. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportnrxdea>.**

**Residue Warnings:** Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:

Turkeys..... 0 days  
Cattle..... 1 day  
Sheep..... 2 days  
Swine and goats..... 3 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

# Neo-Sol<sup>®</sup> 50

(neomycin sulfate soluble powder)

**Antibacterial**

**For Oral Use in Cattle, Swine,  
Sheep, Goats and Turkeys**

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Each Packet Contains:  
1430 gm neomycin sulfate equivalent to  
1000 gm neomycin.

Store between 20°C - 25°C  
(68°F - 77°F) with excursions permitted  
between 15°C - 40°C (59°F - 104°F).

Restricted Drug (California) - Use only  
as directed.

**Not for Use in Humans.**

**Keep Out of Reach of Children.**

**Net wt  
70.5 oz (2000 gm)**



Approved by FDA under ANADA # 200-130

P08-23369BF (Rev. 04-2021)

Huvepharma and Neo-Sol 50 are registered  
trademarks of Huvepharma. EOOD



Manufactured for:  
Huvepharma, Inc.  
525 Westpark Drive, Suite 230  
Peachtree City, GA 30269

## Neo-Sol<sup>®</sup> 50

(neomycin sulfate soluble powder)

Add to drinking water - Not for use in liquid supplements.

**Dosage and Administration:**

Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.  
Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

**Herd/Flock Treatment:** Each packet will treat 143,000 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 143,000 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required.

Fresh medicated water should be prepared each day.

**Individual Animal Treatment:** To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk to reach 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

**Drinking Water: Swine** - Use the quantity indicated below in 1280 gallons of water, or in 10 gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs weighing 25 to 50 lbs..... 2 packets  
Pigs weighing 50 to 100 lbs.... 3 packets  
Pigs weighing over 100 lbs.... 4 packets

**Precautions:** To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

**Important:** Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep, goats, and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

**To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-984-4683 or [www.huvepharma.us](http://www.huvepharma.us). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimal>.**

**Residue Warnings:** Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:

Turkeys..... 0 days  
Cattle..... 1 day  
Sheep..... 2 days  
Swine and goats..... 3 days

A withdrawal period has not been established for this product in pre-milking calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

# Neo-Sol® 50

(neomycin sulfate soluble powder)

## Antibacterial

For Oral Use in Cattle, Swine,  
Sheep, Goats and Turkeys

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Each Packet Contains:  
357.5 gm neomycin sulfate equivalent to  
250 gm neomycin.

Store between 20°C - 25°C  
(68°F - 77°F) with excursions permitted  
between 15°C - 40°C (59°F - 104°F).

Restricted Drug (California) - Use only  
as directed.

**Not for Use in Humans.**

**Keep Out of Reach of Children.**

Net wt  
17.6 oz (500 gm)

Take Time



Observe Label  
Directions



Manufactured for:  
Huvepharma, Inc.  
325 Westpark Drive, Suite 230  
Peachtree City, GA 30269

Approved by FDA under ANADA # 200-130

KLA-N00001 (Rev. 04-2021)

Huvepharma and Neo-Sol 50 are registered  
trademarks of Huvepharma, EOOD

# Neo-Sol® 50

(neomycin sulfate soluble powder)

Add to drinking water - Not for use in liquid supplements:

**Dosage and Administration:**  
Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days.

Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

**Herd/Flock Treatment:** Each packet will treat 35,750 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 35,750 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required.

Fresh medicated water should be prepared each day.

**Individual Animal Treatment:** To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

neomycin sulfate powder, for solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:23243-6755
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	0.715 g in 1 g

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-6755-1	75 in 1 PAIL		
1		100 g in 1 PACKET		
2	NDC:23243-6755-2	10 in 1 CARTON		
2		2000 g in 1 PACKET		
3	NDC:23243-6755-3	20 in 1 CARTON		
3		500 g in 1 PACKET		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANADA	ANADA200130	04/13/2017	

**Labeler** - Huvepharma, Inc. (619153559)

**Registrant** - Huvepharma EOOD (552671651)

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Huvepharma, Inc - St. Louis		883128204	manufacture, analysis, pack, label

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Shandong Qilu King-phar Pharmaceutical Co., Ltd.		421524323	api manufacture

Revised: 6/2023

Huvepharma, Inc.