

**MAXIMUM STRENGTH ANTACID- aluminum hydroxide, magnesium hydroxide, dimethicone suspension**

**Chain Drug Marketing Association**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**QC ant max**

**Active ingredients (in each 5 mL teaspoonful)**

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)

Magnesium hydroxide 400 mg

Simethicone 40mg

**Purposes**

Antacid

Antigas

**Uses**

relieves

- heartburn
- sour stomach
- acid indigestion
- upset stomach due to these symptoms
- the symptoms referred to as gas

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug.

Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if** symptoms last more than 2 weeks

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep this and all drugs out of the reach of children.**

**Directions**

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor

- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

**Other information**

- **each 5 mL teaspoonful contains:** magnesium 165 mg, sodium 5 mg
- store at room temperature and avoid freezing
- close cap tightly after use

**Inactive ingredients**

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

**Questions or comments?**

**1-800-540-3765**

**package Label**

NDC 63868-715-57



\*Compare to the  
Active Ingredients  
in Mylanta®  
Maximum Strength

# Maximum Strength Antacid

## Antigas

Alumina, Magnesia, and  
Simethicone  
Oral Suspension USP

Alcohol: 0.5%

Fast Soothing Relief of:  
Acid Indigestion  
Heartburn  
Sour Stomach  
Pressure & Bloating



Original Flavor

12 FL OZ (355 mL)

REV GC-0618  
231-06122-2

### Drug Facts

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

#### Active ingredients (in each 5 mL teaspoonful) Purposes

Aluminum hydroxide (equivalent to dried gel, USP) 400 mg ..... Antacid  
Magnesium hydroxide 400 mg ..... Antacid  
Simethicone 40 mg ..... Antigas

**Uses** relieves: • heartburn • sour stomach • acid indigestion  
• upset stomach due to these symptoms  
• the symptoms referred to as gas

#### Warnings

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Antacids may interact with certain prescription drugs.

**Stop use and ask a doctor if symptoms last for more than 2 weeks if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

#### Directions

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than two weeks
- children under 12 years: ask a doctor

#### Other information

- each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg
- store at room temperature and avoid freezing
- close cap tightly after use

**Inactive ingredients** benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

**Questions or comments? 1-800-540-3765**

\*This product is not manufactured or distributed by Johnson & Johnson Merck, distributor of Mylanta®.



REV GC-0618  
931-06122-1



6 35515 99921 3

Distributed by C.D.M.A., Inc.®  
43157 W. Nine Mile  
Novi, MI 48375-0995  
www.qualitychoice.com  
Questions: 248-449-9300

## MAXIMUM STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-715
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDRO XIDE	400 mg in 5 mL

<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>BUTYLPARABEN</b> (UNII: 3QP1U3FV8)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-715-57	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2014	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	03/01/2014	

**Labeler** - Chain Drug Marketing Association (011920774)

**Registrant** - GCP Laboratories (965480861)

### Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(63868-715)