

MOTION SICKNESS- dimenhydrinate tablet
Strategic Sourcing Services, LLC (Health Mart)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Health Mart 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-800-426-9391

Principal display panel

Health Mart®

NDC 62011-0341-1

Compare to Dramamine® Original Formula active ingredient*

ORIGINAL FORMULA

Motion Sickness

Dimenhydrinate 50 mg
Antiemetic

Prevents Nausea, Vomiting & Dizziness for Children & Adults

Actual Size

12 Tablets

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Medtech Products

Inc., owner of the registered trademark Dramamine® Original Formula.
50844 REV0518B19802

McKESSON

Another Quality Product

Distributed by McKesson

One Post Street, San Francisco, CA 94104

Money Back Guarantee

healthmart.com



HealthMart.



NDC 62011-0341-1

Compare to Dramamine® Original Formula active ingredient*

ORIGINAL FORMULA

Motion Sickness

Dimenhydrinate 50 mg
Antiemetic

12 Tablets

Prevents Nausea, Vomiting & Dizziness
for Children & Adults



HealthMart.

NDC 62011-0341-1

Compare to Dramamine® Original Formula active ingredient*

ORIGINAL FORMULA

Motion Sickness

Dimenhydrinate 50 mg
Antiemetic

Actual Size

12 Tablets

Prevents Nausea, Vomiting & Dizziness
for Children & Adults

MCKESSON
Another Quality Product
Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
healthmart.com

*This product is not manufactured or distributed by Medtech Products
Inc., owner of the registered trademark Dramamine® Original Formula.
S0844 REV0518A19802

B-1241-198-02-R
REV0518A19802



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF PULSTER UNIT IS TORN, BROKEN OR
SHOWS ANY SIGNS OF TAMPERING

Active ingredient (in each tablet) Purpose
Dimenhydrinate 50 mg. Antiemetic

Uses for prevention and treatment of these symptoms associated with motion sickness:
nausea ■ vomiting ■ dizziness


Warnings
Do not use for children under 2 years of age unless directed by a doctor.
Ask a doctor before use if you have
■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma
■ difficulty in urination due to enlargement of the prostate gland
■ a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
When using this product
■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur ■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.


Directions
■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity
adults and children 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours;
12 years and over or as directed by a doctor
children 6 to under 12 years ½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours;
or as directed by a doctor
children 2 to under 6 years ½ tablet every 6-8 hours; do not exceed 1 ½ tablets in 24 hours;
or as directed by a doctor

Other information
■ each tablet contains: calcium 35 mg
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR PULSTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture ■ see end flap for expiration date and lot number

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments? 1-800-426-9391





Healthmart 44-198

MOTION SICKNESS

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0341
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, CHLORTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0341-1	2 in 1 CARTON	12/01/1992	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part336	12/01/1992	

Labeler - Strategic Sourcing Services, LLC (Health Mart) (116956644)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(62011-0341)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(62011-0341)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(62011-0341)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(62011-0341)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(62011-0341)

Revised: 1/2020

Strategic Sourcing Services, LLC (Health Mart)