

**PAIN RELIEF EXTRA STRENGTH- acetaminophen tablet, coated**  
**Chain Drug Marketing Association**

-----  
**QCH - 1004 150CT - 2019-1007**

***Drug Facts***

**Active ingredient (in each caplet)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease.

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

### Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> <li>• take 2 caplets every 6 hours while symptoms last</li> <li>• do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>• do not use for more than 10 days unless directed by a doctor</li> </ul>
children under 12 years	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>

### Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

### Inactive ingredients

hypromellose, mineral oil, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

### PRINCIPAL DISPLAY PANEL

NDC 63868-088-02

QUALITY CHOICE

†Compare to TYLENOL(R) Extra Strength Caplets active ingredient

Extra Strength

Pain Relief

Pain Reliever / Fever Reducer

Acetaminophen

100 Caplets - 500 mg each

TWIN PACK 2



**Drug Facts**  
**Active ingredient**  
(in each caplet)  
Acetaminophen 500 mg. Pain reliever/fever reducer.

**Purpose**  
Temporarily relieves minor aches and pains due to:  
■ the common cold ■ headache  
■ backache ■ minor pain of arthritis  
■ toothache  
■ temporarily reduces fever

**Uses**  
Temporarily relieves minor aches and pains due to:  
■ the common cold ■ headache  
■ backache ■ minor pain of arthritis  
■ toothache  
■ temporarily reduces fever

**Warnings**  
Liver warnings: This product contains acetaminophen. The maximum daily dose of this product is 6 caplets (3,000 mg) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have liver disease.**  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**Stop use and ask a doctor if**  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ new symptoms occur  
■ redness or swelling is present  
These could be signs of a serious condition.

**Drug Facts (continued)**  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children.  
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions**  
■ do not take more than directed (see overdose warning)  
adults and children 12 years and over  
■ take 2 caplets every 6 hours while symptoms last  
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor  
do not use for more than 10 days unless directed by a doctor  
ask a doctor under 12 years

**Other Information**  
■ store between 20-25°C (68-77°F)  
■ retain carton for complete product information

**Inactive ingredients**  
hypromellose, mineral oil, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

NC

NC

NC

TWIN PACK 2

PACK 2

INK AND COATING FREE FOR LOT AND EXPIRATION STAMPING



DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING



This product Medication, d



## PAIN RELIEF EXTRA STRENGTH

acetaminophen tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-088
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm

<b>Flavor</b>		<b>Imprint Code</b>	M2A4;57344	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-088-02	2 in 1 CARTON	08/27/2012	04/30/2018
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63868-088-15	150 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/27/2012	05/31/2026
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	08/27/2012	05/31/2026	

**Labeler** - Chain Drug Marketing Association (011920774)

Revised: 10/2024

Chain Drug Marketing Association