

# QUETIAPINE- quetiapine tablet

## Solco Healthcare US, LLC

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use QUETIAPINE TABLETS safely and effectively. See full prescribing information for QUETIAPINE TABLETS.

QUETIAPINE tablets, for oral use  
Initial U.S. Approval: 1997

#### **WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS; and SUICIDAL THOUGHTS AND BEHAVIORS**

*See full prescribing information for complete boxed warning.*

##### **Increased Mortality in Elderly Patients with Dementia-Related Psychosis**

- **Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Quetiapine tablets are not approved for elderly patients with dementia-related psychosis (5.1)**

##### **Suicidal Thoughts and Behaviors**

- **Increased risk of suicidal thoughts and behavior in children, adolescents and young adults taking antidepressants (5.2)**
- **Monitor for worsening and emergence of suicidal thoughts and behaviors (5.2)**

### RECENT MAJOR CHANGES

Warnings and Precautions (5.15) 1/2025

### INDICATIONS AND USAGE

Quetiapine tablets are atypical antipsychotic indicated for the treatment of:

- Schizophrenia (1.1)
- Bipolar I disorder manic episodes (1.2)
- Bipolar disorder, depressive episodes (1.2)

### DOSAGE AND ADMINISTRATION

- Quetiapine tablets can be taken with or without food (2.1)

Indication	Initial Dose	Recommended Dose	Maximum Dose
Schizophrenia - Adults (2.2)	25 mg twice daily	150-750 mg/day	750 mg/day
Schizophrenia - Adolescents (13-17 years) (2.2)	25 mg twice daily	400-800 mg/day	800 mg/day
Bipolar Mania - Adults Monotherapy or as an adjunct to lithium or divalproex (2.2)	50 mg twice daily	400-800 mg/day	800 mg/day
Bipolar Mania - Children and Adolescents (10-17 years), Monotherapy (2.2)	25 mg twice daily	400-600 mg/day	600 mg/day
Bipolar Depression - Adults (2.2)	50 mg once daily at bedtime	300 mg/day	300 mg/day

- *Geriatric Use:* Consider a lower starting dose (50 mg/day), slower titration and careful monitoring during the initial dosing period in the elderly (2.3, 8.5)
- *Hepatic Impairment:* Lower starting dose (25 mg/day) and slower titration may be needed (2.4, 8.7, 12.3)

### DOSAGE FORMS AND STRENGTHS

Tablets: 25 mg, 50 mg, 100 mg, 200 mg, 300 mg, and 400 mg (3)

### CONTRAINDICATIONS

Known hypersensitivity to quetiapine tablets or any components in the formulation. (4)

---

## WARNINGS AND PRECAUTIONS

---

- **Cerebrovascular Adverse Reactions:** Increased incidence of cerebrovascular adverse reactions (e.g., stroke, transient ischemic attack) has been seen in elderly patients with dementia-related psychoses treated with atypical antipsychotic drugs (5.3)
- **Neuroleptic Malignant Syndrome (NMS):** Manage with immediate discontinuation and close monitoring (5.4)
- **Metabolic Changes:** Atypical antipsychotics have been associated with metabolic changes. These metabolic changes include hyperglycemia, dyslipidemia, and weight gain (5.5)
- **Hyperglycemia and Diabetes Mellitus:** Monitor patients for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Monitor glucose regularly in patients with diabetes or at risk for diabetes
- **Dyslipidemia:** Undesirable alterations have been observed in patients treated with atypical antipsychotics. Appropriate clinical monitoring is recommended, including fasting blood lipid testing at the beginning of, and periodically, during treatment
- **Weight Gain:** Gain in body weight has been observed; clinical monitoring of weight is recommended
- **Tardive Dyskinesia:** Discontinue if clinically appropriate (5.6)
- **Hypotension:** Use with caution in patients with known cardiovascular or cerebrovascular disease (5.7)
- **Increased Blood Pressure in Children and Adolescents:** Monitor blood pressure at the beginning of, and periodically during treatment in children and adolescents (5.9)
- **Leukopenia, Neutropenia and Agranulocytosis:** Monitor complete blood count frequently during the first few months of treatment in patients with a pre-existing low white cell count or a history of leukopenia/neutropenia and discontinue quetiapine tablets at the first sign of a decline in WBC in absence of other causative factors (5.10)
- **Cataracts:** Lens changes have been observed in patients during long-term quetiapine treatment. Lens examination is recommended when starting treatment and at 6-month intervals during chronic treatment (5.11)
- **Anticholinergic (antimuscarinic) Effects:** Use with caution with other anticholinergic drugs and in patients with urinary retention, prostatic hypertrophy, or constipation (5.20)

---

## ADVERSE REACTIONS

---

Most common adverse reactions (incidence  $\geq$ 5% and twice placebo):

- Adults: somnolence, dry mouth, dizziness, constipation, asthenia, abdominal pain, postural hypotension, pharyngitis, weight gain, lethargy, ALT increased, dyspepsia (6.1)
- Children and Adolescents: somnolence, dizziness, fatigue, increased appetite, nausea, vomiting, dry mouth, tachycardia, weight increased (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Solco Healthcare US, LLC at 1-866-257-2597 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

---

## DRUG INTERACTIONS

---

- **Concomitant use of strong CYP3A4 inhibitors:** Reduce quetiapine dose to one sixth when coadministered with strong CYP3A4 inhibitors (e.g., ketoconazole, ritonavir) (2.5, 7.1, 12.3)
- **Concomitant use of strong CYP3A4 inducers:** Increase quetiapine dose up to 5 fold when used in combination with a chronic treatment (more than 7-14 days) of potent CYP3A4 inducers (e.g., phenytoin, rifampin, St. John's wort) (2.6, 7.1, 12.3)
- **Discontinuation of strong CYP3A4 inducers:** Reduce quetiapine dose by 5-fold within 7-14 days of discontinuation of CYP3A4 inducers (2.6, 7.1, 12.3)

---

## USE IN SPECIFIC POPULATIONS

---

- **Pregnancy:** May cause extrapyramidal and/or withdrawal symptoms in neonates with third trimester exposure. (8.1)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 7/2025**

---

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS; and SUICIDAL THOUGHTS AND BEHAVIORS

#### 1 INDICATIONS AND USAGE

- 1.1 Schizophrenia
- 1.2 Bipolar Disorder
- 1.3 Special Considerations in Treating Pediatric Schizophrenia and Bipolar I Disorder

## **2 DOSAGE AND ADMINISTRATION**

- 2.1 Important Administration Instructions
- 2.2 Recommended Dosing
- 2.3 Dose Modifications in Elderly Patients
- 2.4 Dose Modifications in Hepatically Impaired Patients
- 2.5 Dose Modifications when used with CYP3A4 Inhibitors
- 2.6 Dose Modifications when used with CYP3A4 Inducers
- 2.7 Re-initiation of Treatment in Patients Previously Discontinued
- 2.8 Switching from Antipsychotics

## **3 DOSAGE FORMS AND STRENGTHS**

## **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

- 5.1 Increased Mortality in Elderly Patients with Dementia-Related Psychosis
- 5.2 Suicidal Thoughts and Behaviors in Adolescents and Young Adults
- 5.3 Cerebrovascular Adverse Reactions, Including Stroke, in Elderly Patients with Dementia-Related Psychosis
- 5.4 Neuroleptic Malignant Syndrome (NMS)
- 5.5 Metabolic Changes
- 5.6 Tardive Dyskinesia
- 5.7 Hypotension
- 5.8 Falls
- 5.9 Increases in Blood Pressure (Children and Adolescents)
- 5.10 Leukopenia, Neutropenia, and Agranulocytosis
- 5.11 Cataracts
- 5.12 QT Prolongation
- 5.13 Seizures
- 5.14 Hypothyroidism
- 5.15 Hyperprolactinemia
- 5.16 Potential for Cognitive and Motor Impairment
- 5.17 Body Temperature Regulation
- 5.18 Dysphagia
- 5.19 Discontinuation Syndrome
- 5.20 Anticholinergic (antimuscarinic) Effects

## **6 ADVERSE REACTIONS**

- 6.1 Clinical Study Experience
- 6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

- 7.1 Effect of Other Drugs on Quetiapine
- 7.2 Effect of Quetiapine on Other Drugs

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

## **9 DRUG ABUSE AND DEPENDENCE**

- 9.1 Controlled Substance
- 9.2 Abuse

## **10 OVERDOSAGE**

- 10.1 Human Experience
- 10.2 Management of Overdosage

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

### **14 CLINICAL STUDIES**

14.1 Schizophrenia

14.2 Bipolar Disorder

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **WARNING: INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS; and SUICIDAL THOUGHTS AND BEHAVIORS**

#### **Increased Mortality in Elderly Patients with Dementia-Related Psychosis**

**Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death [see *Warnings and Precautions* (5.1)]. Quetiapine tablets are not approved for the treatment of patients with dementia-related psychosis [see *Warnings and Precautions* (5.1)].**

#### **Suicidal Thoughts and Behaviors**

**Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term studies. These studies did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in patients over age 24; there was a reduction in risk with antidepressant use in patients aged 65 and older [see *Warnings and Precautions* (5.2)].**

**In patients of all ages who are started on antidepressant therapy, monitor closely for worsening, and for emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication with the prescriber [see *Warnings and Precautions* (5.2)].**

**Quetiapine tablets are not approved for use in pediatric patients under ten years of age [see *Use in Specific Populations* (8.4)].**

## **1 INDICATIONS AND USAGE**

### **1.1 Schizophrenia**

Quetiapine tablets are indicated for the treatment of schizophrenia. The efficacy of quetiapine tablets in schizophrenia was established in three 6-week trials in adults and one 6-week trial in adolescents (13-17 years). The effectiveness of quetiapine tablets for the maintenance treatment of schizophrenia has not been systematically evaluated in controlled clinical trials [see *Clinical Studies* (14.1)].

## 1.2 Bipolar Disorder

Quetiapine tablets are indicated for the acute treatment of manic episodes associated with bipolar I disorder, both as monotherapy and as an adjunct to lithium or divalproex. Efficacy was established in two 12-week monotherapy trials in adults, in one 3-week adjunctive trial in adults, and in one 3-week monotherapy trial in pediatric patients (10-17 years) [see *Clinical Studies (14.2)*].

Quetiapine tablets are indicated as monotherapy for the acute treatment of depressive episodes associated with bipolar disorder. Efficacy was established in two 8-week monotherapy trials in adult patients with bipolar I and bipolar II disorder [see *Clinical Studies (14.2)*].

Quetiapine tablets are indicated for the maintenance treatment of bipolar I disorder, as an adjunct to lithium or divalproex. Efficacy was established in two maintenance trials in adults. The effectiveness of quetiapine tablets as monotherapy for the maintenance treatment of bipolar disorder has not been systematically evaluated in controlled clinical trials [see *Clinical Studies (14.2)*].

## 1.3 Special Considerations in Treating Pediatric Schizophrenia and Bipolar I Disorder

Pediatric schizophrenia and bipolar I disorder are serious mental disorders, however, diagnosis can be challenging. For pediatric schizophrenia, symptom profiles can be variable, and for bipolar I disorder, patients may have variable patterns of periodicity of manic or mixed symptoms. It is recommended that medication therapy for pediatric schizophrenia and bipolar I disorder be initiated only after a thorough diagnostic evaluation has been performed and careful consideration given to the risks associated with medication treatment. Medication treatment for both pediatric schizophrenia and bipolar I disorder is indicated as part of a total treatment program that often includes psychological, educational and social interventions.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Important Administration Instructions

Quetiapine tablets can be taken with or without food.

### 2.2 Recommended Dosing

The recommended initial dose, titration, dose range and maximum quetiapine tablets dose for each approved indication is displayed in Table 1. After initial dosing, adjustments can be made upwards or downwards, if necessary, depending upon the clinical response and tolerability of the patient [see *Clinical Studies (14.1 and 14.2)*].

**Table 1: Recommended Dosing for Quetiapine Tablets**

Indication	Initial Dose and Titration	Recommended Dose	Maximum Dose
Schizophrenia - Adults	Day 1: 25 mg twice daily. Increase in increments of 25 mg-50 mg divided two or three times on Days 2 and 3 to range of 300-400 mg by Day 4. Further adjustments can be made in increments of 25-50 mg twice a day, in intervals of not less than 2 days.	150-750 mg/day	750 mg/day
Schizophrenia - Adolescents	Day 1: 25 mg twice daily. Day 2: Twice daily dosing totaling 100 mg.		

(13-17 years)	Day 3: Twice daily dosing totaling 200 mg. Day 4: Twice daily dosing totaling 300 mg. Day 5: Twice daily dosing totaling 400 mg. Further adjustments should be in increments no greater than 100 mg/day within the recommended dose range of 400-800 mg/day. Based on response and tolerability, may be administered three times daily.	400-800 mg/day	800 mg/day
Schizophrenia - Maintenance	Not applicable.	400-800 mg/day	800 mg/day
Bipolar Mania - Adults Monotherapy or as an adjunct to lithium or divalproex	Day 1: Twice daily dosing totaling 100 mg. Day 2: Twice daily dosing totaling 200 mg. Day 3: Twice daily dosing totaling 300 mg. Day 4: Twice daily dosing totaling 400 mg. Further dosage adjustments up to 800 mg/day by Day 6 should be in increments of no greater than 200 mg/day.	400-800 mg/day	800 mg/day
Bipolar Mania - Children and Adolescents (10 to 17 years), Monotherapy	Day 1: 25 mg twice daily. Day 2: Twice daily dosing totaling 100 mg. Day 3: Twice daily dosing totaling 200 mg. Day 4: Twice daily dosing totaling 300 mg. Day 5: Twice daily dosing totaling 400 mg. Further adjustments should be in increments no greater than 100 mg/day within the recommended dose range of 400-600 mg/day. Based on response and tolerability, may be administered three times daily.	400-600 mg/day	600 mg/day
Bipolar Depression - Adults	Administer once daily at bedtime. Day 1: 50 mg Day 2: 100 mg Day 3: 200 mg Day 4: 300 mg	300 mg/day	300 mg/day
Bipolar I Disorder Maintenance Therapy - Adults	Administer twice daily totaling 400-800 mg/day as adjunct to lithium or divalproex. Generally, in the maintenance phase, patients continued on the same dose on which they were stabilized.	400-800 mg/day	800 mg/day

## Maintenance Treatment for Schizophrenia and Bipolar I Disorder

**Maintenance Treatment** - Patients should be periodically reassessed to determine the need for maintenance treatment and the appropriate dose for such treatment [see *Clinical Studies (14.2)*].

### 2.3 Dose Modifications in Elderly Patients

Consideration should be given to a slower rate of dose titration and a lower target dose in the elderly and in patients who are debilitated or who have a predisposition to hypotensive reactions [see *Clinical Pharmacology (12.3)*]. When indicated, dose escalation should be performed with caution in these patients.

Elderly patients should be started on quetiapine tablets 50 mg/day and the dose can be increased in increments of 50 mg/day depending on the clinical response and tolerability

of the individual patient.

## **2.4 Dose Modifications in Hepatically Impaired Patients**

Patients with hepatic impairment should be started on 25 mg/day. The dose should be increased daily in increments of 25 mg/day - 50 mg/day to an effective dose, depending on the clinical response and tolerability of the patient.

## **2.5 Dose Modifications when used with CYP3A4 Inhibitors**

Quetiapine tablets dose should be reduced to one sixth of original dose when co-medicated with a potent CYP3A4 inhibitor (e.g., ketoconazole, itraconazole, indinavir, ritonavir, nefazodone, etc.). When the CYP3A4 inhibitor is discontinued, the dose of quetiapine tablets should be increased by 6-fold[see *Clinical Pharmacology (12.3) and Drug Interactions (7.1)*].

## **2.6 Dose Modifications when used with CYP3A4 Inducers**

Quetiapine tablets dose should be increased up to 5-fold of the original dose when used in combination with a chronic treatment (e.g., greater than 7-14 days) of a potent CYP3A4 inducer (e.g., phenytoin, carbamazepine, rifampin, avasimibe, St. John's wort etc.). The dose should be titrated based on the clinical response and tolerability of the individual patient. When the CYP3A4 inducer is discontinued, the dose of quetiapine tablets should be reduced to the original level within 7-14 days [see *Clinical Pharmacology (12.3) and Drug Interactions (7.1)*].

## **2.7 Re-initiation of Treatment in Patients Previously Discontinued**

Although there are no data to specifically address re-initiation of treatment, it is recommended that when restarting therapy of patients who have been off quetiapine tablets for more than one-week, the initial dosing schedule should be followed. When restarting patients who have been off quetiapine tablets for less than one-week, gradual dose escalation may not be required and the maintenance dose may be re-initiated.

## **2.8 Switching from Antipsychotics**

There are no systematically collected data to specifically address switching patients with schizophrenia from antipsychotics to quetiapine tablets, or concerning concomitant administration with antipsychotics. While immediate discontinuation of the previous antipsychotic treatment may be acceptable for some patients with schizophrenia, more gradual discontinuation may be most appropriate for others. In all cases, the period of overlapping antipsychotic administration should be minimized. When switching patients with schizophrenia from depot antipsychotics, if medically appropriate, initiate quetiapine tablets therapy in place of the next scheduled injection. The need for continuing existing EPS medication should be re-evaluated periodically.

## **3 DOSAGE FORMS AND STRENGTHS**

- 25 mg tablets are pink, round, biconvex, film-coated tablets, debossed with "793" on one side and "S" on the other side.
- 50 mg tablets are white, round, biconvex, film-coated tablets, debossed with "794" on one side and "S" on the other side.
- 100 mg tablets are yellow, round, biconvex, film-coated tablets, debossed with "795" on one side and "S" on the other side.
- 200 mg tablets are white, round, biconvex, film-coated tablets, debossed with "796" on one side and "S" on the other side.
- 300 mg tablets are white, capsule-shaped, biconvex, film-coated tablets, debossed with "797" on one side and "S" on the other side.

400 mg tablets are yellow, capsule-shaped, biconvex, film-coated tablets, debossed with "798" on one side and "S" on the other side.

## **4 CONTRAINDICATIONS**

Hypersensitivity to quetiapine or to any excipients in the quetiapine tablets formulation. Anaphylactic reactions have been reported in patients treated with quetiapine tablets.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Increased Mortality in Elderly Patients with Dementia-Related Psychosis**

Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Analysis of 17 placebo-controlled trials (modal duration of 10 weeks), largely in patients taking atypical antipsychotic drugs, revealed a risk of death in drug-treated patients of between 1.6 to 1.7 times the risk of death in placebo-treated patients. Over the course of a typical 10-week controlled trial, the rate of death in drug-treated patients was about 4.5%, compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. Observational studies suggest that, similar to atypical antipsychotic drugs, treatment with conventional antipsychotic drugs may increase mortality. The extent to which the findings of increased mortality in observational studies may be attributed to the antipsychotic drug as opposed to some characteristic(s) of the patients is not clear. Quetiapine tablets are not approved for the treatment of patients with dementia-related psychosis [see *Boxed Warning*].

### **5.2 Suicidal Thoughts and Behaviors in Adolescents and Young Adults**

Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a long-standing concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment. Pooled analyses of short-term placebo-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18-24) with major depressive disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction with antidepressants compared to placebo in adults aged 65 and older.

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, obsessive-compulsive disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of 9 antidepressant drugs in over 4,400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients. There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger patients for almost all drugs studied. There were differences in absolute risk of suicidality across the different indications, with the highest incidence in MDD. The risk differences (drug vs. placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number of cases of suicidality per 1,000

patients treated) are provided in Table 2.

**Table 2: Drug-Placebo Difference in Number of Cases of Suicidality per 1,000 Patients Treated**

<b>Age Range</b>	<b>Drug-Placebo Difference in Number of Cases of Suicidality per 1,000 Patients Treated</b>
	<b>Increases Compared to Placebo</b>
<18	14 additional cases
18-24	5 additional cases
	<b>Decreases Compared to Placebo</b>
25-64	1 fewer case
≥65	6 fewer cases

No suicides occurred in any of the pediatric trials. There were suicides in the adult trials, but the number was not sufficient to reach any conclusion about drug effect on suicide.

It is unknown whether the suicidality risk extends to longer-term use, i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression.

**All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases.**

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for major depressive disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms.

**Families and caregivers of patients being treated with antidepressants for major depressive disorder or other indications, both psychiatric and non-psychiatric, should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to healthcare providers. Such monitoring should include daily observation by families and caregivers.**

Prescriptions for quetiapine tablets should be written for the smallest quantity of tablets consistent with good patient management, in order to reduce the risk of overdose.

**Screening Patients for Bipolar Disorder:** A major depressive episode may be the initial presentation of bipolar disorder. It is generally believed (though not established in controlled trials) that treating such an episode with an antidepressant alone may increase the likelihood of precipitation of a mixed/manic episode in patients at risk for

bipolar disorder. Whether any of the symptoms described above represent such a conversion is unknown. However, prior to initiating treatment with an antidepressant, including quetiapine tablets, patients with depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression.

### **5.3 Cerebrovascular Adverse Reactions, Including Stroke, in Elderly Patients with Dementia-Related Psychosis**

In placebo-controlled trials with risperidone, aripiprazole, and olanzapine in elderly subjects with dementia, there was a higher incidence of cerebrovascular adverse reactions (cerebrovascular accidents and transient ischemic attacks) including fatalities compared to placebo-treated subjects. Quetiapine tablets are not approved for the treatment of patients with dementia-related psychosis [see also *Boxed Warning and Warnings and Precautions (5.1)*].

### **5.4 Neuroleptic Malignant Syndrome (NMS)**

A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with administration of antipsychotic drugs, including quetiapine tablets. Rare cases of NMS have been reported with quetiapine tablets. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatinine phosphokinase, myoglobinuria (rhabdomyolysis) and acute renal failure.

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to exclude cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system (CNS) pathology.

The management of NMS should include: 1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy; 2) intensive symptomatic treatment and medical monitoring; and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, the potential reintroduction of drug therapy should be carefully considered. The patient should be carefully monitored since recurrences of NMS have been reported.

### **5.5 Metabolic Changes**

Atypical antipsychotic drugs have been associated with metabolic changes that include hyperglycemia/diabetes mellitus, dyslipidemia, and body weight gain. While all of the drugs in the class have been shown to produce some metabolic changes, each drug has its own specific risk profile. In some patients, a worsening of more than one of the metabolic parameters of weight, blood glucose, and lipids was observed in clinical studies. Changes in these metabolic profiles should be managed as clinically appropriate.

#### **Hyperglycemia and Diabetes Mellitus**

Hyperglycemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics, including quetiapine. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an

increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycemia-related adverse reactions is not completely understood. However, epidemiological studies suggest an increased risk of hyperglycemia-related adverse reactions in patients treated with the atypical antipsychotics. Precise risk estimates for hyperglycemia-related adverse reactions in patients treated with atypical antipsychotics are not available.

Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g., obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycemia including polydipsia, polyuria, polyphagia, and weakness. Patients who develop symptoms of hyperglycemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug.

Adults:

**Table 3: Fasting Glucose-Proportion of Patients Shifting to  $\geq 126$  mg/dL in Short-Term ( $\leq 12$  weeks) Placebo-Controlled Studies<sup>1</sup>**

Laboratory Analyte	Category Change (At Least Once) from Baseline	Treatment Arm	N	Patients n (%)
Fasting Glucose	Normal to High (<100 mg/dL to $\geq 126$ mg/dL)	Quetiapine	2,907	71 (2.4%)
		Placebo	1,346	19 (1.4%)
	Borderline to High ( $\geq 100$ mg/dL and <126 mg/dL to $\geq 126$ mg/dL)	Quetiapine	572	67 (11.7%)
		Placebo	279	33 (11.8%)

1. Includes quetiapine tablets and quetiapine extended-release tablets data.

In a 24-week trial (active-controlled, 115 patients treated with quetiapine tablets) designed to evaluate glycemic status with oral glucose tolerance testing of all patients, at Week 24 the incidence of a post-glucose challenge glucose level  $\geq 200$  mg/dL was 1.7% and the incidence of a fasting blood glucose level  $\geq 126$  mg/dL was 2.6%. The mean change in fasting glucose from baseline was 3.2 mg/dL and mean change in 2-hour glucose from baseline was -1.8 mg/dL for quetiapine.

In 2 long-term placebo-controlled randomized withdrawal clinical trials for bipolar I disorder maintenance, mean exposure of 213 days for quetiapine tablets (646 patients) and 152 days for placebo (680 patients), the mean change in glucose from baseline was +5.0 mg/dL for quetiapine tablets and -0.05 mg/dL for placebo. The exposure-adjusted rate of any increased blood glucose level ( $\geq 126$  mg/dL) for patients more than 8 hours since a meal (however, some patients may not have been precluded from calorie intake from fluids during fasting period) was 18.0 per 100 patient years for quetiapine tablets (10.7% of patients; n=556) and 9.5 for placebo per 100 patient years (4.6% of patients; n=581).

*Children and Adolescents:*

In a placebo-controlled quetiapine tablets monotherapy study of adolescent patients (13-17 years of age) with schizophrenia (6 weeks duration), the mean change in fasting glucose levels for quetiapine tablets (n=138) compared to placebo (n=67) was -0.75 mg/dL versus -1.70 mg/dL. In a placebo-controlled quetiapine tablets monotherapy study of children and adolescent patients (10-17 years of age) with bipolar mania (3 weeks duration), the mean change in fasting glucose level for quetiapine tablets (n=170) compared to placebo (n=81) was 3.62 mg/dL versus -1.17 mg/dL. No patient in either study with a baseline normal fasting glucose level (<100 mg/dL) or a baseline borderline fasting glucose level (≥100 mg/dL and <126 mg/dL) had a blood glucose level of ≥126 mg/dL.

In a placebo-controlled quetiapine extended-release tablets monotherapy study (8 weeks duration) of children and adolescent patients (10-17 years of age) with bipolar depression, in which efficacy was not established, the mean change in fasting glucose levels for quetiapine extended-release tablets (n=60) compared to placebo (n=62) was 1.8 mg/dL versus 1.6 mg/dL. In this study, there were no patients in the quetiapine extended-release tablets or placebo-treated groups with a baseline normal fasting glucose level (<100 mg/dL) that had an increase in blood glucose level >126 mg/dL. There was one patient in the quetiapine extended-release tablets group with a baseline borderline fasting glucose level (>100 mg/dL and <126 mg/dL) who had an increase in blood glucose level of >126 mg/dL compared to zero patients in the placebo group.

Dyslipidemia

*Adults:*

Table 4 shows the percentage of adult patients with changes in total cholesterol, triglycerides, LDL-cholesterol, and HDL-cholesterol from baseline by indication in clinical trials with quetiapine tablets.

**Table 4: Percentage of Adult Patients with Shifts in Total Cholesterol, Triglycerides, LDL-Cholesterol, and HDL-Cholesterol from Baseline to Clinically Significant Levels by Indication**

Laboratory Analyte	Indication	Treatment Arm	N	Patients n (%)
Total Cholesterol ≥240 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	137	24 (18%)
		Placebo	92	6 (7%)
	Bipolar Depression <sup>2</sup>	Quetiapine tablets	463	41 (9%)
		Placebo	250	15 (6%)
Triglycerides ≥200 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	120	26 (22%)
		Placebo	70	11 (16%)
	Bipolar Depression <sup>2</sup>	Quetiapine tablets	436	59 (14%)
		Placebo	232	20 (9%)
LDL-Cholesterol ≥160 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	na <sup>3</sup>	na <sup>3</sup>
		Placebo	na <sup>3</sup>	na <sup>3</sup>
	Bipolar Depression <sup>2</sup>	Quetiapine tablets	465	29 (6%)
		Placebo	256	12 (5%)
HDL-Cholesterol ≤40 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	na <sup>3</sup>	na <sup>3</sup>
		Placebo	na <sup>3</sup>	na <sup>3</sup>
	Bipolar Depression <sup>2</sup>	Quetiapine tablets	393	56 (14%)
		Placebo	214	29 (14%)

1. 6 weeks duration

2. 8 weeks duration

3. Parameters not measured in the quetiapine tablets registration studies for

schizophrenia.

*Children and Adolescents:*

Table 5 shows the percentage of children and adolescents with changes in total cholesterol, triglycerides, LDL-cholesterol, and HDL-cholesterol from baseline in clinical trials with quetiapine tablets.

**Table 5: Percentage of Children and Adolescents with Shifts in Total Cholesterol, Triglycerides, LDL-Cholesterol, and HDL-Cholesterol from Baseline to Clinically Significant Levels**

Laboratory Analyte	Indication	Treatment Arm	N	Patients n (%)
Total Cholesterol ≥200 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	107	13 (12%)
		Placebo	56	1 (2%)
	Bipolar Mania <sup>2</sup>	Quetiapine tablets	159	16 (10%)
		Placebo	66	2 (3%)
Triglycerides ≥150 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	103	17 (17%)
		Placebo	51	4 (8%)
	Bipolar Mania <sup>2</sup>	Quetiapine tablets	149	32 (22%)
		Placebo	60	8 (13%)
LDL-Cholesterol ≥130 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	112	4 (4%)
		Placebo	60	1 (2%)
	Bipolar Mania <sup>2</sup>	Quetiapine tablets	169	13 (8%)
		Placebo	74	4 (5%)
HDL-Cholesterol ≤40 mg/dL	Schizophrenia <sup>1</sup>	Quetiapine tablets	104	16 (15%)
		Placebo	54	10 (19%)
	Bipolar Mania <sup>2</sup>	Quetiapine tablets	154	16 (10%)
		Placebo	61	4 (7%)

1. 13-17 years, 6 weeks duration

2. 10-17 years, 3 weeks duration

In a placebo-controlled quetiapine extended-release tablets monotherapy study (8 weeks duration) of children and adolescent patients (10 to 17 years of age) with bipolar depression, in which efficacy was not established, the percentage of children and adolescents with shifts in total cholesterol (≥200 mg/dL), triglycerides (≥150 mg/dL), LDL-cholesterol (≥130 mg/dL) and HDL-cholesterol (≤40 mg/dL) from baseline to clinically significant levels were: total cholesterol 8% (7/83) for quetiapine extended-release tablets vs. 6% (5/84) for placebo; triglycerides 28% (22/80) for quetiapine extended-release tablets vs. 9% (7/82) for placebo; LDL-cholesterol 2% (2/86) for quetiapine extended-release tablets vs. 4% (3/85) for placebo and HDL-cholesterol 20% (13/65) for quetiapine extended-release tablets vs. 15% (11/74) for placebo.

**Weight Gain**

Increases in weight have been observed in clinical trials. Patients receiving quetiapine should receive regular monitoring of weight.

*Adults:* In clinical trials with quetiapine tablets the following increases in weight have been reported.

**Table 6: Proportion of Patients with Weight Gain ≥7% of Body Weight (Adults)**

Vital Sign	Indication	Treatment Arm	N	Patients n (%)
<b>Weight Gain <math>\geq</math>7% of Body Weight</b>	Schizophrenia <sup>1</sup>	Quetiapine tablets	391	89 (23%)
		Placebo	206	11 (6%)
	Bipolar Mania (monotherapy) <sup>2</sup>	Quetiapine tablets	209	44 (21%)
		Placebo	198	13 (7%)
	Bipolar Mania (adjunct therapy) <sup>3</sup>	Quetiapine tablets	196	25 (13%)
		Placebo	203	8 (4%)
	Bipolar Depression <sup>4</sup>	Quetiapine tablets	554	47 (8%)
		Placebo	295	7 (2%)

1. up to 6 weeks duration

2. up to 12 weeks duration

3. up to 3 weeks duration

4. up to 8 weeks duration

#### *Children and Adolescents:*

In two clinical trials with quetiapine tablets, one in bipolar mania and one in schizophrenia, reported increases in weight are included in Table 7.

**Table 7: Proportion of Patients with Weight Gain  $\geq$ 7% of Body Weight (Children and Adolescents)**

Vital Sign	Indication	Treatment Arm	N	Patients n (%)
Weight Gain $\geq$ 7% of Body Weight	Schizophrenia <sup>1</sup>	Quetiapine tablets	111	23 (21%)
		Placebo	44	3 (7%)
	Bipolar Mania <sup>2</sup>	Quetiapine tablets	157	18 (12%)
		Placebo	68	0 (0%)

1. 6 weeks duration

2. 3 weeks duration

The mean change in body weight in the schizophrenia trial was 2.0 kg in the quetiapine tablets group and -0.4 kg in the placebo group and in the bipolar mania trial, it was 1.7 kg in the quetiapine tablets group and 0.4 kg in the placebo group.

In an open-label study that enrolled patients from the above two pediatric trials, 63% of patients (241/380) completed 26 weeks of therapy with quetiapine tablets. After 26 weeks of treatment, the mean increase in body weight was 4.4 kg. Forty-five percent of the patients gained  $\geq$ 7% of their body weight, not adjusted for normal growth. In order to adjust for normal growth over 26 weeks, an increase of at least 0.5 standard deviation from baseline in BMI was used as a measure of a clinically significant change; 18.3% of patients on quetiapine tablets met this criterion after 26 weeks of treatment.

In a clinical trial for quetiapine extended-release tablets in children and adolescents (10-17 years of age) with bipolar depression, in which efficacy was not established, the percentage of patients with weight gain  $\geq$ 7% of body weight at any time was 15% (14/92) for quetiapine extended-release tablets vs. 10% (10/100) for placebo. The mean change in body weight was 1.4 kg in the quetiapine extended-release tablets group vs. 0.6 kg in the placebo group.

When treating pediatric patients with quetiapine tablets for any indication, weight gain should be assessed against that expected for normal growth.

## **5.6 Tardive Dyskinesia**

A syndrome of potentially irreversible, involuntary, dyskinetic movements may develop in patients treated with antipsychotic drugs, including quetiapine. Although the prevalence of the syndrome appears to be highest among the elderly, especially elderly women, it is impossible to rely upon prevalence estimates to predict, at the inception of antipsychotic treatment, which patients are likely to develop the syndrome. Whether antipsychotic drug products differ in their potential to cause tardive dyskinesia is unknown.

The risk of developing tardive dyskinesia and the likelihood that it will become irreversible are believed to increase as the duration of treatment and the total cumulative dose of antipsychotic drugs administered to the patient increase. However, the syndrome can develop, although much less commonly, after relatively brief treatment periods at low doses or may even arise after discontinuation of treatment.

Tardive dyskinesia may remit, partially or completely, if antipsychotic treatment is withdrawn. Antipsychotic treatment, itself, however, may suppress (or partially suppress) the signs and symptoms of the syndrome and thereby may possibly mask the underlying process. The effect that symptomatic suppression has upon the long-term course of the syndrome is unknown.

Given these considerations, quetiapine tablets should be prescribed in a manner that is most likely to minimize the occurrence of tardive dyskinesia. Chronic antipsychotic treatment should generally be reserved for patients who appear to suffer from a chronic illness that (1) is known to respond to antipsychotic drugs, and (2) for whom alternative, equally effective, but potentially less harmful treatments are not available or appropriate. In patients who do require chronic treatment, the smallest dose and the shortest duration of treatment producing a satisfactory clinical response should be sought. The need for continued treatment should be reassessed periodically.

If signs and symptoms of tardive dyskinesia appear in a patient on quetiapine tablets, drug discontinuation should be considered. However, some patients may require treatment with quetiapine tablets despite the presence of the syndrome.

## **5.7 Hypotension**

Quetiapine may induce orthostatic hypotension associated with dizziness, tachycardia and, in some patients, syncope, especially during the initial dose-titration period, probably reflecting its  $\alpha_1$ -adrenergic antagonist properties. Syncope was reported in 1% (28/3,265) of the patients treated with quetiapine tablets, compared with 0.2% (2/954) on placebo and about 0.4% (2/527) on active control drugs. Orthostatic hypotension, dizziness, and syncope may lead to falls.

Quetiapine tablets should be used with particular caution in patients with known cardiovascular disease (history of myocardial infarction or ischemic heart disease, heart failure, or conduction abnormalities), cerebrovascular disease or conditions which would predispose patients to hypotension (dehydration, hypovolemia, and treatment with antihypertensive medications). The risk of orthostatic hypotension and syncope may be minimized by limiting the initial dose to 25 mg twice daily [see *Dosage and Administration* (2.2)]. If hypotension occurs during titration to the target dose, a return to the previous dose in the titration schedule is appropriate.

## **5.8 Falls**

Atypical antipsychotic drugs, including quetiapine tablets, may cause somnolence, postural hypotension, motor, and sensory instability, which may lead to falls and, consequently, fractures or other injuries. For patients with diseases, conditions, or medications that could exacerbate these effects, complete fall risk assessments when initiating antipsychotic treatment and recurrently for patients on long-term antipsychotic therapy.

## **5.9 Increases in Blood Pressure (Children and Adolescents)**

In placebo-controlled trials in children and adolescents with schizophrenia (6-week duration) or bipolar mania (3-week duration), the incidence of increases at any time in systolic blood pressure ( $\geq 20$  mmHg) was 15.2% (51/335) for quetiapine tablets and 5.5% (9/163) for placebo; the incidence of increases at any time in diastolic blood pressure ( $\geq 10$  mmHg) was 40.6% (136/335) for quetiapine tablets and 24.5% (40/163) for placebo. In the 26-week open-label clinical trial, one child with a reported history of hypertension experienced a hypertensive crisis. Blood pressure in children and adolescents should be measured at the beginning of, and periodically during treatment.

In a placebo-controlled quetiapine extended-release tablets clinical trial (8 weeks duration) in children and adolescents (10-17 years of age) with bipolar depression, in which efficacy was not established, the incidence of increases at any time in systolic blood pressure ( $\geq 20$  mmHg) was 6.5% (6/92) for quetiapine extended-release tablets and 6.0% (6/100) for placebo; the incidence of increases at any time in diastolic blood pressure ( $\geq 10$  mmHg) was 46.7% (43/92) for quetiapine extended-release tablets and 36.0% (36/100) for placebo.

### **5.10 Leukopenia, Neutropenia, and Agranulocytosis**

In clinical trial and postmarketing experience, events of leukopenia/neutropenia have been reported temporally related to atypical antipsychotic agents, including quetiapine tablets. Agranulocytosis has been reported.

Agranulocytosis (defined as absolute neutrophil count  $< 500/\text{mm}^3$ ) has been reported with quetiapine, including fatal cases and cases in patients without pre-existing risk factors. Neutropenia should be considered in patients presenting with infection, particularly in the absence of obvious predisposing factor(s), or in patients with unexplained fever, and should be managed as clinically appropriate.

Possible risk factors for leukopenia/neutropenia include pre-existing low white cell count (WBC) and history of drug induced leukopenia/neutropenia. Patients with a pre-existing low WBC or a history of drug induced leukopenia/neutropenia should have their complete blood count (CBC) monitored frequently during the first few months of therapy and should discontinue quetiapine tablets at the first sign of a decline in WBC in absence of other causative factors.

Patients with neutropenia should be carefully monitored for fever or other symptoms or signs of infection and treated promptly if such symptoms or signs occur. Patients with severe neutropenia (absolute neutrophil count  $< 1,000/\text{mm}^3$ ) should discontinue quetiapine tablets and have their WBC followed until recovery.

### **5.11 Cataracts**

The development of cataracts was observed in association with quetiapine treatment in chronic dog studies [see *Nonclinical Toxicology (13.2)*]. Lens changes have also been observed in adults, children, and adolescents during long-term quetiapine tablets treatment, but a causal relationship to quetiapine tablets use has not been established. Nevertheless, the possibility of lenticular changes cannot be excluded at this time. Therefore, examination of the lens by methods adequate to detect cataract formation, such as slit lamp exam or other appropriately sensitive methods, is recommended at initiation of treatment or shortly thereafter, and at 6-month intervals during chronic treatment.

### **5.12 QT Prolongation**

In clinical trials, quetiapine was not associated with a persistent increase in QT intervals. However, the QT effect was not systematically evaluated in a thorough QT study. In post marketing experience, there were cases reported of QT prolongation in patients who overdosed on quetiapine [see *Overdosage (10.1)*], in patients with concomitant illness,

and in patients taking medicines known to cause electrolyte imbalance or increase QT interval [see *Drug Interactions (7.1)*].

The use of quetiapine should be avoided in combination with other drugs that are known to prolong QTc including Class 1A antiarrhythmics (e.g., quinidine, procainamide) or Class III antiarrhythmics (e.g., amiodarone, sotalol), antipsychotic medications (e.g., ziprasidone, chlorpromazine, thioridazine), antibiotics (e.g., gatifloxacin, moxifloxacin), or any other class of medications known to prolong the QTc interval (e.g., pentamidine, levomethadyl acetate, methadone).

Quetiapine should also be avoided in circumstances that may increase the risk of occurrence of torsade de pointes and/or sudden death including (1) a history of cardiac arrhythmias such as bradycardia; (2) hypokalemia or hypomagnesemia; (3) concomitant use of other drugs that prolong the QTc interval; and (4) presence of congenital prolongation of the QT interval.

Caution should also be exercised when quetiapine is prescribed in patients with increased risk of QT prolongation (e.g., cardiovascular disease, family history of QT prolongation, the elderly, congestive heart failure, and heart hypertrophy).

### 5.13 Seizures

During clinical trials, seizures occurred in 0.5% (20/3,490) of patients treated with quetiapine tablets compared to 0.2% (2/954) on placebo and 0.7% (4/527) on active control drugs. As with other antipsychotics, quetiapine tablets should be used cautiously in patients with a history of seizures or with conditions that potentially lower the seizure threshold, e.g., Alzheimer's dementia. Conditions that lower the seizure threshold may be more prevalent in a population of 65 years or older.

### 5.14 Hypothyroidism

*Adults:* Clinical trials with quetiapine demonstrated dose-related decreases in thyroid hormone levels. The reduction in total and free thyroxine (T<sub>4</sub>) of approximately 20% at the higher end of the therapeutic dose range was maximal in the first six weeks of treatment and maintained without adaptation or progression during more chronic therapy. In nearly all cases, cessation of quetiapine treatment was associated with a reversal of the effects on total and free T<sub>4</sub>, irrespective of the duration of treatment. The mechanism by which quetiapine effects the thyroid axis is unclear. If there is an effect on the hypothalamic-pituitary axis, measurement of TSH alone may not accurately reflect a patient's thyroid status. Therefore, both TSH and free T<sub>4</sub>, in addition to clinical assessment, should be measured at baseline and at follow-up.

In the mania adjunct studies, where quetiapine tablets were added to lithium or divalproex, 12% (24/196) of quetiapine tablets treated patients compared to 7% (15/203) of placebo-treated patients had elevated TSH levels. Of the quetiapine tablets treated patients with elevated TSH levels, 3 had simultaneous low free T<sub>4</sub> levels (free T<sub>4</sub> <0.8 LLN).

About 0.7% (26/3,489) of quetiapine tablets patients did experience TSH increases in monotherapy studies. Some patients with TSH increases needed replacement thyroid treatment.

In all quetiapine trials, the incidence of shifts in thyroid hormones and TSH were<sup>1</sup>: decrease in free T<sub>4</sub> (<0.8 LLN), 2.0% (357/17,513); decrease in total T<sub>4</sub> (<0.8 LLN), 4.0% (75/1,861); decrease in free T<sub>3</sub> (<0.8 LLN), 0.4% (53/13,766); decrease in total T<sub>3</sub> (<0.8 LLN), 2.0% (26/1,312), and increase in TSH (>5 mIU/L), 4.9% (956/19,412). In eight patients, where TBG was measured, levels of TBG were unchanged.

Table 8 shows the incidence of these shifts in short-term placebo-controlled clinical trials.

**Table 8: Incidence of Shifts in Thyroid Hormone Levels and TSH in Short-Term Placebo-Controlled Clinical Trials<sup>1, 2</sup>**

Total T <sub>4</sub>		Free T <sub>4</sub>		Total T <sub>3</sub>		Free T <sub>3</sub>		TSH	
Quetiapine	Placebo	Quetiapine	Placebo	Quetiapine	Placebo	Quetiapine	Placebo	Quetiapine	Placebo
3.4%	0.6%	0.7%	0.1%	0.5%	0.0%	0.2%	0.0%	3.2%	2.7%
(37/1,097)	(4/651)	(52/7,218)	(4/3,668)	(2/369)	(0/113)	(11/5,673)	(1/2,679)	(240/7,587)	(105/3,912)

1. Based on shifts from normal baseline to potentially clinically important value at any time post-baseline. Shifts in total T<sub>4</sub>, free T<sub>4</sub>, total T<sub>3</sub>, and free T<sub>3</sub> are defined as <0.8 x LLN (pmol/L) and shift in TSH is >5 mIU/L at any time.

2. Includes quetiapine tablets and quetiapine extended-release tablets data.

In short-term placebo-controlled monotherapy trials, the incidence of reciprocal, shifts in T<sub>3</sub> and TSH was 0.0 % for both quetiapine (1/4,800) and placebo (0/2,190) and for T<sub>4</sub> and TSH the shifts were 0.1% (7/6,154) for quetiapine versus 0.0% (1/3,007) for placebo.

*Children and Adolescents:*

In acute placebo-controlled trials in children and adolescent patients with schizophrenia (6-week duration) or bipolar mania (3-week duration), the incidence of shifts for thyroid function values at any time for quetiapine tablets treated patients and placebo-treated patients for elevated TSH was 2.9% (8/280) vs. 0.7% (1/138), respectively, and for decreased total thyroxine was 2.8% (8/289) vs. 0% (0/145), respectively. Of the quetiapine tablets treated patients with elevated TSH levels, 1 had simultaneous low free T<sub>4</sub> level at end of treatment.

**5.15 Hyperprolactinemia**

*Adults:* During clinical trials with quetiapine, the incidence of shifts in prolactin levels to a clinically significant value occurred in 3.6% (158/4,416) of patients treated with quetiapine compared to 2.6% (51/1,968) on placebo.

*Children and Adolescents:*

In acute placebo-controlled trials in children and adolescent patients with bipolar mania (3-week duration) or schizophrenia (6-week duration), the incidence of shifts in prolactin levels to a value (>20 µg/L males; >26 µg/L females at any time) was 13.4% (18/134) for quetiapine tablets compared to 4% (3/75) for placebo in males and 8.7% (9/104) for quetiapine tablets compared to 0% (0/39) for placebo in females.

Like other drugs that antagonize dopamine D<sub>2</sub> receptors, quetiapine tablets elevate prolactin levels in some patients and the elevation may persist during chronic administration. Hyperprolactinemia, regardless of etiology, may suppress hypothalamic GnRH, resulting in reduced pituitary gonadotrophin secretion. This, in turn, may inhibit reproductive function by impairing gonadal steroidogenesis in both female and male patients. Galactorrhea, amenorrhea, gynecomastia, and impotence have been reported in patients receiving prolactin-elevating compounds. Long-standing hyperprolactinemia when associated with hypogonadism may lead to decreased bone density in both female and male subjects.

Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin dependent in vitro, a factor of potential importance if the prescription of these drugs is considered in a patient with previously detected breast cancer. As is common with compounds which increase prolactin release, mammary gland, and pancreatic islet cell neoplasia (mammary adenocarcinomas, pituitary, and pancreatic adenomas) was observed in carcinogenicity studies conducted in mice and rats [see Nonclinical Toxicology (13.1)]. Published epidemiologic studies have shown inconsistent results when exploring the potential association between

hyperprolactinemia and breast cancer.

### **5.16 Potential for Cognitive and Motor Impairment**

Somnolence was a commonly reported adverse event reported in patients treated with quetiapine tablets especially during the 3-5 day period of initial dose-titration. In schizophrenia trials, somnolence was reported in 18% (89/510) of patients on quetiapine tablets compared to 11% (22/206) of placebo patients. In acute bipolar mania trials using quetiapine tablets as monotherapy, somnolence was reported in 16% (34/209) of patients on quetiapine tablets compared to 4% of placebo patients. In acute bipolar mania trials using quetiapine tablets as adjunct therapy, somnolence was reported in 34% (66/196) of patients on quetiapine tablets compared to 9% (19/203) of placebo patients. In bipolar depression trials, somnolence was reported in 57% (398/698) of patients on quetiapine tablets compared to 15% (51/347) of placebo patients. Since quetiapine tablets have the potential to impair judgment, thinking, or motor skills, patients should be cautioned about performing activities requiring mental alertness, such as operating a motor vehicle (including automobiles) or operating hazardous machinery until they are reasonably certain that quetiapine tablets therapy does not affect them adversely. Somnolence may lead to falls.

### **5.17 Body Temperature Regulation**

Although not reported with quetiapine tablets, disruption of the body's ability to reduce core body temperature has been attributed to antipsychotic agents. Appropriate care is advised when prescribing quetiapine tablets for patients who will be experiencing conditions which may contribute to an elevation in core body temperature, e.g., exercising strenuously, exposure to extreme heat, receiving concomitant medication with anticholinergic activity, or being subject to dehydration.

### **5.18 Dysphagia**

Esophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in elderly patients, in particular those with advanced Alzheimer's dementia. Quetiapine tablets and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia.

### **5.19 Discontinuation Syndrome**

Acute withdrawal symptoms, such as insomnia, nausea, and vomiting have been described after abrupt cessation of atypical antipsychotic drugs, including quetiapine tablets. In short-term placebo-controlled, monotherapy clinical trials with quetiapine extended-release tablets that included a discontinuation phase which evaluated discontinuation symptoms, the aggregated incidence of patients experiencing one or more discontinuation symptoms after abrupt cessation was 12.1% (241/1,993) for quetiapine extended-release tablets and 6.7% (71/1,065) for placebo. The incidence of the individual adverse reactions (i.e., insomnia, nausea, headache, diarrhea, vomiting, dizziness, and irritability) did not exceed 5.3% in any treatment group and usually resolved after 1 week post-discontinuation. Gradual withdrawal is advised. [see *Use in Specific Populations* (8.1)].

### **5.20 Anticholinergic (antimuscarinic) Effects**

Norquetiapine, an active metabolite of quetiapine, has moderate to strong affinity for several muscarinic receptor subtypes. This contributes to anticholinergic adverse reactions when quetiapine tablets are used at therapeutic doses, taken concomitantly with other anticholinergic medications, or taken in overdose. Quetiapine tablets should be used with caution in patients receiving medications having anticholinergic (antimuscarinic) effects [see *Drug Interactions* (7.1), *Overdosage* (10.1), and *Clinical*

*Pharmacology (12.1)].*

Constipation was a commonly reported adverse event in patients treated with quetiapine and represents a risk factor for intestinal obstruction. Intestinal obstruction has been reported with quetiapine, including fatal reports in patients who were receiving multiple concomitant medications that decrease intestinal motility.

Quetiapine tablets should be used with caution in patients with a current diagnosis or prior history of urinary retention, clinically significant prostatic hypertrophy, or constipation.

## **6 ADVERSE REACTIONS**

The following adverse reactions are discussed in more detail in other sections of the labeling:

- Increased mortality in elderly patients with dementia-related psychosis [see *Warnings and Precautions (5.1)*]
- Suicidal thoughts and behaviors in adolescents and young adults [see *Warnings and Precautions (5.2)*]
- Cerebrovascular adverse reactions, including stroke in elderly patients with dementia-related psychosis [see *Warnings and Precautions (5.3)*]
- Neuroleptic Malignant Syndrome (NMS) [see *Warnings and Precautions (5.4)*]
- Metabolic changes (hyperglycemia, dyslipidemia, weight gain) [see *Warnings and Precautions (5.5)*]
- Tardive dyskinesia [see *Warnings and Precautions (5.6)*]
- Hypotension [see *Warnings and Precautions (5.7)*]
- Falls [see *Warnings and Precautions (5.8)*]
- Increases in blood pressure (children and adolescents) [see *Warnings and Precautions (5.9)*]
- Leukopenia, neutropenia and agranulocytosis [see *Warnings and Precautions (5.10)*]
- Cataracts [see *Warnings and Precautions (5.11)*]
- QT Prolongation [see *Warnings and Precautions (5.12)*]
- Seizures [see *Warnings and Precautions (5.13)*]
- Hypothyroidism [see *Warnings and Precautions (5.14)*]
- Hyperprolactinemia [see *Warnings and Precautions (5.15)*]
- Potential for cognitive and motor impairment [see *Warnings and Precautions (5.16)*]
- Body temperature regulation [see *Warnings and Precautions (5.17)*]
- Dysphagia [see *Warnings and Precautions (5.18)*]
- Discontinuation Syndrome [see *Warnings and Precautions (5.19)*]
- Anticholinergic (antimuscarinic) Effects [see *Warnings and Precautions (5.20)*]

### **6.1 Clinical Study Experience**

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

#### **Adults:**

The information below is derived from a clinical trial database for quetiapine tablets consisting of over 4,300 patients. This database includes 698 patients exposed to quetiapine tablets for the treatment of bipolar depression, 405 patients exposed to quetiapine tablets for the treatment of acute bipolar mania (monotherapy and adjunct therapy), 646 patients exposed to quetiapine tablets for the maintenance treatment of bipolar I disorder as adjunct therapy, and approximately 2,600 patients and/or normal subjects exposed to 1 or more doses of quetiapine tablets for the treatment of

schizophrenia.

Of these approximately 4,300 subjects, approximately 4,000 (2,300 in schizophrenia, 405 in acute bipolar mania, 698 in bipolar depression, and 646 for the maintenance treatment of bipolar I disorder) were patients who participated in multiple dose effectiveness trials, and their experience corresponded to approximately 2,400 patient-years. The conditions and duration of treatment with quetiapine tablets varied greatly and included (in overlapping categories) open-label and double-blind phases of studies, inpatients and outpatients, fixed-dose and dose-titration studies, and short-term or longer-term exposure. Adverse reactions were assessed by collecting adverse reactions, results of physical examinations, vital signs, weights, laboratory analyses, ECGs, and results of ophthalmologic examinations.

The stated frequencies of adverse reactions represent the proportion of individuals who experienced, at least once, an adverse reaction of the type listed.

### **Adverse Reactions Associated with Discontinuation of Treatment in Short-Term, Placebo-Controlled Trials**

*Schizophrenia:* Overall, there was little difference in the incidence of discontinuation due to adverse reactions (4% for quetiapine tablets vs. 3% for placebo) in a pool of controlled trials. However, discontinuations due to somnolence (0.8% quetiapine tablets vs. 0% placebo) and hypotension (0.4% quetiapine tablets vs. 0% placebo) were considered to be drug related [see *Warnings and Precautions (5.7 and 5.19)*].

*Bipolar Disorder:*

*Mania:* Overall, discontinuations due to adverse reactions were 5.7% for quetiapine tablets vs. 5.1% for placebo in monotherapy and 3.6% for quetiapine tablets vs. 5.9% for placebo in adjunct therapy.

*Depression:* Overall, discontinuations due to adverse reactions were 12.3% for quetiapine tablets 300 mg vs. 19.0% for quetiapine tablets 600 mg and 5.2% for placebo.

*Commonly Observed Adverse Reactions in Short-Term, Placebo-Controlled Trials:*

In the acute therapy of schizophrenia (up to 6 weeks) and bipolar mania (up to 12 weeks) trials, the most commonly observed adverse reactions associated with the use of quetiapine tablets monotherapy (incidence of 5% or greater) and observed at a rate on quetiapine tablets at least twice that of placebo were somnolence (18%), dizziness (11%), dry mouth (9%), constipation (8%), ALT increased (5%), weight gain (5%), and dyspepsia (5%).

*Adverse Reactions Occurring at an Incidence of 2% or More Among Quetiapine Tablets Treated Patients in Short-Term, Placebo-Controlled Trials:*

The prescriber should be aware that the figures in the tables and tabulations cannot be used to predict the incidence of side effects in the course of usual medical practice where patient characteristics and other factors differ from those that prevailed in the clinical trials. Similarly, the cited frequencies cannot be compared with figures obtained from other clinical investigations involving different treatments, uses, and investigators. The cited figures, however, do provide the prescribing physician with some basis for estimating the relative contribution of drug and non-drug factors to the side effect incidence in the population studied.

Table 9 enumerates the incidence, rounded to the nearest percent, of adverse reactions that occurred during acute therapy of schizophrenia (up to 6 weeks) and bipolar mania (up to 12 weeks) in 2% or more of patients treated with quetiapine tablets (doses ranging from 75 to 800 mg/day) where the incidence in patients treated with quetiapine tablets was greater than the incidence in placebo-treated patients.

**Table 9: Adverse Reaction Incidence in 3- to 12-Week Placebo-Controlled Clinical Trials for the Treatment of Schizophrenia and Bipolar Mania (Monotherapy)**

<b>Preferred Term</b>	<b>Quetiapine Tablets (n=719)</b>	<b>Placebo (n=404)</b>
Headache	21%	14%
Agitation	20%	17%
Somnolence	18%	8%
Dizziness	11%	5%
Dry Mouth	9%	3%
Constipation	8%	3%
Pain	7%	5%
Tachycardia	6%	4%
Vomiting	6%	5%
Asthenia	5%	3%
Dyspepsia	5%	1%
Weight Gain	5%	1%
ALT Increased	5%	1%
Anxiety	4%	3%
Pharyngitis	4%	3%
Rash	4%	2%
Abdominal Pain	4%	1%
Postural Hypotension	4%	1%
Back Pain	3%	1%
AST Increased	3%	1%
Rhinitis	3%	1%
Fever	2%	1%
Gastroenteritis	2%	0%
Amblyopia	2%	1%

In the acute adjunct therapy of bipolar mania (up to 3 weeks) studies, the most commonly observed adverse reactions associated with the use of quetiapine tablets (incidence of 5% or greater) and observed at a rate on quetiapine tablets at least twice that of placebo were somnolence (34%), dry mouth (19%), asthenia (10%), constipation (10%), abdominal pain (7%), postural hypotension (7%), pharyngitis (6%), and weight gain (6%).

Table 10 enumerates the incidence, rounded to the nearest percent, of adverse reactions that occurred during therapy (up to 3 weeks) of acute mania in 2% or more of patients treated with quetiapine tablets (doses ranging from 100 to 800 mg/day) used as adjunct therapy to lithium and divalproex where the incidence in patients treated with quetiapine tablets was greater than the incidence in placebo-treated patients.

**Table 10: Adverse Reaction Incidence in 3-Week Placebo-Controlled Clinical Trials for the Treatment of Bipolar Mania (Adjunct Therapy)**

<b>Preferred Term</b>	<b>Quetiapine Tablets (n=196)</b>	<b>Placebo (n=203)</b>
Somnolence	34%	9%
Dry Mouth	19%	3%
Headache	17%	13%
Asthenia	10%	4%
Constipation	10%	5%

Dizziness	9%	6%
Tremor	8%	7%
Abdominal Pain	7%	3%
Postural Hypotension	7%	2%
Agitation	6%	4%
Weight Gain	6%	3%
Pharyngitis	6%	3%
Back Pain	5%	3%
Hypertonia	4%	3%
Rhinitis	4%	2%
Peripheral Edema	4%	2%
Twitching	4%	1%
Dyspepsia	4%	3%
Depression	3%	2%
Amblyopia	3%	2%
Speech Disorder	3%	1%
Hypotension	3%	1%
Hormone Level Altered	3%	0%
Heaviness	2%	1%
Infection	2%	1%
Fever	2%	1%
Hypertension	2%	1%
Tachycardia	2%	1%
Increased Appetite	2%	1%
Hypothyroidism	2%	1%
Incoordination	2%	1%
Thinking Abnormal	2%	0%
Anxiety	2%	0%
Ataxia	2%	0%
Sinusitis	2%	1%
Sweating	2%	1%
Urinary Tract Infection	2%	1%

In bipolar depression studies (up to 8 weeks), the most commonly observed adverse reactions associated with the use of quetiapine tablets (incidence of 5% or greater) and observed at a rate on quetiapine tablets at least twice that of placebo were somnolence (57%), dry mouth (44%), dizziness (18%), constipation (10%), and lethargy (5%).

Table 11 enumerates the incidence, rounded to the nearest percent, of adverse reactions that occurred during therapy (up to 8 weeks) of bipolar depression in 2% or more of patients treated with quetiapine tablets (doses of 300 and 600 mg/day) where the incidence in patients treated with quetiapine tablets was greater than the incidence in placebo-treated patients.

**Table 11: Adverse Reaction Incidence in 8-Week Placebo-Controlled Clinical Trials for the Treatment of Bipolar Depression**

<b>Preferred Term</b>	<b>Quetiapine Tablets (n=698)</b>	<b>Placebo (n=347)</b>
Somnolence <sup>1</sup>	57%	15%
Dry Mouth	44%	13%
Dizziness	18%	7%

Constipation	10%	4%
Fatigue	10%	8%
Dyspepsia	7%	4%
Vomiting	5%	4%
Increased Appetite	5%	3%
Lethargy	5%	2%
Nasal Congestion	5%	3%
Orthostatic Hypotension	4%	3%
Akathisia	4%	1%
Palpitations	4%	1%
Vision Blurred	4%	2%
Weight increased	4%	1%
Arthralgia	3%	2%
Paraesthesia	3%	2%
Cough	3%	1%
Extrapyramidal Disorder	3%	1%
Irritability	3%	1%
Dysarthria	3%	0%
Hypersomnia	3%	0%
Sinus Congestion	2%	1%
Abnormal Dreams	2%	1%
Tremor	2%	1%
Gastroesophageal Reflux Disease	2%	1%
Pain in Extremity	2%	1%
Asthenia	2%	1%
Balance Disorder	2%	1%
Hypoesthesia	2%	1%
Dysphagia	2%	0%
Restless Legs Syndrome	2%	0%

1. Somnolence combines adverse reaction terms somnolence and sedation

Explorations for interactions on the basis of gender, age, and race did not reveal any clinically meaningful differences in the adverse reaction occurrence on the basis of these demographic factors.

Dose Dependency of Adverse Reactions in Short-Term, Placebo-Controlled Trials

*Dose-related Adverse Reactions:* Spontaneously elicited adverse reaction data from a study of schizophrenia comparing five fixed doses of quetiapine tablets (75 mg, 150 mg, 300 mg, 600 mg, and 750 mg/day) to placebo were explored for dose-relatedness of adverse reactions. Logistic regression analyses revealed a positive dose response ( $p < 0.05$ ) for the following adverse reactions: dyspepsia, abdominal pain, and weight gain.

Adverse Reactions in clinical trials with quetiapine and not listed elsewhere in the label:

The following adverse reactions have also been reported with quetiapine: nightmares, hypersensitivity, and elevations in serum creatine phosphokinase (not associated with NMS), galactorrhea, bradycardia (which may occur at or near initiation of treatment and be associated with hypotension and/or syncope) decreased platelets, somnambulism (and other related events), elevations in gamma-GT levels, hypothermia, dyspnea, eosinophilia, urinary retention, intestinal obstruction and priapism.

Extrapyramidal Symptoms (EPS):

## Dystonia

**Class Effect:** Symptoms of dystonia, prolonged abnormal contractions of muscle groups, may occur in susceptible individuals during the first few days of treatment. Dystonic symptoms include: spasm of the neck muscles, sometimes progressing to tightness of the throat, swallowing difficulty, difficulty breathing, and/or protrusion of the tongue. While these symptoms can occur at low doses, they occur more frequently and with greater severity with high potency and at higher doses of first generation antipsychotic drugs. An elevated risk of acute dystonia is observed in males and younger age groups.

Four methods were used to measure EPS: (1) Simpson-Angus total score (mean change from baseline) which evaluates Parkinsonism and akathisia, (2) Barnes Akathisia Rating Scale (BARS) Global Assessment Score, (3) incidence of spontaneous complaints of EPS (akathisia, akinesia, cogwheel rigidity, extrapyramidal syndrome, hypertonia, hypokinesia, neck rigidity, and tremor), and (4) use of anticholinergic medications to treat EPS.

**Adults:** Data from one 6-week clinical trial of schizophrenia comparing five fixed doses of quetiapine tablets (75, 150, 300, 600, 750 mg/day) provided evidence for the lack of extrapyramidal symptoms (EPS) and dose-relatedness for EPS associated with quetiapine tablets treatment. Three methods were used to measure EPS: (1) Simpson-Angus total score (mean change from baseline) which evaluates Parkinsonism and akathisia, (2) incidence of spontaneous complaints of EPS (akathisia, akinesia, cogwheel rigidity, extrapyramidal syndrome, hypertonia, hypokinesia, neck rigidity, and tremor), and (3) use of anticholinergic medications to EPS.

In Table 12, dystonic event included nuchal rigidity, hypertonia, dystonia, muscle rigidity, oculogyration; parkinsonism included cogwheel rigidity, tremor, drooling, hypokinesia; akathisia included akathisia, psychomotor agitation; dyskinesic event included tardive dyskinesia, dyskinesia, choreoathetosis; and other extrapyramidal event included restlessness, extrapyramidal disorder, movement disorder.

**Table 12: Adverse Reactions Associated with EPS in a Short-Term, Placebo-Controlled Multiple Fixed-Dose Phase III Schizophrenia Trial (6 weeks duration)**

Preferred Term	Quetiapine Tablets 75 mg/day (N=53)		Quetiapine Tablets 150 mg/day (N=48)		Quetiapine Tablets 300 mg/day (N=52)		Quetiapine Tablets 600 mg/day (N=51)		Quetiapine Tablets 750 mg/day (N=54)		Placebo (N=51)	
	n	%	n	%	n	%	n	%	n	%	n	%
Dystonic event	2	3.8	2	4.2	0	0.0	2	3.9	3	5.6	4	7.8
Parkinsonism	2	3.8	0	0.0	1	1.9	1	2.0	1	1.9	4	7.8
Akathisia	1	1.9	1	2.1	0	0.0	0	0.0	1	1.9	4	7.8
Dyskinetic event	2	3.8	0	0.0	0	0.0	1	2.0	0	0.0	0	0.0
Other extrapyramidal event	2	3.8	0	0.0	3	5.8	3	5.9	1	1.9	4	7.8

Parkinsonism incidence rates as measured by the Simpson-Angus total score for placebo and the five fixed doses (75, 150, 300, 600, 750 mg/day) were: -0.6; -1.0, -1.2; -1.6; -1.8, and -1.8. The rate of anticholinergic medication use to treat EPS for placebo and the five fixed doses was: 14%; 11%; 10%; 8%; 12%, and 11%.

In six additional placebo-controlled clinical trials (3 in acute mania and 3 in schizophrenia) using variable doses of quetiapine tablets, there were no differences between the quetiapine tablets and placebo treatment groups in the incidence of EPS, as assessed by Simpson-Angus total scores, spontaneous complaints of EPS and the use of concomitant anticholinergic medications to treat EPS.

In two placebo-controlled clinical trials for the treatment of bipolar depression using 300 mg and 600 mg of quetiapine tablets, the incidence of adverse reactions potentially related to EPS was 12% in both dose groups and 6% in the placebo group. In these studies, the incidence of the individual adverse reactions (akathisia, extrapyramidal disorder, tremor, dyskinesia, dystonia, restlessness, muscle contractions involuntary, psychomotor hyperactivity, and muscle rigidity) were generally low and did not exceed 4% in any treatment group.

The 3 treatment groups were similar in mean change in SAS total score and BARS Global Assessment score at the end of treatment. The use of concomitant anticholinergic medications was infrequent and similar across the three treatment groups.

### **Children and Adolescents**

The information below is derived from a clinical trial database for quetiapine tablets consisting of over 1,000 pediatric patients. This database includes 677 patients exposed to quetiapine tablets for the treatment of schizophrenia and 393 children and adolescents (10-17 years old) exposed to quetiapine tablets for the treatment of acute bipolar mania.

#### **Adverse Reactions Associated with Discontinuation of Treatment in Short-Term, Placebo-Controlled Trials**

**Schizophrenia:** The incidence of discontinuation due to adverse reactions for quetiapine-treated and placebo-treated patients was 8.2% and 2.7%, respectively. The adverse event leading to discontinuation in 1% or more of patients on quetiapine tablets and at a greater incidence than placebo was somnolence (2.7% and 0% for placebo).

**Bipolar I Mania:** The incidence of discontinuation due to adverse reactions for quetiapine-treated and placebo-treated patients was 11.4% and 4.4%, respectively. The adverse reactions leading to discontinuation in 2% or more of patients on quetiapine tablets and at a greater incidence than placebo were somnolence (4.1% vs. 1.1%) and fatigue (2.1% vs. 0%).

#### **Commonly Observed Adverse Reactions in Short-Term, Placebo-Controlled Trials**

In therapy for schizophrenia (up to 6 weeks), the most commonly observed adverse reactions associated with the use of quetiapine in adolescents (incidence of 5% or greater and quetiapine incidence at least twice that for placebo) were somnolence (34%), dizziness (12%), dry mouth (7%), tachycardia (7%).

In bipolar mania therapy (up to 3 weeks) the most commonly observed adverse reactions associated with the use of quetiapine in children and adolescents (incidence of 5% or greater and quetiapine incidence at least twice that for placebo) were somnolence (53%), dizziness (18%), fatigue (11%), increased appetite (9%), nausea (8%), vomiting (8%), tachycardia (7%), dry mouth (7%), and weight increased (6%).

In an acute (8-week) quetiapine extended-release tablets trial in children and adolescents (10-17 years of age) with bipolar depression, in which efficacy was not established, the most commonly observed adverse reactions associated with the use of quetiapine extended-release tablets (incidence of 5% or greater and at least twice that for placebo) were dizziness 7%, diarrhea 5%, fatigue 5%, and nausea 5%.

#### **Adverse Reactions Occurring at an Incidence of $\geq 2\%$ among Quetiapine Tablets Treated Patients in Short-Term, Placebo-Controlled Trials**

## Schizophrenia (Adolescents, 13-17 years old)

The following findings were based on a 6-week placebo-controlled trial in which quetiapine was administered in either doses of 400 or 800 mg/day.

Table 13 enumerates the incidence, rounded to the nearest percent, of adverse reactions that occurred during therapy (up to 6 weeks) of schizophrenia in 2% or more of patients treated with quetiapine tablets (doses of 400 or 800 mg/day) where the incidence in patients treated with quetiapine tablets was at least twice the incidence in placebo-treated patients.

Adverse reactions that were potentially dose-related with higher frequency in the 800 mg group compared to the 400 mg group included dizziness (8% vs. 15%), dry mouth (4% vs. 10%), and tachycardia (6% vs. 11%).

**Table 13: Adverse Reaction Incidence in a 6-Week Placebo-Controlled Clinical Trial for the Treatment of Schizophrenia in Adolescent Patients**

Preferred Term	Quetiapine Tablets 400 mg (n=73)	Quetiapine Tablets 800 mg (n=74)	Placebo (n=75)
Somnolence <sup>1</sup>	33%	35%	11%
Dizziness	8%	15%	5%
Dry Mouth	4%	10%	1%
Tachycardia <sup>2</sup>	6%	11%	0%
Irritability	3%	5%	0%
Arthralgia	1%	3%	0%
Asthenia	1%	3%	1%
Back Pain	1%	3%	0%
Dyspnea	0%	3%	0%
Abdominal Pain	3%	1%	0%
Anorexia	3%	1%	0%
Tooth Abscess	3%	1%	0%
Dyskinesia	3%	0%	0%
Epistaxis	3%	0%	1%
Muscle Rigidity	3%	0%	0%

1. Somnolence combines adverse reaction terms somnolence and sedation.

2. Tachycardia combines adverse reaction terms tachycardia and sinus tachycardia.

## Bipolar I Mania (Children and Adolescents 10-17 years old)

The following findings were based on a 3-week placebo-controlled trial in which quetiapine was administered in either doses of 400 or 600 mg/day.

### Commonly Observed Adverse Reactions

In bipolar mania therapy (up to 3 weeks) the most commonly observed adverse reactions associated with the use of quetiapine in children and adolescents (incidence of 5% or greater and quetiapine incidence at least twice that for placebo) were somnolence (53%), dizziness (18%), fatigue (11%), increased appetite (9%), nausea (8%), vomiting (8%), tachycardia (7%), dry mouth (7%), and weight increased (6%).

Table 14 enumerates the incidence, rounded to the nearest percent, of adverse reactions that occurred during therapy (up to 3 weeks) of bipolar mania in 2% or more of patients treated with quetiapine tablets (doses of 400 or 600 mg/day) where the incidence in patients treated with quetiapine tablets was greater than the incidence in placebo-treated patients.

Adverse reactions that were potentially dose-related with higher frequency in the 600

mg group compared to the 400 mg group included somnolence (50% vs. 57%), nausea (6% vs. 10%), and tachycardia (6% vs. 9%).

**Table 14: Adverse Reactions in a 3-Week Placebo-Controlled Clinical Trial for the Treatment of Bipolar Mania in Children and Adolescent Patients**

Preferred Term	Quetiapine Tablets 400 mg (n=95)	Quetiapine Tablets 600 mg (n=98)	Placebo (n=90)
Somnolence <sup>1</sup>	50%	57%	14%
Dizziness	19%	17%	2%
Nausea	6%	10%	4%
Fatigue	14%	9%	4%
Increased Appetite	10%	9%	1%
Tachycardia <sup>2</sup>	6%	9%	1%
Dry Mouth	7%	7%	0%
Vomiting	8%	7%	3%
Nasal Congestion	3%	6%	2%
Weight Increased	6%	6%	0%
Irritability	3%	5%	1%
Pyrexia	1%	4%	1%
Aggression	1%	3%	0%
Musculoskeletal Stiffness	1%	3%	1%
Accidental Overdose	0%	2%	0%
Acne	3%	2%	0%
Arthralgia	4%	2%	1%
Lethargy	2%	2%	0%
Pallor	1%	2%	0%
Stomach Discomfort	4%	2%	1%
Syncope	2%	2%	0%
Vision Blurred	3%	2%	0%
Constipation	4%	2%	0%
Ear Pain	2%	0%	0%
Paresthesia	2%	0%	0%
Sinus Congestion	3%	0%	0%
Thirst	2%	0%	0%

1. Somnolence combines adverse reactions terms somnolence and sedation.

2. Tachycardia combines adverse reaction terms tachycardia and sinus tachycardia.

#### *Extrapyramidal Symptoms:*

In a short-term placebo-controlled monotherapy trial in adolescent patients with schizophrenia (6-week duration), the aggregated incidence of extrapyramidal symptoms was 12.9% (19/147) for quetiapine tablets and 5.3% (4/75) for placebo, though the incidence of the individual adverse reactions (akathisia, tremor, extrapyramidal disorder, hypokinesia, restlessness, psychomotor hyperactivity, muscle rigidity, dyskinesia) did not exceed 4.1% in any treatment group. In a short-term placebo-controlled monotherapy trial in children and adolescent patients with bipolar mania (3-week duration), the aggregated incidence of extrapyramidal symptoms was 3.6% (7/193) for quetiapine tablets and 1.1% (1/90) for placebo.

Table 15 presents a listing of patients with adverse reactions potentially associated with extrapyramidal symptoms in the short-term placebo-controlled monotherapy trial in adolescent patients with schizophrenia (6-week duration).

In Tables 15-16 dystonic event included nuchal rigidity, hypertonia, and muscle rigidity; parkinsonism included cogwheel rigidity and tremor; akathisia included akathisia only; dyskinetic event included tardive dyskinesia, dyskinesia, and choreoathetosis; and other extrapyramidal event included restlessness and extrapyramidal disorder.

**Table 15: Adverse Reactions Associated with Extrapyramidal Symptoms in the Placebo-Controlled Trial in Adolescent Patients with Schizophrenia (6-week duration)**

Preferred Term	Quetiapine Tablets 400 mg/day (N=73)		Quetiapine Tablets 800 mg/day (N=74)		All Quetiapine Tablets (N=147)		Placebo (N=75)	
	n	%	n	%	n	%	n	%
Dystonic event	2	2.7	0	0.0	2	1.4	0	0.0
Parkinsonism	4	5.5	4	5.4	8	5.4	2	2.7
Akathisia	3	4.1	4	5.4	7	4.8	3	4.0
Dyskinetic event	2	2.7	0	0.0	2	1.4	0	0.0
Other Extrapyramidal Event	2	2.7	2	2.7	4	2.7	0	0.0

Table 16 presents a listing of patients with adverse reactions associated with extrapyramidal symptoms in a short-term placebo-controlled monotherapy trial in children and adolescent patients with bipolar mania (3-week duration).

**Table 16: Adverse Reactions Associated with Extrapyramidal Symptoms in a Placebo-Controlled Trial in Children and Adolescent Patients with Bipolar I Mania (3-week duration)**

Preferred Term <sup>1</sup>	Quetiapine Tablets 400 mg/day (N=95)		Quetiapine Tablets 600 mg/day (N=98)		All Quetiapine Tablets (N=193)		Placebo (N=90)	
	n	%	n	%	n	%	n	%
Parkinsonism	2	2.1	1	1.0	3	1.6	1	1.1
Akathisia	1	1.0	1	1.0	2	1.0	0	0.0
Other Extrapyramidal Event	1	1.1	1	1.0	2	1.0	0	0.0

1. There were no adverse reactions with the preferred term of dystonic or dyskinetic events.

## Laboratory, ECG, and vital sign changes observed in clinical studies

Laboratory Changes:

Neutrophil Counts

*Adults:* In placebo-controlled monotherapy clinical trials involving 3,368 patients on quetiapine and 1,515 on placebo, the incidence of at least one occurrence of neutrophil count  $<1.0 \times 10^9/L$  among patients with a normal baseline neutrophil count and at least one available follow up laboratory measurement was 0.3% (10/2,967) in patients treated with quetiapine, compared to 0.1% (2/1,349) in patients treated with placebo [see *Warnings and Precautions (5.10)*].

Transaminase Elevations

*Adults:* Asymptomatic, transient, and reversible elevations in serum transaminases (primarily ALT) have been reported. In schizophrenia trials in adults, the proportions of patients with transaminase elevations of >3 times the upper limits of the normal reference range in a pool of 3- to 6-week placebo-controlled trials were approximately 6% (29/483) for quetiapine tablets compared to 1% (3/194) for placebo. In acute bipolar mania trials in adults, the proportions of patients with transaminase elevations of >3 times the upper limits of the normal reference range in a pool of 3- to 12-week placebo-controlled trials were approximately 1% for both quetiapine tablets (3/560) and placebo (3/294). These hepatic enzyme elevations usually occurred within the first 3 weeks of drug treatment and promptly returned to pre-study levels with ongoing treatment with quetiapine tablets. In bipolar depression trials, the proportions of patients with transaminase elevations of >3 times the upper limits of the normal reference range in two 8-week placebo-controlled trials was 1% (5/698) for quetiapine tablets and 2% (6/347) for placebo.

#### Decreased Hemoglobin

*Adults:* In short-term placebo-controlled trials, decreases in hemoglobin to  $\leq 13$  g/dL males,  $\leq 12$  g/dL females on at least one occasion occurred in 8.3% (594/7,155) of quetiapine-treated patients compared to 6.2% (219/3,536) of patients treated with placebo. In a database of controlled and uncontrolled clinical trials, decreases in hemoglobin to  $\leq 13$  g/dL males,  $\leq 12$  g/dL females on at least one occasion occurred in 11% (2,277/20,729) of quetiapine-treated patients.

#### Interference with Urine Drug Screens

There have been literature reports suggesting false positive results in urine enzyme immunoassays for methadone and tricyclic antidepressants in patients who have taken quetiapine. Caution should be exercised in the interpretation of positive urine drug screen results for these drugs, and confirmation by alternative analytical technique (e.g., chromatographic methods) should be considered.

#### ECG Changes

*Adults:* Between-group comparisons for pooled placebo-controlled trials revealed no statistically significant quetiapine tablets /placebo differences in the proportions of patients experiencing potentially important changes in ECG parameters, including QT, QTc, and PR intervals. However, the proportions of patients meeting the criteria for tachycardia were compared in four 3- to 6-week placebo-controlled clinical trials for the treatment of schizophrenia revealing a 1% (4/399) incidence for quetiapine tablets compared to 0.6% (1/156) incidence for placebo. In acute (monotherapy) bipolar mania trials the proportions of patients meeting the criteria for tachycardia was 0.5% (1/192) for quetiapine tablets compared to 0% (0/178) incidence for placebo. In acute bipolar mania (adjunct) trials the proportions of patients meeting the same criteria was 0.6% (1/166) for quetiapine tablets compared to 0% (0/171) incidence for placebo. In bipolar depression trials, no patients had heart rate increases to >120 beats per minute. Quetiapine tablets use was associated with a mean increase in heart rate, assessed by ECG, of 7 beats per minute compared to a mean increase of 1 beat per minute among placebo patients. This slight tendency to tachycardia in adults may be related to quetiapine tablets' potential for inducing orthostatic changes [see *Warnings and Precautions* (5.7)].

*Children and Adolescents:* In the acute (6-week) schizophrenia trial in adolescents, increases in heart rate (>110 bpm) occurred in 5.2% (3/73) of patients receiving quetiapine tablets 400 mg and 8.5% (5/74) of patients receiving quetiapine tablets 800 mg compared to 0% (0/75) of patients receiving placebo. Mean increases in heart rate were 3.8 bpm and 11.2 bpm for quetiapine tablets 400 mg and 800 mg groups, respectively, compared to a decrease of 3.3 bpm in the placebo group [see *Warnings and Precautions* (5.7)].

In the acute (3-week) bipolar mania trial in children and adolescents, increases in heart rate (>110 bpm) occurred in 1.1% (1/89) of patients receiving quetiapine tablets 400 mg and 4.7% (4/85) of patients receiving quetiapine tablets 600 mg compared to 0% (0/98) of patients receiving placebo. Mean increases in heart rate were 12.8 bpm and 13.4 bpm for quetiapine tablets 400 mg and 600 mg groups, respectively, compared to a decrease of 1.7 bpm in the placebo group [see *Warnings and Precautions (5.7)*].

In an acute (8-week) quetiapine extended-release tablets trial in children and adolescents (10-17 years of age) with bipolar depression, in which efficacy was not established, increases in heart rate (>110 bpm 10-12 years and 13-17 years) occurred in 0% of patients receiving quetiapine extended-release tablets and 1.2% of patients receiving placebo. Mean increases in heart rate were 3.4 bpm for quetiapine extended-release tablets, compared to 0.3 bpm in the placebo group [see *Warnings and Precautions (5.7)*].

## **6.2 Postmarketing Experience**

The following adverse reactions were identified during post approval of quetiapine tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Adverse reactions reported since market introduction which were temporally related to quetiapine therapy include anaphylactic reaction, cardiomyopathy, drug reaction with eosinophilia and systemic symptoms (DRESS), hyponatremia, myocarditis, nocturnal enuresis, pancreatitis, retrograde amnesia, rhabdomyolysis, syndrome of inappropriate antidiuretic hormone secretion (SIADH), Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), decreased platelet count, serious liver reactions (including hepatitis, liver necrosis, and hepatic failure), agranulocytosis, intestinal obstruction, ileus, colon ischemia, urinary retention, sleep apnea, acute generalized exanthematous pustulosis (AGEP), confusional state, cutaneous vasculitis, and fecal incontinence.

## **7 DRUG INTERACTIONS**

### **7.1 Effect of Other Drugs on Quetiapine**

The risks of using quetiapine tablets in combination with other drugs have not been extensively evaluated in systematic studies. Given the primary CNS effects of quetiapine tablets, caution should be used when it is taken in combination with other centrally acting drugs. Quetiapine tablets potentiated the cognitive and motor effects of alcohol in a clinical trial in subjects with selected psychotic disorders, and alcoholic beverages should be limited while taking quetiapine.

Quetiapine exposure is increased by the prototype CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, indinavir, ritonavir, nefazodone, etc.) and decreased by the prototype CYP3A4 inducers (e.g., phenytoin, carbamazepine, rifampin, avasimibe, St. John's wort etc.). Dose adjustment of quetiapine will be necessary if it is co-administered with potent CYP3A4 inducers or inhibitors.

CYP3A4 inhibitors:

Coadministration of ketoconazole, a potent inhibitor of cytochrome CYP3A4, resulted in significant increase in quetiapine exposure. The dose of quetiapine tablets should be reduced to one sixth of the original dose if co-administered with a strong CYP3A4 inhibitor [see *Dosage and Administration (2.5)* and *Clinical Pharmacology (12.3)*].

CYP3A4 inducers:

Coadministration of quetiapine and phenytoin, a CYP3A4 inducer increased the mean oral clearance of quetiapine by 5-fold. Increased doses of quetiapine tablets up to 5 fold

may be required to maintain control of symptoms of schizophrenia in patients receiving quetiapine and phenytoin, or other known potent CYP3A4 inducers [see *Dosage and Administration (2.6) and Clinical Pharmacology (12.3)*]. When the CYP3A4 inducer is discontinued, the dose of quetiapine tablets should be reduced to the original level within 7-14 days [see *Dosage and Administration (2.6)*].

#### Anticholinergic Drugs:

Concomitant treatment with quetiapine and other drugs with anticholinergic activity can increase the risk for severe gastrointestinal adverse reactions related to hypomotility. Quetiapine tablets should be used with caution in patients receiving medications having anticholinergic (antimuscarinic) effects [see *Warnings and Precautions (5.20)*].

The potential effects of several concomitant medications on quetiapine pharmacokinetics were studied [see *Clinical Pharmacology (12.3)*].

### 7.2 Effect of Quetiapine on Other Drugs

Because of its potential for inducing hypotension, quetiapine tablets may enhance the effects of certain antihypertensive agents.

Quetiapine tablets may antagonize the effects of levodopa and dopamine agonists.

There are no clinically relevant pharmacokinetic interactions of quetiapine tablets on other drugs based on the CYP pathway. Quetiapine tablets and its metabolites are non-inhibitors of major metabolizing CYP's (1A2, 2C9, 2C19, 2D6, and 3A4).

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Neonates exposed to antipsychotic drugs (including quetiapine tablets) during the third trimester are at risk for extrapyramidal and/or withdrawal symptoms following delivery (see *Clinical Considerations*). Overall available data from published epidemiologic studies of pregnant women exposed to quetiapine have not established a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes (see *Data*). There are risks to the mother associated with untreated schizophrenia, bipolar I, or major depressive disorder, and with exposure to antipsychotics, including quetiapine tablets, during pregnancy (see *Clinical Considerations*).

In animal studies, embryo-fetal toxicity occurred including delays in skeletal ossification at approximately 1 and 2 times the maximum recommended human dose (MRHD) of 800 mg/day in both rats and rabbits, and an increased incidence of carpal/tarsal flexure (minor soft tissue anomaly) in rabbit fetuses at approximately 2 times the MRHD. In addition, fetal weights were decreased in both species. Maternal toxicity (observed as decreased body weights and/or death) occurred at 2 times the MRHD in rats and approximately 1-2 times the MRHD in rabbits.

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

#### Clinical Considerations

##### *Disease-associated maternal and/or fetal risk*

There is a risk to the mother from untreated schizophrenia, or bipolar I disorder, including increased risk of relapse, hospitalization, and suicide. Schizophrenia and bipolar

I disorder are associated with increased adverse perinatal outcomes, including preterm birth. It is not known if this is a direct result of the illness or other comorbid factors.

A prospective, longitudinal study followed 201 pregnant women with a history of major depressive disorder who were euthymic and taking antidepressants at the beginning of pregnancy. The women who discontinued antidepressants during pregnancy were more likely to experience a relapse of major depression than women who continued antidepressants. Consider the risk of untreated depression when discontinuing or changing treatment with antidepressant medication during pregnancy and postpartum.

#### *Fetal/neonatal adverse reactions*

Extrapyramidal and/or withdrawal symptoms, including agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, and feeding disorder have been reported in neonates who were exposed to antipsychotic drugs, including quetiapine tablets, during the third trimester of pregnancy. These symptoms varied in severity. Monitor neonates for extrapyramidal and/or withdrawal symptoms and manage symptoms appropriately. Some neonates recovered within hours or days without specific treatment; others required prolonged hospitalization.

#### Data

##### *Human Data*

Published data from observational studies, birth registries, and case reports on the use of atypical antipsychotics during pregnancy do not report a clear association with antipsychotics and major birth defects. A retrospective cohort study from a Medicaid database of 9,258 women exposed to antipsychotics during pregnancy did not indicate an overall increased risk of major birth defects.

##### *Animal Data*

When pregnant rats and rabbits were exposed to quetiapine during organogenesis, there was no teratogenic effect in fetuses. Doses were 25, 50 and 200 mg/kg in rats and 25, 50 and 100 mg/kg in rabbits which are approximately 0.3, 0.6 and 2-times (rats) and 0.6, 1 and 2-times (rabbits) the MRHD for schizophrenia of 800 mg/day based on mg/m<sup>2</sup> body surface area. However, there was evidence of embryo-fetal toxicity including delays in skeletal ossification at approximately 1 and 2 times the MRHD of 800 mg/day in both rats and rabbits, and an increased incidence of carpal/tarsal flexure (minor soft tissue anomaly) in rabbit fetuses at approximately 2 times the MRHD. In addition, fetal weights were decreased in both species. Maternal toxicity (observed as decreased body weights and/or death) occurred at 2 times the MRHD in rats and approximately 1-2 times the MRHD (all doses tested) in rabbits.

In a peri/postnatal reproductive study in rats, no drug-related effects were observed when pregnant dams were treated with quetiapine at doses 0.01, 0.1, and 0.2 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area. However, in a preliminary peri/postnatal study, there were increases in fetal and pup death, and decreases in mean litter weight at 3 times the MRHD.

## **8.2 Lactation**

### Risk Summary

Limited data from published literature report the presence of quetiapine in human breast milk at relative infant dose of <1% of the maternal weight-adjusted dosage. There are no consistent adverse events that have been reported in infants exposed to quetiapine through breast milk. There is no information on the effects of quetiapine on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for quetiapine tablets and any potential adverse effects on the breastfed child from quetiapine tablets or from the mother's

underlying condition.

### **8.3 Females and Males of Reproductive Potential**

#### Infertility

##### *Females*

Based on the pharmacologic action of quetiapine (D2 antagonism), treatment with quetiapine tablets may result in an increase in serum prolactin levels, which may lead to a reversible reduction in fertility in females of reproductive potential [see *Warnings and Precautions (5.15)*].

### **8.4 Pediatric Use**

In general, the adverse reactions observed in children and adolescents during the clinical trials were similar to those in the adult population with few exceptions. Increases in systolic and diastolic blood pressure occurred in children and adolescents and did not occur in adults. Orthostatic hypotension occurred more frequently in adults (4-7%) compared to children and adolescents (<1%) [see *Warnings and Precautions (5.7)* and *Adverse Reactions (6.1)*].

#### Schizophrenia

The efficacy and safety of quetiapine tablets in the treatment of schizophrenia in adolescents aged 13-17 years were demonstrated in one 6-week, double-blind, placebo-controlled trial [see *Indications and Usage (1.1)*, *Dosage and Administration (2.2)*, *Adverse Reactions (6.1)*, and *Clinical Studies (14.1)*].

Safety and effectiveness of quetiapine tablets in pediatric patients less than 13 years of age with schizophrenia have not been established.

#### Maintenance

The safety and effectiveness of quetiapine tablets in the maintenance treatment of bipolar disorder has not been established in pediatric patients less than 18 years of age. The safety and effectiveness of quetiapine tablets in the maintenance treatment of schizophrenia has not been established in any patient population, including pediatric patients.

#### Bipolar Mania

The efficacy and safety of quetiapine tablets in the treatment of mania in children and adolescents ages 10-17 years with bipolar I disorder was demonstrated in a 3-week, double-blind, placebo-controlled, multicenter trial [see *Indications and Usage (1.2)*, *Dosage and Administration (2.3)*, *Adverse Reactions (6.1)*, and *Clinical Studies (14.2)*].

Safety and effectiveness of quetiapine tablets in pediatric patients less than 10 years of age with bipolar mania have not been established.

#### Bipolar Depression

Safety and effectiveness of quetiapine tablets in pediatric patients less than 18 years of age with bipolar depression have not been established. A clinical trial with quetiapine extended-release tablets was conducted in children and adolescents (10- 17 years of age) with bipolar depression, efficacy was not established.

Some differences in the pharmacokinetics of quetiapine were noted between children/adolescents (10-17 years of age) and adults. When adjusted for weight, the AUC and C<sub>max</sub> of quetiapine were 41% and 39% lower, respectively, in children and adolescents compared to adults. The pharmacokinetics of the active metabolite, norquetiapine, were similar between children/adolescents and adults after adjusting for weight [see *Clinical Pharmacology (12.3)*].

## **8.5 Geriatric Use**

Of the approximately 3,700 patients in clinical studies with quetiapine tablets, 7% (232) were 65 years of age or over. In general, there was no indication of any different tolerability of quetiapine tablets in the elderly compared to younger adults. Nevertheless, the presence of factors that might decrease pharmacokinetic clearance, increase the pharmacodynamic response to quetiapine tablets, or cause poorer tolerance or orthostasis, should lead to consideration of a lower starting dose, slower titration, and careful monitoring during the initial dosing period in the elderly. The mean plasma clearance of quetiapine tablets was reduced by 30% to 50% in elderly patients when compared to younger patients [see *Clinical Pharmacology (12.3) and Dosage and Administration (2.3)*].

## **8.6 Renal Impairment**

Clinical experience with quetiapine tablets in patients with renal impairment is limited [see *Clinical Pharmacology (12.3)*].

## **8.7 Hepatic Impairment**

Since quetiapine is extensively metabolized by the liver, higher plasma levels are expected in patients with hepatic impairment. In this population, a low starting dose of 25 mg/day is recommended and the dose may be increased in increments of 25 mg/day-50 mg/day [see *Dosage and Administration (2.4) and Clinical Pharmacology (12.3)*].

# **9 DRUG ABUSE AND DEPENDENCE**

## **9.1 Controlled Substance**

Quetiapine fumarate is not a controlled substance.

## **9.2 Abuse**

Quetiapine tablets have not been systematically studied, in animals or humans, for its potential for abuse, tolerance, or physical dependence. While the clinical trials did not reveal any tendency for any drug-seeking behavior, these observations were not systematic and it is not possible to predict on the basis of this limited experience the extent to which a CNS-active drug will be misused, diverted, and/or abused once marketed. Consequently, patients should be evaluated carefully for a history of drug abuse, and such patients should be observed closely for signs of misuse or abuse of quetiapine tablets, e.g., development of tolerance, increases in dose, drug-seeking behavior.

# **10 OVERDOSAGE**

## **10.1 Human Experience**

In clinical trials, survival has been reported in acute overdoses of up to 30 grams of quetiapine. Most patients who overdosed experienced no adverse reactions or recovered fully from the reported reactions. Death has been reported in a clinical trial following an overdose of 13.6 grams of quetiapine alone. In general, reported signs and symptoms were those resulting from an exaggeration of the drug's known pharmacological effects, i.e., drowsiness, sedation, tachycardia, hypotension, and anticholinergic toxicity including coma and delirium. Patients with pre-existing severe cardiovascular disease may be at an increased risk of the effects of overdose [see *Warnings and Precautions (5.12)*]. One case, involving an estimated overdose of 9,600



## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The mechanism of action of quetiapine in the listed indications is unclear. However, the efficacy of quetiapine in these indications could be mediated through a combination of dopamine type 2 (D<sub>2</sub>) and serotonin type 2 (5HT<sub>2</sub>) antagonism. The active metabolite, N-desalkyl quetiapine (norquetiapine), has similar activity at D<sub>2</sub>, but greater activity at 5HT<sub>2A</sub> receptors, than the parent drug (quetiapine).

### 12.2 Pharmacodynamics

Quetiapine and its metabolite, norquetiapine, have affinity for multiple neurotransmitter receptors with norquetiapine binding with higher affinity than quetiapine in general. The K<sub>i</sub> values for quetiapine and norquetiapine at the dopamine D<sub>1</sub> are 428/99.8 nM, at D<sub>2</sub> 626/489 nM, at serotonin 5HT<sub>1A</sub> 1,040/191 nM at 5HT<sub>2A</sub> 38/2.9 nM, at histamine H<sub>1</sub> 4.4/1.1 nM, at muscarinic M<sub>1</sub> 1,086/38.3 nM, and at adrenergic α<sub>1b</sub> 14.6/46.4 nM and, at α<sub>2</sub> receptors 617/1,290 nM, respectively. Quetiapine and norquetiapine lack appreciable affinity to the benzodiazepine receptors.

#### Effect on QT Interval

In clinical trials, quetiapine was not associated with a persistent increase in QT intervals. However, the QT effect was not systematically evaluated in a thorough QT study. In post marketing experience, there were cases reported of QT prolongation in patients who overdosed on quetiapine [see *Overdosage (10.1)*], in patients with concomitant illness, and in patients taking medicines known to cause electrolyte imbalance or increase QT interval.

### 12.3 Pharmacokinetics

#### Adults

Quetiapine activity is primarily due to the parent drug. The multiple-dose pharmacokinetics of quetiapine are dose-proportional within the proposed clinical dose range, and quetiapine accumulation is predictable upon multiple dosing. Elimination of quetiapine is mainly via hepatic metabolism with a mean terminal half-life of about 6 hours within the proposed clinical dose range. Steady-state concentrations are expected to be achieved within two days of dosing. Quetiapine is unlikely to interfere with the metabolism of drugs metabolized by cytochrome P450 enzymes.

#### Children and Adolescents

At steady state the pharmacokinetics of the parent compound, in children and adolescents (10-17 years of age), were similar to adults. However, when adjusted for dose and weight, AUC and C<sub>max</sub> of the parent compound were 41% and 39% lower, respectively, in children and adolescents than in adults. For the active metabolite, norquetiapine, AUC and C<sub>max</sub> were 45% and 31% higher, respectively, in children and adolescents than in adults. When adjusted for dose and weight, the pharmacokinetics of the metabolite, norquetiapine, was similar between children and adolescents and adults [see *Use in Specific Populations (8.4)*].

#### Absorption

Quetiapine is rapidly absorbed after oral administration, reaching peak plasma concentrations in 1.5 hours. The tablet formulation is 100% bioavailable relative to solution. The bioavailability of quetiapine is marginally affected by administration with food, with C<sub>max</sub> and AUC values increased by 25% and 15%, respectively.

#### Distribution

Quetiapine is widely distributed throughout the body with an apparent volume of distribution of  $10 \pm 4$  L/kg. It is 83% bound to plasma proteins at therapeutic concentrations. *In vitro*, quetiapine did not affect the binding of warfarin or diazepam to human serum albumin. In turn, neither warfarin nor diazepam altered the binding of quetiapine.

#### *Metabolism and Elimination*

Following a single oral dose of  $^{14}\text{C}$ -quetiapine, less than 1% of the administered dose was excreted as unchanged drug, indicating that quetiapine is highly metabolized. Approximately 73% and 20% of the dose was recovered in the urine and feces, respectively.

Quetiapine is extensively metabolized by the liver. The major metabolic pathways are sulfoxidation to the sulfoxide metabolite and oxidation to the parent acid metabolite; both metabolites are pharmacologically inactive. *In vitro* studies using human liver microsomes revealed that the cytochrome P450 3A4 isoenzyme is involved in the metabolism of quetiapine to its major, but inactive, sulfoxide metabolite and in the metabolism of its active metabolite N-desalkyl quetiapine.

#### *Age*

Oral clearance of quetiapine was reduced by 40% in elderly patients ( $\geq 65$  years,  $n=9$ ) compared to young patients ( $n=12$ ), and dosing adjustment may be necessary [see *Dosage and Administration (2.3)*].

#### *Gender*

There is no gender effect on the pharmacokinetics of quetiapine.

#### *Race*

There is no race effect on the pharmacokinetics of quetiapine.

#### *Smoking*

Smoking has no effect on the oral clearance of quetiapine.

#### *Renal Insufficiency*

Patients with severe renal impairment ( $\text{Clcr}=10\text{-}30$  mL/min/ $1.73$  m<sup>2</sup>,  $n=8$ ) had a 25% lower mean oral clearance than normal subjects ( $\text{Clcr} > 80$  mL/min/ $1.73$  m<sup>2</sup>,  $n=8$ ), but plasma quetiapine concentrations in the subjects with renal insufficiency were within the range of concentrations seen in normal subjects receiving the same dose. Dosage adjustment is therefore not needed in these patients [see *Use in Specific Populations (8.6)*].

#### *Hepatic Insufficiency*

Hepatically impaired patients ( $n=8$ ) had a 30% lower mean oral clearance of quetiapine than normal subjects. In two of the 8 hepatically impaired patients, AUC and  $C_{\text{max}}$  were 3 times higher than those observed typically in healthy subjects. Since quetiapine is extensively metabolized by the liver, higher plasma levels are expected in the hepatically impaired population, and dosage adjustment may be needed [see *Dosage and Administration (2.4)* and *Use in Specific Populations (8.7)*].

#### *Drug-Drug Interaction Studies*

The *in vivo* assessments of effect of other drugs on the pharmacokinetics of quetiapine are summarized in Table 17 [see *Dosage and Administration (2.5 and 2.6)* and *Drug Interactions (7.1)*].

---

### **Table 17: The Effect of Other Drugs on the Pharmacokinetics of Quetiapine**

Coadministered drug	Dose schedules		Effect on quetiapine pharmacokinetics
	Coadministered Drug	Quetiapine	
Phenytoin	100 mg three times daily	250 mg three times daily	5-fold increase in oral clearance
Divalproex	500 mg twice daily	150 mg twice daily	17% increase mean max plasma concentration at steady state. No effect on absorption or mean oral clearance
Thioridazine	200 mg twice daily	300 mg twice daily	65% increase in oral clearance
Cimetidine	400 mg three times daily for 4 days	150 mg three times daily	20% decrease in mean oral clearance
Ketoconazole (potent CYP 3A4 inhibitor)	200 mg once daily for 4 days	25 mg single dose	84% decrease in oral clearance resulting in a 6.2-fold increase in AUC of quetiapine
Fluoxetine	60 mg once daily	300 mg twice daily	No change in steady state PK
Imipramine	75 mg twice daily	300 mg twice daily	No change in steady state PK
Haloperidol	7.5 mg twice daily	300 mg twice daily	No change in steady state PK
Risperidone	3 mg twice daily	300 mg twice daily	No change in steady state PK

*In vitro* enzyme inhibition data suggest that quetiapine and 9 of its metabolites would have little inhibitory effect on *in vivo* metabolism mediated by cytochromes CYP 1A2, 2C9, 2C19, 2D6, and 3A4. Quetiapine at doses of 750 mg/day did not affect the single dose pharmacokinetics of antipyrine, lithium, or lorazepam (Table 18) [see *Drug Interactions* (7.2)].

**Table 18: The Effect of Quetiapine on the Pharmacokinetics of Other Drugs**

Coadministered drug	Dose schedules		Effect on other drugs pharmacokinetics
	Coadministered drug	Quetiapine	
Lorazepam	2 mg, single dose	250 mg three times daily	Oral clearance of lorazepam reduced by 20%
Divalproex	500 mg twice daily	150 mg twice daily	C <sub>max</sub> and AUC of free valproic acid at steady-state was decreased by 10-12%
Lithium	Up to 2,400 mg/day given in twice daily doses	250 mg three times daily	No effect on steady-state pharmacokinetics of lithium
Antipyrine	1 g, single dose	250 mg three times daily	No effect on clearance of antipyrine or urinary recovery of its metabolites

## 13 NONCLINICAL TOXICOLOGY

## 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### *Carcinogenesis*

Carcinogenicity studies were conducted in C57BL mice and Wistar rats. Quetiapine was administered in the diet to mice at doses of 20, 75, 250, and 750 mg/kg and to rats by gavage at doses of 25, 75, and 250 mg/kg for two years. These doses are equivalent to 0.1, 0.5, 1.5, and 4.5 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area (mice) or 0.3, 1, and 3 times the MRHD based on mg/m<sup>2</sup> body surface area (rats). There were statistically significant increases in thyroid gland follicular adenomas in male mice at doses 1.5 and 4.5 times the MRHD based on mg/m<sup>2</sup> body surface area and in male rats at a dose of 3 times the MRHD on mg/m<sup>2</sup> body surface area. Mammary gland adenocarcinomas were statistically significantly increased in female rats at all doses tested (0.3, 1, and 3 times the MRHD based on mg/m<sup>2</sup> body surface area).

Thyroid follicular cell adenomas may have resulted from chronic stimulation of the thyroid gland by thyroid stimulating hormone (TSH) resulting from enhanced metabolism and clearance of thyroxine by rodent liver. Changes in TSH, thyroxine, and thyroxine clearance consistent with this mechanism were observed in subchronic toxicity studies in rat and mouse and in a 1-year toxicity study in rat; however, the results of these studies were not definitive. The relevance of the increases in thyroid follicular cell adenomas to human risk, through whatever mechanism, is unknown.

Antipsychotic drugs have been shown to chronically elevate prolactin levels in rodents. Serum measurements in a 1-year toxicity study showed that quetiapine increased median serum prolactin levels a maximum of 32- and 13-fold in male and female rats, respectively. Increases in mammary neoplasms have been found in rodents after chronic administration of other antipsychotic drugs and are considered to be prolactin-mediated. The relevance of this increased incidence of prolactin-mediated mammary gland tumors in rats to human risk is unknown [see *Warnings and Precautions (5.15)*].

### *Mutagenesis*

Quetiapine was not mutagenic or clastogenic in standard genotoxicity tests. The mutagenic potential of quetiapine was tested in the *in vitro* Ames bacterial gene mutation assay and in the *in vitro* mammalian gene mutation assay in Chinese Hamster Ovary cells. The clastogenic potential of quetiapine was tested in the *in vitro* chromosomal aberration assay in cultured human lymphocytes and in the *in vivo* bone marrow micronucleus assay in rats up to 500 mg/kg, which is 6 times the MRHD based on mg/m<sup>2</sup> body surface area.

### *Impairment of Fertility*

Quetiapine decreased mating and fertility in male Sprague-Dawley rats at oral doses of 50 and 150 mg/kg or approximately 1 and 3 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area. Drug-related effects included increases in interval to mate and in the number of matings required for successful impregnation. These effects continued to be observed at 3 times the MRHD even after a two-week period without treatment. The no-effect dose for impaired mating and fertility in male rats was 25 mg/kg, or 0.3 times the MRHD based on mg/m<sup>2</sup> body surface area. Quetiapine adversely affected mating and fertility in female Sprague-Dawley rats at an oral dose approximately 1 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area. Drug-related effects included decreases in matings and in matings resulting in pregnancy, and an increase in the interval to mate. An increase in irregular estrus cycles was observed at doses of 10 and 50 mg/kg, or approximately 0.1 and 1 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area. The no-effect dose in female rats was 1 mg/kg, or 0.01 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area.

## 13.2 Animal Toxicology and/or Pharmacology

Quetiapine caused a dose-related increase in pigment deposition in thyroid gland in rat

toxicity studies which were 4 weeks in duration or longer and in a mouse 2-year carcinogenicity study. Doses were 10, 25, 50, 75, 150 and 250 mg/kg in rat studies which are approximately 0.1, 0.3, 0.6, 1, 2 and 3-times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area, respectively. Doses in the mouse carcinogenicity study were 20, 75, 250 and 750 mg/kg which are approximately 0.1, 0.5, 1.5, and 4.5 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area. Pigment deposition was shown to be irreversible in rats. The identity of the pigment could not be determined, but was found to be co-localized with quetiapine in thyroid gland follicular epithelial cells. The functional effects and the relevance of this finding to human risk are unknown.

In dogs receiving quetiapine for 6 or 12 months, but not for 1-month, focal triangular cataracts occurred at the junction of posterior sutures in the outer cortex of the lens at a dose of 100 mg/kg, or 4 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area. This finding may be due to inhibition of cholesterol biosynthesis by quetiapine. Quetiapine caused a dose-related reduction in plasma cholesterol levels in repeat-dose dog and monkey studies; however, there was no correlation between plasma cholesterol and the presence of cataracts in individual dogs. The appearance of delta-8-cholestanol in plasma is consistent with inhibition of a late stage in cholesterol biosynthesis in these species. There also was a 25% reduction in cholesterol content of the outer cortex of the lens observed in a special study in quetiapine treated female dogs. Drug-related cataracts have not been seen in any other species; however, in a 1-year study in monkeys, a striated appearance of the anterior lens surface was detected in 2/7 females at a dose of 225 mg/kg or 5.5 times the MRHD of 800 mg/day based on mg/m<sup>2</sup> body surface area.

## **14 CLINICAL STUDIES**

### **14.1 Schizophrenia**

#### **Short-term Trials-Adults**

The efficacy of quetiapine tablets in the treatment of schizophrenia was established in 3 short-term (6-week) controlled trials of inpatients with schizophrenia who met DSM III-R criteria for schizophrenia. Although a single fixed dose haloperidol arm was included as a comparative treatment in one of the three trials, this single haloperidol dose group was inadequate to provide a reliable and valid comparison of quetiapine tablets and haloperidol.

Several instruments were used for assessing psychiatric signs and symptoms in these studies, among them the Brief Psychiatric Rating Scale (BPRS), a multi-item inventory of general psychopathology traditionally used to evaluate the effects of drug treatment in schizophrenia. The BPRS psychosis cluster (conceptual disorganization, hallucinatory behavior, suspiciousness, and unusual thought content) is considered a particularly useful subset for assessing actively psychotic schizophrenic patients. A second traditional assessment, the Clinical Global Impression (CGI), reflects the impression of a skilled observer, fully familiar with the manifestations of schizophrenia, about the overall clinical state of the patient.

The results of the trials follow:

1. In a 6-week, placebo-controlled trial (n=361) (Study 1) involving 5 fixed doses of quetiapine tablets (75 mg/day, 150 mg/day, 300 mg/day, 600 mg/day, and 750 mg/day given in divided doses three times per day), the 4 highest doses of quetiapine tablets were generally superior to placebo on the BPRS total score, the BPRS psychosis cluster and the CGI severity score, with the maximal effect seen at 300 mg/day, and the effects of doses of 150 mg/day to 750 mg/day were generally indistinguishable.
2. In a 6-week, placebo-controlled trial (n=286) (Study 2) involving titration of quetiapine tablets in high (up to 750 mg/day given in divided doses three times per day) and low

(up to 250 mg/day given in divided doses three times per day) doses, only the high dose quetiapine tablets group (mean dose, 500 mg/day) was superior to placebo on the BPRS total score, the BPRS psychosis cluster, and the CGI severity score.

3. In a 6-week dose and dose regimen comparison trial (n=618) (Study 3) involving two fixed doses of quetiapine tablets (450 mg/day given in divided doses both twice daily and three times daily and 50 mg/day given in divided doses twice daily), only the 450 mg/day (225 mg given twice daily) dose group was superior to the 50 mg/day (25 mg given twice daily) quetiapine tablets dose group on the BPRS total score, the BPRS psychosis cluster, and the CGI severity score.

The primary efficacy results of these three studies in the treatment of schizophrenia in adults is presented in Table 19.

Examination of population subsets (race, gender, and age) did not reveal any differential responsiveness on the basis of race or gender, with an apparently greater effect in patients under the age of 40 years compared to those older than 40. The clinical significance of this finding is unknown.

### Adolescents (ages 13-17)

The efficacy of quetiapine tablets in the treatment of schizophrenia in adolescents (13-17 years of age) was demonstrated in a 6-week, double-blind, placebo-controlled trial (Study 4). Patients who met DSM-IV diagnostic criteria for schizophrenia were randomized into one of three treatment groups: quetiapine tablets 400 mg/day (n=73), quetiapine tablets 800 mg/day (n=74), or placebo (n=75). Study medication was initiated at 50 mg/day and on day 2 increased to 100 mg/per day (divided and given two or three times per day). Subsequently, the dose was titrated to the target dose of 400 mg/day or 800 mg/day using increments of 100 mg/day, divided and given two or three times daily. The primary efficacy variable was the mean change from baseline in total Positive and Negative Syndrome Scale (PANSS).

Quetiapine tablets at 400 mg/day and 800 mg/day was superior to placebo in the reduction of PANSS total score. The primary efficacy results of this study in the treatment of schizophrenia in adolescents is presented in Table 19.

**Table 19: Schizophrenia Short-Term Trials**

Study Number	Treatment Group	Primary Efficacy Endpoint: BPRS Total		
		Mean Baseline Score (SD)	LS Mean Change from Baseline (SE)	Placebo-subtracted Difference <sup>1</sup> (95% CI)
Study 1	Quetiapine tablets (75 mg/day)	45.7 (10.9)	-2.2 (2.0)	-4.0 (-11.2, 3.3)
	Quetiapine tablets (150 mg/day) <sup>2</sup>	47.2 (10.1)	-8.7 (2.1)	-10.4 (-17.8, -3.0)
	Quetiapine tablets (300 mg/day) <sup>2</sup>	45.3 (10.9)	-8.6 (2.1)	-10.3 (-17.6, -3.0)
	Quetiapine tablets (600 mg/day) <sup>2</sup>	43.5 (11.3)	-7.7 (2.1)	-9.4 (-16.7, -2.1)
	Quetiapine tablets (750 mg/day) <sup>2</sup>	45.7 (11.0)	-6.3 (2.0)	-8.0 (-15.2, -0.8)

	Placebo	45.3 (9.2)	1.7 (2.1)	--
Study 2	Quetiapine tablets (250 mg/day)	38.9 (9.8)	-4.2 (1.6)	-3.2 (-7.6, 1.2)
	Quetiapine tablets (750 mg/day) <sup>2</sup>	41.0 (9.6)	-8.7 (1.6)	-7.8 (-12.2, -3.4)
	Placebo	38.4 (9.7)	-1.0 (1.6)	--
Study 3	Quetiapine tablets (450 mg/day BID)	42.1 (10.7)	-10.0 (1.3)	-4.6 (-7.8, -1.4)
	Quetiapine tablets (450 mg/day TID) <sup>3</sup>	42.7 (10.4)	-8.6 (1.3)	-3.2 (-6.4, 0.0)
	Quetiapine tablets (50 mg BID)	41.7 (10.0)	-5.4 (1.3)	--
		<b>Primary Efficacy Endpoint: PANSS Total</b>		
		<b>Mean Baseline Score (SD)</b>	<b>LS Mean Change from Baseline (SE)</b>	<b>Placebo-subtracted Difference<sup>1</sup> (95% CI)</b>
Study 4	Quetiapine tablets (400 mg/day) <sup>2</sup>	96.2 (17.7)	-27.3 (2.6)	-8.2 (-16.1, -0.3)
	Quetiapine tablets (800 mg/day) <sup>2</sup>	96.9 (15.3)	-28.4 (1.8)	-9.3 (-16.2, -2.4)
	Placebo	96.2 (17.7)	-19.2 (3.0)	--

SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: unadjusted confidence interval.

1. Difference (drug minus placebo) in least-squares mean change from baseline.

2. Doses that are statistically significant superior to placebo.

3. Doses that are statistically significant superior to quetiapine tablets 50 mg BID.

## 14.2 Bipolar Disorder

Bipolar I disorder, manic or mixed episodes

Adults

The efficacy of quetiapine tablets in the acute treatment of manic episodes was established in 3 placebo-controlled trials in patients who met DSM-IV criteria for bipolar I disorder with manic episodes. These trials included patients with or without psychotic features and excluded patients with rapid cycling and mixed episodes. Of these trials, 2 were monotherapy (12 weeks) and 1 was adjunct therapy (3 weeks) to either lithium or divalproex. Key outcomes in these trials were change from baseline in the Young Mania Rating Scale (YMRS) score at 3 and 12 weeks for monotherapy and at 3 weeks for adjunct therapy. Adjunct therapy is defined as the simultaneous initiation or subsequent administration of quetiapine tablets with lithium or divalproex.

The primary rating instrument used for assessing manic symptoms in these trials was YMRS, an 11-item clinician-rated scale traditionally used to assess the degree of manic symptomatology (irritability, disruptive/aggressive behavior, sleep, elevated mood, speech, increased activity, sexual interest, language/thought disorder, thought content, appearance, and insight) in a range from 0 (no manic features) to 60 (maximum score).

The results of the trials follow:

### *Monotherapy*

The efficacy of quetiapine tablets in the acute treatment of bipolar mania was established in 2 placebo-controlled trials. In two 12-week trials (n=300, n=299) comparing quetiapine tablets to placebo, quetiapine tablets were superior to placebo in the reduction of the YMRS total score at weeks 3 and 12. The majority of patients in these trials taking quetiapine tablets were dosed in a range between 400 mg/day and 800 mg per day (studies 1 and 2 in Table 20).

### *Adjunct Therapy*

In this 3-week placebo-controlled trial, 170 patients with bipolar mania (YMRS  $\geq 20$ ) were randomized to receive quetiapine tablets or placebo as adjunct treatment to lithium or divalproex. Patients may or may not have received an adequate treatment course of lithium or divalproex prior to randomization. Quetiapine tablets were superior to placebo when added to lithium or divalproex alone in the reduction of YMRS total score (Study 3 in Table 20).

The majority of patients in this trial taking quetiapine tablets were dosed in a range between 400 mg/day and 800 mg per day. In a similarly designed trial (n=200), quetiapine tablets were associated with an improvement in YMRS scores but did not demonstrate superiority to placebo, possibly due to a higher placebo effect.

The primary efficacy results of these studies in the treatment of mania in adults is presented in Table 20.

### Children and Adolescents (ages 10-17)

The efficacy of quetiapine tablets in the acute treatment of manic episodes associated with bipolar I disorder in children and adolescents (10-17 years of age) was demonstrated in a 3-week, double-blind, placebo-controlled, multicenter trial (Study 4 in Table 20). Patients who met DSM-IV diagnostic criteria for a manic episode were randomized into one of three treatment groups: quetiapine tablets 400 mg/day (n=95), quetiapine tablets 600 mg/day (n=98), or placebo (n=91). Study medication was initiated at 50 mg/day and on day 2 increased to 100 mg/day (divided doses given two or three times daily). Subsequently, the dose was titrated to a target dose of 400 mg/day or 600 mg/day using increments of 100 mg/day, given in divided doses two or three times daily. The primary efficacy variable was the mean change from baseline in total YMRS score.

Quetiapine tablets 400 mg/day and 600 mg/day were superior to placebo in the reduction of YMRS total score (Table 20).

**Table 20: Mania Trials**

Study Number	Treatment Group	Primary Efficacy Measure: YMRS Total		
		Mean Baseline Score (SD) <sup>4</sup>	LS Mean Change from Baseline (SE)	Placebo-subtracted Difference <sup>2</sup> (95% CI)
Study 1	Quetiapine tablets (200-800 mg/day) <sup>1, 3</sup>	34.0 (6.1)	-12.3 (1.3)	-4.0 (-7.0, -1.0)
	Haloperidol <sup>1, 3</sup>	32.3 (6.0)	-15.7 (1.3)	-7.4 (-10.4, -4.4)
	Placebo	33.1 (6.6)	-8.3 (1.3)	--
Study 2	Quetiapine tablets (200-800 mg/day) <sup>1</sup>	32.7 (6.5)	-14.6 (1.5)	-7.9 (-10.9, -5.0)
	Lithium <sup>1, 3</sup>	33.3 (7.1)	-15.2 (1.6)	-8.5 (-11.5, -5.5)

	Placebo	34.0 (6.9)	-6.7 (1.6)	--
Study 3	Quetiapine tablets (200-800 mg/day) <sup>1</sup> + mood stabilizer	31.5(5.8)	-13.8(1.6)	-3.8(-7.1,-0.6)
	Placebo + mood stabilizer	31.1(5.5)	-10(1.5)	--
Study 4	Quetiapine tablets (400 mg/day) <sup>1</sup>	29.4 (5.9)	-14.3 (0.96)	-5.2 (-8.1, -2.3)
	Quetiapine tablets (600 mg/day) <sup>1</sup>	29.6 (6.4)	-15.6 (0.97)	-6.6 (-9.5, -3.7)
	Placebo	30.7 (5.9)	-9.0 (1.1)	--

Mood stabilizer: lithium or divalproex; SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: unadjusted confidence interval.

1. Doses that are statistically significantly superior to placebo.

2. Difference (drug minus placebo) in least-squares mean change from baseline.

3. Included in the trial as an active comparator.

4. Adult data mean baseline score is based on patients included in the primary analysis; pediatric mean baseline score is based on all patients in the ITT population.

## Bipolar Disorder, Depressive Episodes

### Adults

The efficacy of quetiapine tablets for the acute treatment of depressive episodes associated with bipolar disorder was established in 2 identically designed 8-week, randomized, double-blind, placebo-controlled studies (N=1,045) (studies 5 and 6 in Table 21). These studies included patients with either bipolar I or II disorder and those with or without a rapid cycling course. Patients randomized to quetiapine tablets were administered fixed doses of either 300 mg or 600 mg once daily.

The primary rating instrument used to assess depressive symptoms in these studies was the Montgomery-Asberg Depression Rating Scale (MADRS), a 10-item clinician-rated scale with scores ranging from 0 to 60. The primary endpoint in both studies was the change from baseline in MADRS score at week 8. In both studies, quetiapine tablets were superior to placebo in reduction of MADRS score. Improvement in symptoms, as measured by change in MADRS score relative to placebo, was seen in both studies at Day 8 (week 1) and onwards. In these studies, no additional benefit was seen with the 600 mg dose. For the 300 mg dose group, statistically significant improvements over placebo were seen in overall quality of life and satisfaction related to various areas of functioning, as measured using the Q-LES-Q(SF).

The primary efficacy results of these studies in the acute treatment of depressive episodes associated with bipolar disorder in adults is presented in Table 21.

**Table 21: Depressive Episodes Associated with Bipolar Disorder**

Study Number	Treatment Group	Primary Efficacy Measure: MADRS Total		
		Mean Baseline Score (SD)	LS Mean Change from Baseline (SE)	Placebo-subtracted Difference <sup>2</sup> (95% CI)
Study 5	Quetiapine tablets (300 mg/day) <sup>1</sup>	30.3 (5.0)	-16.4 (0.9)	-6.1 (-8.3, -3.9)
	Quetiapine tablets (600)	30.3 (5.3)	-16.7 (0.9)	-6.5 (-8.7, -4.3)

	mg/day) <sup>1</sup>			
	Placebo	30.6 (5.3)	-10.3 (0.9)	--
Study 6	Quetiapine tablets (300 mg/day) <sup>1</sup>	31.1 (5.7)	-16.9 (1.0)	-5.0 (-7.3, -2.7)
	Quetiapine tablets (600 mg/day) <sup>1</sup>	29.9 (5.6)	-16.0 (1.0)	-4.1 (-6.4, -1.8)
	Placebo	29.6 (5.4)	-11.9 (1.0)	--

SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: unadjusted confidence interval.

1. Doses that are statistically significantly superior to placebo.


2. Difference (drug minus placebo) in least-squares mean change from baseline.

## Maintenance Treatment as an Adjunct to Lithium or Divalproex


The efficacy of quetiapine tablets in the maintenance treatment of bipolar I disorder was established in 2 placebo-controlled trials in patients (n=1,326) who met DSM-IV criteria for bipolar I disorder (studies 7 and 8 in Figures 1 and 2). The trials included patients whose most recent episode was manic, depressed, or mixed, with or without psychotic features. In the open-label phase, patients were required to be stable on quetiapine tablets plus lithium or divalproex for at least 12 weeks in order to be randomized. On average, patients were stabilized for 15 weeks. In the randomization phase, patients continued treatment with lithium or divalproex and were randomized to receive either quetiapine tablets (administered twice daily totaling 400 mg/day to 800 mg/day) or placebo. Approximately 50% of the patients had discontinued from the quetiapine tablets group by day 280 and 50% of the placebo group had discontinued by day 117 of double-blind treatment. The primary endpoint in these studies was time to recurrence of a mood event (manic, mixed, or depressed episode). A mood event was defined as medication initiation or hospitalization for a mood episode; YMRS score  $\geq 20$  or MADRS score  $\geq 20$  at 2 consecutive assessments; or study discontinuation due to a mood event (Figure 1 and Figure 2).

In both studies, quetiapine tablets were superior to placebo in increasing the time to recurrence of any mood event. The treatment effect was present for increasing time to recurrence of both manic and depressed episodes. The effect of quetiapine tablets was independent of any specific subgroup (assigned mood stabilizer, sex, age, race, most recent bipolar episode, or rapid cycling course).

### Figure 1 Kaplan-Meier Curves of Time to Recurrence of Mood Event (Study 7)



**Figure 2 Kaplan-Meier Curves of Time to Recurrence of A Mood Event (Study 8)**



**16 HOW SUPPLIED/STORAGE AND HANDLING**

25 mg Tablets pink, round, biconvex, film coated tablets, debossed with “793” on one side and “S” on the other side, are supplied in:

Bottles of 100 tablets (NDC 43547-518-10)

Bottles of 1,000 tablets (NDC 43547-518-11).

50 mg Tablets white, round, biconvex, film coated tablets, debossed with “794” on one side and “S” on the other side, are supplied in:

Bottles of 100 tablets (NDC 43547-519-10)

Bottles of 1,000 tablets (NDC 43547-519-11).

100 mg Tablets yellow, round, biconvex, film coated tablets, debossed with “795” on one side and “S” on the other side, are supplied in:

Bottles of 100 tablets (NDC 43547-520-10)

Bottles of 1,000 tablets (NDC 43547-520-11).

200 mg Tablets white, round, biconvex, film coated tablets, debossed with “796” on one side and “S” on the other side, are supplied in:

Bottles of 100 tablets (NDC 43547-521-10)

Bottles of 1,000 tablets (NDC 43547-521-11).

300 mg Tablets white, capsule-shaped, biconvex, film coated tablets, debossed with “797” on one side and “S” on the other side, are supplied in:

Bottles of 60 tablets (NDC 43547-522-06)

Bottles of 100 tablets (NDC 43547-522-10)

Bottles of 500 tablets (NDC 43547-522-50).

400 mg Tablets yellow, capsule-shaped, biconvex, film coated tablets, debossed with “798” on one side and “S” on the other side, are supplied in:

Bottles of 100 tablets (NDC 43547-523-10)

Bottles of 500 tablets (NDC 43547-523-50).

Store at 20°C-25°C (68°F-77°F); excursions permitted to 15°C-30°C (59°F-86°F) [See USP Controlled Room Temperature].

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Patients should be advised of the following issues and asked to alert their prescriber if these occur while taking quetiapine tablets.

### **Increased Mortality in Elderly Patients with Dementia-Related Psychosis**

Patients and caregivers should be advised that elderly patients with dementia-related psychosis treated with atypical antipsychotic drugs are at increased risk of death compared with placebo. Quetiapine is not approved for elderly patients with dementia-related psychosis [see *Warnings and Precautions* (5.1)].

### **Suicidal Thoughts and Behaviors**

Patients, their families, and their caregivers should be encouraged to be alert to the emergence of anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, mania, other unusual changes in behavior, worsening of depression, and suicidal ideation, especially early during antidepressant treatment and when the dose is adjusted up or down. Families and caregivers of patients should be advised to look for the emergence of such symptoms on a day-to-day basis, since changes may be abrupt. Such symptoms should be reported to the patient's prescriber or health professional, especially if they are severe, abrupt in onset, or were not part of the patient's presenting symptoms. Symptoms such as these may be associated with an increased risk for

suicidal thinking and behavior and indicate a need for very close monitoring and possibly changes in the medication [see *Warnings and Precautions (5.2)*].

#### Neuroleptic Malignant Syndrome (NMS)

Patients should be advised to report to their physician any signs or symptoms that may be related to NMS. These may include muscle stiffness and high fever [see *Warnings and Precautions (5.4)*].

#### Hyperglycemia and Diabetes Mellitus

Patients should be aware of the symptoms of hyperglycemia (high blood sugar) and diabetes mellitus. Patients who are diagnosed with diabetes, those with risk factors for diabetes, or those that develop these symptoms during treatment should have their blood glucose monitored at the beginning of and periodically during treatment [see *Warnings and Precautions (5.5)*].

#### Hyperlipidemia

Patients should be advised that elevations in total cholesterol, LDL-cholesterol and triglycerides and decreases in HDL-cholesterol may occur. Patients should have their lipid profile monitored at the beginning of and periodically during treatment [see *Warnings and Precautions (5.5)*].

#### Weight Gain

Patients should be advised that they may experience weight gain. Patients should have their weight monitored regularly [see *Warnings and Precautions (5.5)*].

#### Orthostatic Hypotension

Patients should be advised of the risk of orthostatic hypotension (symptoms include feeling dizzy or lightheaded upon standing, which may lead to falls), especially during the period of initial dose titration, and also at times of re-initiating treatment or increases in dose [see *Warnings and Precautions (5.7)*].

#### Increased Blood Pressure in Children and Adolescents

Children and adolescent patients should have their blood pressure measured at the beginning of, and periodically during, treatment [see *Warnings and Precautions (5.9)*].

#### Leukopenia/Neutropenia

Patients with a pre-existing low WBC or a history of drug induced leukopenia/neutropenia should be advised that they should have their CBC monitored while taking quetiapine tablets. Patients should be advised to talk to their doctor as soon as possible if they have a fever, flu-like symptoms, sore throat, or any other infection as this could be a result of a very low WBC, which may require quetiapine tablets to be stopped and/or treatment to be given [see *Warnings and Precautions (5.10)*].

#### Interference with Cognitive and Motor Performance

Patients should be advised of the risk of somnolence or sedation (which may lead to falls), especially during the period of initial dose titration. Patients should be cautioned about performing any activity requiring mental alertness, such as operating a motor vehicle (including automobiles) or operating machinery, until they are reasonably certain quetiapine therapy does not affect them adversely [see *Warnings and Precautions (5.16)*].

#### Heat Exposure and Dehydration

Patients should be advised regarding appropriate care in avoiding overheating and dehydration [see *Warnings and Precautions (5.17)*].

#### Concomitant Medication

As with other medications, patients should be advised to notify their physicians if they are taking, or plan to take, any prescription or over-the-counter drugs [see *Drug Interactions (7.1)*].

#### Pregnancy

Advise pregnant women to notify their healthcare provider if they become pregnant or intend to become pregnant during treatment with quetiapine tablets. Advise patients that quetiapine tablets may cause extrapyramidal and/or withdrawal symptoms (agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, and feeding disorder) in a neonate. [see *Use in Specific Populations (8.1)*].

#### Infertility

Advise females of reproductive potential that quetiapine tablets may impair fertility due to an increase in serum prolactin levels. The effects on fertility are reversible [see *Use in Specific Populations (8.3)*].

#### Need for Comprehensive Treatment Program

Quetiapine tablets are indicated as an integral part of a total treatment program for adolescents with schizophrenia and pediatric bipolar disorder that may include other measures (psychological, educational, and social). Effectiveness and safety of quetiapine tablets have not been established in pediatric patients less than 13 years of age for schizophrenia or less than 10 years of age for bipolar mania. Appropriate educational placement is essential and psychosocial intervention is often helpful. The decision to prescribe atypical antipsychotic medication will depend upon the physician's assessment of the chronicity and severity of the patient's symptoms [see *Indications and Usage (1.3)*].

#### **Distributed by:**

Solco Healthcare US, LLC

Somerset, NJ 08873, USA

#### **Manufactured by:**

Zhejiang Huahai Pharmaceutical Co., Ltd.

Xunqiao, Linhai, Zhejiang 317024, China

Revised: 07/2025

203228-04

## **MEDGUIDE SECTION**

### **Medication Guide**

#### **Quetiapine (kwe-TYE-a-peen) Tablets**

Read this Medication Guide before you start taking quetiapine tablets and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

#### **What is the most important information I should know about quetiapine tablets?**

##### **Quetiapine tablets may cause serious side effects, including:**

**1. risk of death in the elderly with dementia. Medicines like quetiapine tablets can increase the risk of death in elderly people who have memory loss (dementia).** Quetiapine tablets are not for treating psychosis in the elderly with dementia.

## 2. risk of suicidal thoughts or actions (antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions).

- **Talk to your or your family member's healthcare provider about:**
- all risks and benefits of treatment with antidepressant medicines.
- all treatment choices for depression or other serious mental illness
- **Antidepressant medications may increase suicidal thoughts or actions in some children, teenagers, and young adults within the first few months of treatment.**
- **Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions.** These include people who have (or have a family history of) depression, bipolar illness (also called manic-depressive illness), or suicidal thoughts or actions.
- **How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?**
- Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed.
- Call the healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.
- Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

### **Call a healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:**

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling very agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

### **What else do I need to know about antidepressant medicines?**

- **Never stop an antidepressant medicine without first talking to your healthcare provider.** Stopping an antidepressant medicine suddenly can cause other symptoms.
- **Antidepressants are medicines used to treat depression and other illnesses.** It is important to discuss all the risks of treating depression, and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.
- **Antidepressant medicines have other side effects.** Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member.
- **Antidepressant medicines can interact with other medicines.** Know all of the medicines that you or your family member take. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking

with your healthcare provider.

- **Not all antidepressant medicines prescribed for children are FDA approved for use in children.** Talk to your child's healthcare provider for more information.

### **What are quetiapine tablets?**

Quetiapine tablets are prescription medicine used to treat:

- schizophrenia in people 13 years of age or older
- bipolar disorder in adults, including:
- depressive episodes associated with bipolar disorder
- manic episodes associated with bipolar I disorder alone or with lithium or divalproex
- long-term treatment of bipolar I disorder with lithium or divalproex
- manic episodes associated with bipolar I disorder in children ages 10-17 years old

It is not known if quetiapine tablets are safe and effective in children under 10 years of age.

### **What should I tell my healthcare provider before taking quetiapine tablets?**

**Before you take quetiapine tablets, tell your healthcare provider if you have or have had:**

- diabetes or high blood sugar in you or your family. Your healthcare provider should check your blood sugar before you start quetiapine tablets, and also during therapy
- high levels of total cholesterol, triglycerides or LDL-cholesterol, or low levels of HDL-cholesterol
- low or high blood pressure
- low white blood cell count
- cataracts
- seizures
- abnormal thyroid tests
- high prolactin levels
- heart problems
- liver problems
- any other medical condition
- pregnancy or plans to become pregnant. It is not known if quetiapine tablets will harm your unborn baby.
- If you become pregnant while receiving quetiapine tablets, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics.
- breast-feeding or plans to breast-feed. Quetiapine can pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you receive quetiapine tablets.
- if you have or have had a condition where you cannot completely empty your bladder (urinary retention), have an enlarged prostate, or constipation, or increased pressure inside your eyes.

**Tell the healthcare provider about all the medicines that you take or recently have taken** including prescription medicines, over-the-counter medicines, herbal supplements, and vitamins.

Quetiapine tablets and other medicines may affect each other causing serious side effects. Quetiapine tablets may affect the way other medicines work, and other medicines may affect how quetiapine tablets work.

Tell your healthcare provider if you are having a urine drug screen because quetiapine tablets may affect your test results. Tell those giving the test that you are taking

quetiapine tablets.

### **How should I take quetiapine tablets?**

- Take quetiapine tablets exactly as your healthcare provider tells you to take it. Do not change the dose yourself.
- Take quetiapine tablets by mouth, with or without food.
- **If you feel you need to stop quetiapine tablets, talk with your healthcare provider first.** If you suddenly stop taking quetiapine tablets, you may have side effects such as trouble sleeping or trouble staying asleep (insomnia), nausea, and vomiting.
- If you miss a dose of quetiapine tablets, take it as soon as you remember. If you are close to your next dose, skip the missed dose. Just take the next dose at your regular time. Do not take 2 doses at the same time unless your healthcare provider tells you to. If you are not sure about your dosing, call your healthcare provider.

### **What should I avoid while taking quetiapine tablets?**

- Do not drive, operate machinery, or do other dangerous activities until you know how quetiapine tablets affects you. Quetiapine tablets may make you drowsy.
- Avoid getting overheated or dehydrated.
- Do not over-exercise.
- In hot weather, stay inside in a cool place if possible.
- Stay out of the sun. Do not wear too much or heavy clothing.
- Drink plenty of water.
- Do not drink alcohol while taking quetiapine tablets. It may make some side effects of quetiapine tablets worse.

### **What are possible side effects of quetiapine tablets?**

#### **Quetiapine tablets can cause serious side effects, including:**

- **See “What is the most important information I should know about quetiapine tablets?”**
- **stroke that can lead to death can happen in elderly people with dementia who take medicines like quetiapine tablets**
- **neuroleptic malignant syndrome (NMS).** NMS is a rare but very serious condition that can happen in people who take antipsychotic medicines, including quetiapine tablets. NMS can cause death and must be treated in a hospital. Call your healthcare provider right away if you become severely ill and have some or all of these symptoms:
  - high fever
  - excessive sweating
  - rigid muscles
  - confusion
  - changes in your breathing, heartbeat, and blood pressure
- **falls** can happen in some people who take quetiapine tablets. These falls may cause serious injuries.
- **high blood sugar (hyperglycemia).** High blood sugar can happen if you have diabetes already or if you have never had diabetes. High blood sugar could lead to:
  - build-up of acid in your blood due to ketones (ketoacidosis)
  - coma
  - death

Increases in blood sugar can happen in some people who take quetiapine tablets. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes (such as being overweight or a family history of diabetes) your healthcare provider should check your blood sugar before you start quetiapine tablets

and during therapy.

Call your healthcare provider if you have any of these symptoms of high blood sugar (hyperglycemia) while taking quetiapine tablets:

- feel very thirsty
- need to urinate more than usual
- feel very hungry
- feel weak or tired
- feel sick to your stomach
- feel confused, or your breath smells fruity
- **high fat levels in your blood (increased cholesterol and triglycerides).** High fat levels may happen in people treated with quetiapine tablets. You may not have any symptoms, so your healthcare provider may decide to check your cholesterol and triglycerides during your treatment with quetiapine tablets.
- **increase in weight (weight gain).** Weight gain is common in people who take quetiapine tablets so you and your healthcare provider should check your weight regularly. Talk to your healthcare provider about ways to control weight gain, such as eating a healthy, balanced diet, and exercising.
- **movements you cannot control in your face, tongue, or other body parts (tardive dyskinesia).** These may be signs of a serious condition. Tardive dyskinesia may not go away, even if you stop taking quetiapine tablets. Tardive dyskinesia may also start after you stop taking quetiapine tablets.
- **decreased blood pressure (orthostatic hypotension),** including lightheadedness or fainting caused by a sudden change in heart rate and blood pressure when rising too quickly from a sitting or lying position.
- **increases in blood pressure in children and teenagers.** Your healthcare provider should check blood pressure in children and adolescents before starting quetiapine tablets and during therapy.
- **low white blood cell count.** Tell your healthcare provider as soon as possible if you have a fever, flu-like symptoms, or any other infection, as this could be a result of a very low white blood cell count. Your healthcare provider may check your white blood cell level to determine if further treatment or other action is needed.
- **cataracts**
- **seizures**
- **abnormal thyroid tests.** Your healthcare provider may do blood tests to check your thyroid hormone level.
- **increases in prolactin levels**
- **sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities**
- **increased body temperature**
- **difficulty swallowing**
- **trouble sleeping or trouble staying asleep (insomnia), nausea, or vomiting if you suddenly stop taking quetiapine tablets.** These symptoms usually get better 1 week after you start having them.

**The most common side effects of quetiapine tablets include:**

**In Adults:**

- drowsiness
- sudden drop in blood pressure upon standing
- weight gain
- sluggishness
- abnormal liver tests
- upset stomach
- dry mouth
- dizziness

- weakness
- abdominal pain
- constipation
- sore throat

### **In Children and Adolescents:**

- drowsiness
- dizziness
- fatigue
- nausea
- increased appetite
- dry mouth
- vomiting
- weight gain
- rapid heart beat

These are not all the possible side effects of quetiapine tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store quetiapine tablets?**

- Store quetiapine tablets at room temperature, between 68°F to 77°F (20°C to 25°C).
- **Keep quetiapine tablets and all medicines out of the reach of children.**

### **General information about the safe and effective use of quetiapine tablets**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use quetiapine tablets for a condition for which it was not prescribed. Do not give quetiapine tablets to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about quetiapine tablets. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about quetiapine tablets that is written for health professionals.

For more information, call Solco Healthcare US, LLC at 1-866-257-2597.

### **What are the ingredients in quetiapine tablets?**

**Active ingredient:** Quetiapine fumarate

**Inactive ingredients:** dibasic calcium phosphate dihydrate, fumaric acid, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium starch glycolate, talc, and titanium dioxide. The 25 mg tablets contain iron oxide red and lecithin (soya). The 50 mg, 200 mg and 300 mg tablets contain lecithin (soya). The 100 mg and 400 mg tablets contain D&C yellow #10 aluminum lake and iron oxide red.

**This Medication Guide has been approved by the U.S. Food and Drug Administration.**

### **Distributed by:**

Solco Healthcare US, LLC  
Somerset, NJ 08873, USA

**Manufactured by:**

Zhejiang Huahai Pharmaceutical Co., Ltd.  
Xunqiao, Linhai, Zhejiang, 317024, China  
Revised: 07/2025

**FOOTNOTES**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**Container Label-25 mg-100 tablets**

Rx only  
NDC 43547-518-10  
Quetiapine Tablets, USP

**PHARMACIST: Dispense the enclosed Medication Guide to each patient.**

**Each film-coated tablet contains:** quetiapine ..... 25 mg  
(equivalent to 28.8 mg quetiapine fumarate)

**USUAL DOSAGE:** See accompanying Prescribing Information.

**WARNING:** As with all medications, keep out of the reach of children.

Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F)  
[see USP Controlled Room Temperature].

**Manufactured by:**

Zhejiang Huahai Pharmaceutical Co., Ltd.  
Xunqiao, Linhai, Zhejiang 317024, China

**Distributed by:**

Solco Healthcare US, LLC  
Somerset, NJ 08873, USA

Rev: 07/2024



**Package/Label Display Panel**

**Container Label-50 mg-100 tablets**

Rx only  
NDC 43547-519-10

Quetiapine Tablets, USP

**PHARMACIST: Dispense the enclosed Medication Guide to each patient.**

**Each film-coated tablet contains:** quetiapine ..... 50 mg

(equivalent to 57.65 mg quetiapine fumarate)

**USUAL DOSAGE:** See accompanying Prescribing Information.

**WARNING:** As with all medications, keep out of the reach of children.

Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F)  
[see USP Controlled Room Temperature].

**Manufactured by:**

Zhejiang Huahai Pharmaceutical Co., Ltd.

Xunqiao, Linhai, Zhejiang 317024, China

**Distributed by:**

Solco Healthcare US, LLC

Somerset, NJ 08873, USA

Rev: 07/2024

## Package/Label Display Panel

### Container Label-100 mg-100 tablets

Rx only

NDC 43547-520-10

Quetiapine Tablets, USP

**PHARMACIST: Dispense the enclosed Medication Guide to each patient.**

**Each film-coated tablet contains:** quetiapine ..... 100 mg

(equivalent to 115.25 mg quetiapine fumarate)

**USUAL DOSAGE:** See accompanying Prescribing Information.

**WARNING:** As with all medications, keep out of the reach of children.

Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F)  
[see USP Controlled Room Temperature].

**Manufactured by:**

Zhejiang Huahai Pharmaceutical Co., Ltd.

Xunqiao, Linhai, Zhejiang 317024, China

**Distributed by:**

Solco Healthcare US, LLC  
Somerset, NJ 08873, USA

Rev: 07/2024



**Package/Label Display Panel**

**Container Label-200 mg-100 tablets**

Rx only

NDC 43547-521-10

Quetiapine Tablets, USP

**PHARMACIST: Dispense the enclosed Medication Guide to each patient.**

**Each film-coated tablet contains:** quetiapine ..... 200 mg

(equivalent to 230.50 mg quetiapine fumarate)

**USUAL DOSAGE:** See accompanying Prescribing Information.

**WARNING:** As with all medications, keep out of the reach of children.

Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F) [see USP Controlled Room Temperature].

**Manufactured by:**

Zhejiang Huahai Pharmaceutical Co., Ltd.  
Xunqiao, Linhai, Zhejiang 317024, China

**Distributed by:**

Solco Healthcare US, LLC  
Somerset, NJ 08873, USA

Rev: 07/2024

	<p>Each film-coated tablet contains: quetiapine ..... 200 mg (equivalent to 230.50 mg quetiapine fumarate)</p> <p><b>USUAL DOSAGE:</b> See accompanying Prescribing Information.</p> <p><b>WARNING:</b> As with all medications, keep out of the reach of children. Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F) [see USP Controlled Room Temperature].</p> <p>Print Medication Guide at: <a href="http://www.solcohealthcare.com/medguide/quetiapine-tablets.pdf">www.solcohealthcare.com/medguide/quetiapine-tablets.pdf</a></p> <p><b>Manufactured by:</b> Zhejiang Huahai Pharmaceutical Co., Ltd. Xunqiao, Linhai, Zhejiang 317024, China</p> <p><b>Distributed by:</b> Solco Healthcare US, LLC Somerset, NJ 08873, USA Rev.: 07/2024</p>	<p>NDC 43547-521-10 <b>Rx only</b></p> <p><b>Quetiapine Tablets, USP</b></p> <p><b>200 mg</b></p> <p>PHARMACIST: Dispense the Medication Guide provided separately to each patient.</p> <p>100 Tablets</p>	<p>This box is FPO to indicate the area to be left uncoated.</p>
--	--	--	--

**Package/Label Display Panel**

**Container Label-300 mg-60 tablets**

Rx only

NDC 43547-522-06

Quetiapine Tablets, USP

**PHARMACIST: Dispense the enclosed Medication Guide to each patient.**

**Each film-coated tablet contains:** quetiapine ..... 300 mg (equivalent to 345.75 mg quetiapine fumarate)

**USUAL DOSAGE:** See accompanying Prescribing Information.

**WARNING:** As with all medications, keep out of the reach of children.

Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F) [see USP Controlled Room Temperature].

**Manufactured by:**

Zhejiang Huahai Pharmaceutical Co., Ltd.

Xunqiao, Linhai, Zhejiang 317024, China

**Distributed by:**

Solco Healthcare US, LLC

Somerset, NJ 08873, USA

Rev: 07/2024

	<p>Each film-coated tablet contains: quetiapine ..... 300 mg (equivalent to 345.75 mg quetiapine fumarate)</p> <p><b>USUAL DOSAGE:</b> See accompanying Prescribing Information.</p> <p><b>WARNING:</b> As with all medications, keep out of the reach of children. Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F) [see USP Controlled Room Temperature].</p> <p>Print Medication Guide at: <a href="http://www.solcohealthcare.com/medguide/quetiapine-tablets.pdf">www.solcohealthcare.com/medguide/quetiapine-tablets.pdf</a></p> <p><b>Manufactured by:</b> Zhejiang Huahai Pharmaceutical Co., Ltd. Xunqiao, Linhai, Zhejiang 317024, China</p> <p><b>Distributed by:</b> Solco Healthcare US, LLC Somerset, NJ 08873, USA Rev.: 07/2024</p>	<p>NDC 43547-522-06 <b>Rx only</b></p> <p><b>Quetiapine Tablets, USP</b></p> <p><b>300 mg</b></p> <p>PHARMACIST: Dispense the Medication Guide provided separately to each patient.</p> <p>60 Tablets</p>	<p>This box is FPO to indicate the area to be left uncoated.</p>
---	--	---	--

**Package/Label Display Panel**

## Container Label-400 mg-100 tablets

Rx only

NDC 43547-523-10

Quetiapine Tablets, USP

**PHARMACIST: Dispense the enclosed Medication Guide to each patient.**

**Each film-coated tablet contains:** quetiapine ..... 400 mg

(equivalent to 461.00 mg quetiapine fumarate)

**USUAL DOSAGE:** See accompanying Prescribing Information.

**WARNING:** As with all medications, keep out of the reach of children.

Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F)  
[see USP Controlled Room Temperature].

### Manufactured by:

Zhejiang Huahai Pharmaceutical Co., Ltd.

Xunqiao, Linhai, Zhejiang 317024, China

### Distributed by:

Solco Healthcare US, LLC

Somerset, NJ 08873, USA

Rev: 07/2024

Each film-coated tablet contains: quetiapine ..... 400 mg  
(equivalent to 461.00 mg quetiapine fumarate)  
**USUAL DOSAGE:** See accompanying Prescribing Information.  
**WARNING:** As with all medications, keep out of the reach of children.  
Store at 20°C - 25°C (68°F - 77°F); excursions permitted to 15°C - 30°C (59°F - 86°F) [see USP Controlled Room Temperature].  
Print Medication Guide at:  
[www.solcohealthcare.com/medguide/quetiapine-tablets.pdf](http://www.solcohealthcare.com/medguide/quetiapine-tablets.pdf)  
**Manufactured by:**  
Zhejiang Huahai Pharmaceutical Co., Ltd.  
Xunqiao, Linhai, Zhejiang 317024, China  
**Distributed by:**  
Solco Healthcare US, LLC  
Somerset, NJ 08873, USA  
Rev.: 07/2024

NDC 43547-523-10 Rx only

**Quetiapine  
Tablets, USP**

**400 mg**

PHARMACIST: Dispense the Medication Guide  
provided separately to each patient.

**100 Tablets**

This box is FPO to indicate  
the area to be left uncoated.

## QUETIAPINE

quetiapine tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:43547-518
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
QUETIAPINE FUMARATE (UNII: 2S3PL1B6UJ) (QUETIAPINE - UNII:BGL0JSY5SI)	QUETIAPINE	25 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FUMARIC ACID</b> (UNII: 88XHZ13131)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	793;S
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43547-518-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	
2	NDC:43547-518-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206954	04/12/2022	

## QUETIAPINE

quetiapine tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:43547-519
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>QUETIAPINE FUMARATE</b> (UNII: 2S3PL1B6UJ) (QUETIAPINE - UNII:BGL0JSY5SI)	QUETIAPINE	50 mg

Inactive Ingredients	
Ingredient Name	Strength
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: 07TSZ97GEP)	
FUMARIC ACID (UNII: 88XHZ13131)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	WHITE	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	794;S
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43547-519-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	
2	NDC:43547-519-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206954	04/12/2022	

## QUETIAPINE

quetiapine tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:43547-520
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
QUETIAPINE FUMARATE (UNII: 2S3PL1B6UJ) (QUETIAPINE - UNII:BGL0JSY5SI)	QUETIAPINE	100 mg

Inactive Ingredients	
Ingredient Name	Strength

<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FUMARIC ACID</b> (UNII: 88XHZ13131)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	YELLOW	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	795;S
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43547-520-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	
2	NDC:43547-520-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206954	04/12/2022	

## QUETIAPINE

quetiapine tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:43547-521
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>QUETIAPINE FUMARATE</b> (UNII: 2S3PL1B6UJ) (QUETIAPINE - UNII:BGL0JSY5SI)	QUETIAPINE	200 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>FUMARIC ACID</b> (UNII: 88XHZ13131)	

<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	796;S
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43547-521-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	
2	NDC:43547-521-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206954	04/12/2022	

## QUETIAPINE

quetiapine tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:43547-522
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>QUETIAPINE FUMARATE</b> (UNII: 2S3PL1B6UJ) (QUETIAPINE - UNII:BGL0JSY5SI)	QUETIAPINE	300 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>FUMARIC ACID</b> (UNII: 88XHZ13131)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	797;S
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43547-522-06	60 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	
2	NDC:43547-522-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	
3	NDC:43547-522-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206954	04/12/2022	

## QUETIAPINE

quetiapine tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:43547-523
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>QUETIAPINE FUMARATE</b> (UNII: 2S3PL1B6UJ) (QUETIAPINE - UNII:BGL0JSY5SI)	QUETIAPINE	400 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: 07TSZ97GEP)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FUMARIC ACID</b> (UNII: 88XHZ13131)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	YELLOW	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	798;S
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43547-523-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	
2	NDC:43547-523-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	04/12/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206954	04/12/2022	

**Labeler** - Solco Healthcare US, LLC (828343017)

**Registrant** - Prinston Pharmaceutical Inc. (967289799)

### Establishment

Name	Address	ID/FEI	Business Operations
Zhejiang Huahai Pharmaceutical Co., LTD		530732460	MANUFACTURE(43547-518, 43547-519, 43547-520, 43547-521, 43547-522, 43547-523)

Revised: 7/2025

Solco Healthcare US, LLC