

MINOXIDIL FOR MEN- minoxidil aerosol, foam

BosleyMD

Minoxidil (for Men)

Drug Facts

Active ingredient

Minoxidil 5% w/w (without propellant)

Purpose

Hair regrowth treatment for men

Use

to regrow hair on the top of the scalp (vertex only, see pictures on side of this carton)

Warnings

For external use only. For use by men only.

Extremely Flammable: Avoid fire, flame, or smoking during and immediately following application.

Do not use if

- you are a woman
- your amount of hair loss is different than that shown on side of this carton or your hair loss is on the front of the scalp. Minoxidil Topical Aerosol, 5% (Foam) is not intended for frontal baldness or receding hairline.
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

Ask a doctor before use if you have

heart disease.

When using this product

- do not apply on other parts of the body
- avoid contact with the eyes. In case of accidental contact, rinse eyes with large

amounts of cool tap water.

- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. Results may occur at 2 months with twice a day usage. For some men, you may need to use this product for at least 4 months before you see results.
- the amount of hair regrowth is different for each person. This product will not work for all men.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- apply half a capful 2 times a day to the scalp in the hair loss area
- massage into scalp with fingers, then wash hands well
- see enclosed leaflet for complete directions on how to use
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- hair growth has been shown in a clinical study of men (mostly white) aged 18-49 years who used it for 4 months
- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- store at controlled room temperature 20° to 25°C (68° to 77°F)
- contents under pressure. Do not puncture or incinerate container.
- Do not expose to heat or store at temperatures above 120°F (49°C).

Inactive ingredients

butane, butylated hydroxytoluene, cetyl alcohol, citric acid, glycerin, isobutane, lactic acid, polysorbate 60, propane, purified water, SD alcohol 40-B, stearyl alcohol

Questions?

Call **1-800-BOSLEY1 (1-800-267-5391)**

Principal display panel

Minoxidil Topical Aerosol 5% (For Men)

Hair Regrowth Treatment

Easy-to Use Foam For Men

Not For Use by Women

Reactivates Hair Follicles

Unscented

Clinically Proven To Help Regrow Hair

Minoxidil Topical Aerosol, 5% is for men who have a general thinning of hair on the top of the scalp (vertex only, as shown below). Not intended for frontal baldness or a receding hairline. Minoxidil Topical Aerosol, 5% (For Men) has been shown to regrow hair in men with the following degrees of thinning hair or hair loss.

For best results:

Use twice daily.

Apply directly to the scalp.

Read and keep carton and information leaflet for full product information.

Distributed by: BosleyMD

800.BOSLEY1 bosleymd.com

Manufactured for: Scientific Hair Research LLC,

Los Angeles, CA 90045

Package label

MINOXIDIL FOR MEN

minoxidil aerosol, foam

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69113-501
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)	MINOXIDIL	50 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ALCOHOL (UNII: 3K9958V90M)	
ISOBUTANE (UNII: BXR49TP611)	
LACTIC ACID, UNSPECIFIED FORM (UNII: 33X04XA5AT)	
BUTANE (UNII: 6LV4FOR43R)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
PROPANE (UNII: T75W9911L6)	
WATER (UNII: 059QF0KO0R)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69113-501-01	1 in 1 PACKAGE	08/28/2025	
1		60 g in 1 CAN; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA208092	08/28/2025	
------	------------	------------	--

Labeler - BosleyMD (961897647)

Revised: 8/2025

BosleyMD