

MOTION SICKNESS RELIEF- meclizine hcl tablet
Chain Drug Consortium, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

pv 778

Active ingredient (in each tablet)

Meclizine HCl 25mg

Purpose

Antiemetic

Uses

For the prevention and treatment of these symptoms associated with motion sickness • nausea • vomiting
• dizziness

Warnings

Do not use in children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- breathing problems, such as emphysema or chronic bronchitis
- glaucoma
- difficulty urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, take the first dose one hour before starting activity
- not for frequent or prolonged use except on the advice of a doctor
- do not exceed recommended dose

Adults and children 12 years if age and older	Take 1-2 tablets once daily, or as directed by a doctor. Do not exceed 2 tablets in 24 hours.
---	--

Other information

- **TAMPER EVIDENT:** Do not use if imprinted seal under cap is broken or missing.
- store at room temperature 15°-30°C (59°-86°F)
- protect from humidity

Inactive ingredients

croscarmellose sodium, D and C yellow no. 10 lake, lactose, magnesium stearate, microcrystalline cellulose, silica, sodium starch glycolate, starch, stearic acid

package label

Drug Facts (continued)

Other information

- Tamper Evident: Do not use if imprinted seal under cap is broken or missing.
- store at room temperature 15°-30° C (59°-86° F) • protect from heat and humidity

Inactive ingredients

croscarmellose sodium, D&C yellow #10 lake, lactose, magnesium stearate, microcrystalline cellulose, silica, sodium starch glycolate, starch, stearic acid


*This product is not manufactured or distributed by the owner of the registered trademark DRAMAMINE II®.

Distributed By: Chain Drug Consortium, LLC.
3301 N.W. Boca Raton Blvd., Suite 101
Boca Raton, FL 33431

REV778-0711CT



If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.



Drug Facts

SAVE CARTON FOR COMPLETE DRUG FACTS

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

For the prevention and treatment of these symptoms associated with motion sickness
• nausea • vomiting • dizziness

Warnings

Do not use in children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- breathing problems such as emphysema or chronic bronchitis
- glaucoma • difficulty urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

Drug Facts (continued)

When using this product

- drowsiness may occur • avoid alcoholic drinks
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • to prevent motion sickness, take the first dose one hour before starting activity

- not for frequent or prolonged use except on the advice of a doctor
- do not exceed recommended dose

age	dosage
adults and children 12 years of age and older	Take 1 to 2 tablets once daily, or as directed by a doctor. Do not exceed 2 tablets in 24 hours.



MOTION SICKNESS RELIEF

meclizine hcl tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:68016-778

Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg	
Inactive Ingredients				
	Ingredient Name	Strength		
	ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
	SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
	CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
	STARCH, CORN (UNII: O8232NY3SJ)			
	MAGNESIUM STEARATE (UNII: 70097M6I30)			
	D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
	SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
	CROSCARMELOSE SODIUM (UNII: M28OL1HH48)			
	STEARIC ACID (UNII: 4ELV7Z65AP)			
Product Characteristics				
Color	yellow	Score	no score	
Shape	ROUND	Size	9mm	
Flavor		Imprint Code	TCL086	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-778-02	1 in 1 CARTON	07/01/2011	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part336	07/01/2011		

Labeler - Chain Drug Consortium, LLC (101668460)

Registrant - Geri-Care Pharmaceutical Corp (611196254)