

PAIN RELIEVER- acetaminophen tablet
Shield Pharmaceuticals Corp

Pain Reliever

Drug Facts

Active Ingredient

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours with other drugs
- containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if youhave liver disease

Ask a doctor or pharmacist before use if you aretaking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

store between 20-25°C (68-77°F)

Inactive ingredients

magnesium stearate, potassium sorbate, povidone (K30), pregelatinized starch, purified water, sodium starch glycolate

Questions or comments?

1-800-373-6981

Compare to the active ingredient in Tylenol® Extra Strength

Pain Reliever

Acetaminophen 500 mg

Med365

Compare to the active ingredient in Extra Strength Tylenol®

Extra Strength

Pain Reliever

Acetaminophen 500 mg

Pain reliever / Fever reducer

50 caplets

Not actual Size

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet) Purpose
Acetaminophen 500 mg Pain reliever/fever reducer

Uses
■ temporarily relieves minor aches and pains due to:
■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Drug facts (continued under the label)

Distributed by:
Shield Pharmaceuticals Corp.
Ronkonkoma, NY 11779

6 1141940633 0

LOT:
EXP:

Made in China
ORG: 1025

Drug Facts (continued)

Ask a doctor before use if you have ■ liver disease
Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin
Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not use more than directed (see overdose warning)
adults and children
12 years and over:
■ take 2 caplets every 6 hours
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
■ do not use for more than 10 days unless directed by a doctor
■ ask a doctor
children under 12 years:

Other information
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients
magnesium stearate, potassium sorbate, povidone (K30), pregelatinized starch, purified water, sodium starch glycolate

Questions or comments?
1-800-373-6861 (toll-free)

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc. Model Consumer Healthcare Division, owner of the registered trademark Tylenol® Extra Strength caplets.

PAIN RELIEVER

acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83059-0103
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POTASSIUM SORBATE (UNII: 1VPU26JZ Z 4)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	
WATER (UNII: 059QF0KO0R)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	16mm
Flavor		Imprint Code	PAR500
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83059-0103-5	50 in 1 BOTTLE; Type 0: Not a Combination Product	11/07/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/07/2025	

Labeler - Shield Pharmaceuticals Corp (118724924)

Revised: 11/2025

Shield Pharmaceuticals Corp