

ASPIRIN LOW DOSE- aspirin tablet, coated Bryant Ranch Prepack

gc 981

Active ingredient (in each tablet)

Aspirin 81 mg(NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains
- ask your doctor about other uses for aspirin

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer.
- if you ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you are taking a diuretic
- you have a history of stomach problems, such as heartburn
- you have: -high blood pressure -heart disease -liver cirrhosis -kidney disease -asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding: -feel faint -vomit blood -have bloody or black stools -have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- any new symptoms occur
- ringing in the ears or loss of hearing occurs

These can be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dose
- drink a full glass of water with each dose
- swallow whole, do not chew or crush
- **adults and children 12 years and older:** take 4-8 tablets every 4 hours, as needed, not more than 48 tablets in 24 hours, or as directed by a doctor
- **children under 12:** ask a doctor

Other information

- store at 20-25°C (68-77°F); excursions permitted between 15°C - 30°C (59°F - 86°F)

Inactive ingredients

cellulose, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, PEG, polydextrose, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, starch, talc, titanium dioxide, triacetin, triethyl citrate, wax.

Questions or comments?

1-800-540-3765

HOW SUPPLIED

Aspirin 81 mg EC Tablets

- NDC: 71335-0622-1: 120 Tablets in a BOTTLE
- NDC: 71335-0622-2: 30 Tablets in a BOTTLE
- NDC: 71335-0622-3: 100 Tablets in a BOTTLE
- NDC: 71335-0622-4: 20 Tablets in a BOTTLE
- NDC: 71335-0622-5: 90 Tablets in a BOTTLE
- NDC: 71335-0622-6: 60 Tablets in a BOTTLE
- NDC: 71335-0622-7: 36 Tablets in a BOTTLE
- NDC: 71335-0622-8: 15 Tablets in a BOTTLE
- NDC: 71335-0622-9: 10 Tablets in a BOTTLE

Repackaged/Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504

Aspirin 81 mg EC Tablets



GTIN 0037335062217
Lot 20682026
Exp 7/24/2026
SN 0123456789

Drug Facts	
Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID) ...	Pain reliever
<small>Nonsteroidal anti-inflammatory drug</small>	
Uses	
Temporarily relieves minor aches and pains. Because of its delayed action, this product will not produce fast relief of headaches or other symptoms needing immediate relief. Ask your doctor about other uses for enteric-coated 81 mg Aspirin.	
Warnings	
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be early sign of Reye's syndrome, a rare but serious illness. Allergy Alert: Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, shock, asthma (wheezing). Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for longer time than directed. Do not use, if you are allergic to aspirin or any other pain reliever/fever reducer. Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis; making any other drug under a doctor's care for any serious condition. Stop use and ask a doctor if: an allergic reaction occurs. Seek medical help right away if you experience any of the following signs of stomach bleeding: feel faint, have bloody or black stools, vomit blood, have stomach pain that does not get better, spine gets worse or lasts more than 10 days; redness or swelling is present; new symptoms occur: ringing in the ears or a loss of hearing occurs. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Other information	
Store at 20°C (77°F); excursions permitted between 15°-30°C (59°-86°F) above or below room temperature on package.	
Directions	
• Drink a full glass of water with each dose. • Adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor. • Children under 12 years: consult a doctor.	
Inactive ingredients	
Corn starch, pregelatinized starch, povidone, microcrystalline cellulose, colloidal silicon dioxide, stearic acid, methacrylic acid and ethyl acrylate copolymer, talc, titanium dioxide, triethyl citrate, colloidal silicon dioxide, sodium bicarbonate, sodium lauryl sulfate, D&C yellow #10, hypromellose, bisacetin.	

NDC 71335-0622-1

Aspirin USP

81 mg

120 Tablets



Repackaged by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504 USA

Manufactured by:
Geri-Care
Pharmaceutical Corp.



7133506221

ASPIRIN LOW DOSE

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71335-0622(NDC:57896-981)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71335-0622-1	120 in 1 BOTTLE; Type 0: Not a Combination Product	03/28/2018	
2	NDC:71335-0622-2	30 in 1 BOTTLE; Type 0: Not a Combination Product	02/09/2018	
3	NDC:71335-0622-3	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/11/2018	
4	NDC:71335-0622-4	20 in 1 BOTTLE; Type 0: Not a Combination Product	11/21/2018	
5	NDC:71335-0622-5	90 in 1 BOTTLE; Type 0: Not a Combination Product	02/09/2018	
6	NDC:71335-0622-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	08/22/2018	
7	NDC:71335-0622-7	36 in 1 BOTTLE; Type 0: Not a Combination Product	04/27/2018	
8	NDC:71335-0622-8	15 in 1 BOTTLE; Type 0: Not a Combination Product	08/12/2022	
9	NDC:71335-0622-9	10 in 1 BOTTLE; Type 0: Not a Combination Product	07/09/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2000	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(71335-0622) , RELABEL(71335-0622)

Revised: 7/2024

Bryant Ranch Prepack