

**MUCUS RELIEF COLD, FLU AND SORE THROAT- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl solution
TOPCO ASSOCIATES LLC**

TopCare 44-005

Active ingredients (in each 20 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg
Guaifenesin 400 mg
Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sinus congestion and pressure
 - stuffy nose
 - cough
 - headache
 - nasal congestion
 - sore throat
- temporarily promotes nasal and/or sinus drainage
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash

- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- liver disease
- thyroid disease
- heart disease
- difficulty in urination due to enlargement of the prostate gland
- high blood pressure
- diabetes
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- **each 20 mL contains:** sodium 10 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments?

1-888-423-0139

Principal Display Panel

+TopCare®
health

NDC 76162-050-45

COMPARE TO THE ACTIVE INGREDIENTS IN MAXIMUM STRENGTH
MUCINEX® FAST-MAX® COLD, FLU & SORE THROAT

MAXIMUM STRENGTH

Mucus Relief

Cold, Flu & Sore Throat

ACETAMINOPHEN

PAIN RELIEVER • FEVER REDUCER

DEXTROMETHORPHAN HBr • COUGH SUPPRESSANT

GUAIFENESIN • EXPECTORANT

PHENYLEPHRINE HCl • NASAL DECONGESTANT

MULTI-SYMPTOM RELIEF:

Relieves Body Pain, Headache,

Fever & Sore Throat

- Controls Cough
- Relieves Nasal & Chest Congestion
- Thins & Loosens Mucus
- Relieves Sinus Pressure & Congestion

FOR AGES 12+
6 FL OZ (177 mL)
Mixed Berry
Flavored

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ITASCA, IL 60143

©TOPCO LNKA0525 QUESTIONS? 1-888-423-0139

topcare@topco.com www.topcarebrand.com

Visit here for more information: <http://topbrnds.com/4912jr>

QUALITY GUARANTEED

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat. 50844 REV0724A00545



NDC 76162-050-45

*COMPARE TO THE ACTIVE INGREDIENTS IN MAXIMUM STRENGTH MUCINEX® FAST-MAX® COLD, FLU & SORE THROAT

MAXIMUM STRENGTH

Mucus Relief Cold, Flu & Sore Throat

ACETAMINOPHEN

PAIN RELIEVER • FEVER REDUCER
DEXTROMETHORPHAN HBr • COUGH SUPPRESSANT
GUAIFENESIN • EXPECTORANT
PHENYLEPHRINE HCl • NASAL DECONGESTANT

MULTI-SYMPTOM RELIEF:

- Relieves Body Pain, Headache, Fever & Sore Throat
- Controls Cough
- Relieves Nasal & Chest Congestion
- Thins & Loosens Mucus
- Relieves Sinus Pressure & Congestion

FOR AGES 12+



QUALITY GUARANTEED

F-005-45
REV A

6 FL OZ (177 mL)

Mixed Berry Flavored

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ITASCA, IL 60143
©TOPCO LNK00525 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com
Visit here for more information: <http://topbrnds.com/4912jr>

PARENTS:

Learn about your medicine at www.StopForGood.com

PEEL BACK TAB TO READ COMPLETE DRUG FACTS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms: ■ headache ■ stuffy nose ■ cough ■ minor aches and pains ■ sore throat ■ sinus congestion and pressure ■ nasal congestion ■ temporarily promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ 3 or more alcoholic drinks every day while using this product ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® FAST-MAX® Cold, Flu & Sore Throat. 50844 REV0724A00545



0 36800 38616 7 B-005-45 REV A

No Print / No Varnish Area Lot # and Exp. Info

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ skin reddening ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- liver disease ■ diabetes
- heart disease ■ thyroid disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur

Drug Facts (continued)

- pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- each 20 mL contains: sodium 10 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments? 1-888-423-0139

HINGE

HINGE

Topcare 44-005

MUCUS RELIEF COLD, FLU AND SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:76162-050

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	BERRY (MIXED)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-050-45	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/11/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/11/2023	

Labeler - TOPCO ASSOCIATES LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(76162-050) , pack(76162-050)

Revised: 3/2026

TOPCO ASSOCIATES LLC