

**ALLERGY CHILDRENS- diphenhydramine hydrochloride solution**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Children's Allergy Liquid Drug Facts**

**Active ingredient (in each 5 mL)**

Diphenhydramine HCl 12.5 mg

**Purpose**

Antihistamine

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - sneezing
  - itching of the nose or throat
  - runny nose
  - itchy, watery eyes

**Warnings**

**Do not use**

- with any other product containing diphenhydramine, even one used on skin
- to make a child sleepy

**Ask a doctor before use if the child has**

- a breathing problem such as chronic bronchitis
- glaucoma
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if the child is**

taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

**Keep out of reach of children.**

## Directions

- find right dose on chart below
- mL = milliliter
- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 2 years	do not use
children 2 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL to 10 mL

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

## Other information

- **each 5 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F). Protect from light. Store in outer carton until contents used.
- **do not use if printed neckband is broken or missing**

## Inactive ingredients

anhydrous citric acid, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, poloxamer 407, purified water, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution

## Questions or comments?

**1-800-719-9260**

## Principal Display Panel

meijer®

CHILDREN'S | ages 6-11

allergy liquid

Antihistamine | Diphenhydramine HCl 12.5 mg/5 mL Oral Solution

COMPARE TO CHILDREN'S BENADRYL® ACTIVE INGREDIENT

4-6 HOUR RELIEF

RELIEF FOR SNEEZING; RUNNY NOSE; ITCHY THROAT; ITCHY, WATERY EYES

CHERRY FLAVOR

Multi-Symptom  
 Alcohol Free  
 8 FL OZ (237 mL)



**meijer** NDC 41250-379-34  
**CHILDREN'S | ages 6-11**  
**allergy liquid**  
 Antihistamine | Diphenhydramine  
 HCl 12.5 mg/5 mL Oral Solution

COMPARE TO CHILDREN'S BENADRYL®  
 ACTIVE INGREDIENT\*

**4-6 HOUR RELIEF** RELIEF FOR SNEEZING; RUNNY NOSE;  
 ITCHY THROAT; ITCHY, WATERY EYES

**CHERRY FLAVOR**

Multi-Symptom  
 Alcohol Free



8 FL OZ (237 mL)

**meijer** NDC 41250-379-34  
**CHILDREN'S | ages 6-11**  
**allergy liquid**  
 Antihistamine | Diphenhydramine  
 HCl 12.5 mg/5 mL Oral Solution

COMPARE TO CHILDREN'S BENADRYL®  
 ACTIVE INGREDIENT\*

**4-6 HOUR RELIEF** RELIEF FOR SNEEZING; RUNNY NOSE;  
 ITCHY THROAT; ITCHY, WATERY EYES

**CHERRY FLAVOR**

Multi-Symptom  
 Alcohol Free



8 FL OZ (237 mL)

**meijer**  
**CHILDREN'S | ages 6-11**  
**allergy liquid**  
 Antihistamine | Diphenhydramine  
 HCl 12.5 mg/5 mL Oral Solution  
 RELIEF FOR SNEEZING; RUNNY NOSE;  
 ITCHY THROAT; ITCHY, WATERY EYES  
**4-6 HOUR RELIEF**  
 CHERRY FLAVOR

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
Diphenhydramine HCl 12.5 mg	Antihistamine

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - sneezing
  - itching of the nose or throat
  - runny nose
  - itchy, watery eyes

**Warnings**

Do not use

- with any other product containing diphenhydramine, even one used on skin
- to make a child sleepy

Ask a doctor before use if the child has

- a breathing problem such as chronic bronchitis
- glaucoma
- a sodium-restricted diet

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers

**When using this product**

- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- find right dose on chart below
- mL = milliliter
- take every 4 to 6 hours, or as directed by a doctor
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 2 years	do not use
children 2 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL to 10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information**

- each 5 mL contains sodium 15 mg
- store at 20-25°C (68-77°F). Protect from light. Store in outer carton until contents used.
- do not use if printed neckband is broken or missing

**Inactive ingredients**

anhydrous citric acid, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, poloxamer 407, purified water, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution

**Drug Facts (continued)**  
 Questions or comments? 1-800-719-9260  
 \*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Benadryl®.



DIST. BY MEIJER DISTRIBUTION, INC.  
 GRAND RAPIDS, MI 49544  
 www.meijer.com



37934 6E C11

**ALLERGY CHILDRENS**  
 diphenhydramine hydrochloride solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-379
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII: 8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 5 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB65)	
<b>POLOXAMER 407</b> (UNII: TUF2IVW3M2)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	

## Product Characteristics

<b>Color</b>	RED (Bluish-Red)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-379-26	1 in 1 CARTON	11/15/1987	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-379-34	1 in 1 CARTON	12/12/1995	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	11/15/1987	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 10/2025

Meijer Distribution Inc