EQUATE EFFERVESCENT COLD RELIEF- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent Walmart

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Equate Efferves cent Cold Relief

Active ingredients

Aspirin 325 mg (NSAID)* Chlorpheniramine maleate 2 mg Phenylephrine Bitartrate 7.8 mg *Nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer Antihistamine Nasal decongestant

Uses

temporarily relieves these symptoms of the common cold:

- fever sneezing nasal congestion sore throat headache minor aches and pains
- runny nose sinus congestion and pressure

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic.

Ask a doctor before use if you have ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

■ taking sedatives or tranquilizers ■ presently taking a prescription drug ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

- do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.
- do not use more than directed you may get drowsy avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding
- feel faint vomit blood have bloody or black stools have stomach pain that does not get better

■ an allergic reaction occurs. Seek medical help right away. ■ pain or nasal congestion gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present
- ringing in the ears or a loss of hearing occurs nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause

problems in the unborn child or complications during delivery.

Keep out of reach of children. Keep out of reach of children.

In case of overdose,

get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4-6 hours.
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

each tablet contains: sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Questions or comments?

1 - 888 - 287 - 1915

PACKAGE LABEL. PRINCIPAL DISPLAY PANEL



Aspirin (**NSAID**)/Pain Reliever - Fever Reducer Chlorpheniramine Maleate/Natihistamine Phenylephrine Bitartrate/Nasal Decongestant

Effervescent Cold Relief



EQUATE EFFERVESCENT COLD RELIEF aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent **Product Information** HUMAN OTC DRUG **Product Type** NDC:49035-768 Item Code (Source) **Route of Administration** ORAL **Active Ingredient/Active Moiety Ingredient Name** Strength **Basis of Strength** ASPIRIN (UNII: R16C05Y76E) (ASPIRIN - UNII:R16C05Y76E) ASPIRIN 325 mg CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE -CHLORPHENIRAMINE 2 mg UNII:3U6IO1965U) MALEATE

Ingredient Name				Strength
ACESULFAME POTAS	SIUM (UNII: 230V73Q5G9)		
ASPARTAME (UNII: Z01	H242BBR1)			
ANHYDRO US CITRIC	ACID (UNII: XF417D3PSL)			
DOCUSATE SODIUM (UNII: F05Q2T2JA0)			
MANNITOL (UNII: 30W	/L53L36A)			
POVIDONE K30 (UNII:	U725QWY32X)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SODIUM BICARBONA	TE (UNII: 8 MDF5V39QO)			
Product Character	ristics			
Color	white	Score	Score	
Shape	ROUND	Size	Size	
Flavor	LEMON	Imprint Code	Imprint Code	
Contains				
Packaging				
# Item Code	Package Description		Marketing Start Date	Marketing End Date
1 NDC:49035-768-67	48 in 1 CARTON		0 5/23/20 19	
1	2 in 1 POUCH; Type 0: Not a Combination Product			
Maultating I-fa	umation			
Marketing Info				
Marketing Category	Application Number	or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341			

Labeler - Walmart (051957769)

Revised: 5/2019

Walmart